

Journal Pre-proof

Deep-sea ecosystems of the Indian Ocean >1000 m

Elin A. Thomas, Todd Bond, Jess L. Kolbusz, Yakufu Niyazi,
Denise J.B. Swanborn, Alan J. Jamieson



PII: S0048-9697(24)06951-1

DOI: <https://doi.org/10.1016/j.scitotenv.2024.176794>

Reference: STOTEN 176794

To appear in: *Science of the Total Environment*

Received date: 7 May 2024

Revised date: 21 September 2024

Accepted date: 5 October 2024

Please cite this article as: E.A. Thomas, T. Bond, J.L. Kolbusz, et al., Deep-sea ecosystems of the Indian Ocean >1000 m, *Science of the Total Environment* (2024), <https://doi.org/10.1016/j.scitotenv.2024.176794>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2024 Published by Elsevier B.V.

Deep-sea ecosystems of the Indian Ocean > 1,000 m

Elin A. Thomas¹, Todd Bond¹, Jess L. Kolbusz¹, Yakufu Niyazi¹, Denise J.B. Swanborn¹, Alan J. Jamieson¹

¹Minderoo-UWA Deep-Sea Research Centre, School of Biological Sciences and Oceans Institute, The University of Western Australia, 35 Stirling Highway, Perth, Australia, WA 6009

Corresponding author: Elin A. Thomas elin.thomas@uwa.edu.au

Keywords: Indian Ocean, deep sea, benthic ecology, seafloor geomorphology, knowledge gaps

Abstract

The Indian Ocean is the third largest of the world's oceans, accounting for ~20% for the global marine realm. It is geomorphologically complex, hosting a wide variety of ecosystems across basins, trenches, seamounts, ridges, and fracture zones. While modern exploration has contributed significantly to our knowledge of its coastal ecosystems, deeper waters (> 1,000 m) remain relatively unknown despite accounting for over 90% of its total area. This study provides the first comprehensive review of the Indian Ocean's diverse deep sea, presenting ecosystem knowledge summaries for each major seafloor feature, contextualised with the broader historical, socioeconomic, geological, and oceanographic conditions. Unsurprisingly, some ecosystems are better characterised than others, from the relatively well-surveyed Java (Sunda) Trench and hydrothermal vents of the Carlsberg, Central and Southwest Indian Ridges, to the unexplored Southeast Indian Ridge and hadal features of the western Indian Ocean. Similarly, there is a large depth discrepancy in available records with a clear bias towards shallower sampling. We identify four outstanding problems to be addressed for the advancement of deep-sea research in the Indian Ocean: 1) inconsistencies in research extent and effort over spatial scales, 2) severe lack of data over temporal scales, 3) unexplored deep pelagic environments, and 4) a need to place the Indian Ocean's deep-sea ecosystems in a global

context. By synthesising and championing existing research, identifying knowledge gaps, and presenting the outstanding problems to be addressed, this review provides a platform to ensure this forgotten ocean is prioritised for deep-sea research during the UN Ocean Decade and beyond.

Introduction

The Indian Ocean spans an area of 71,729,338 km², of which ~92% is deep sea > 1,000 m (Table 1). It is the third largest of the world's oceans and is second only to the Pacific in its proportion of deep-sea area. Despite this, it is the least understood of the oceans, having been largely neglected scientifically until the late-1950s (Rogers 2017). Modern exploration has greatly contributed to our knowledge of coastal biodiversity, yet its deeper waters (> 1,000 m) are still relatively unknown (Demopoulos et al. 2003, Wafar et al. 2011). Earlier reviews of deep Indian Ocean research have primarily focussed on the benthic communities of continental margins, slopes, and abyssal plains (Neyman et al. 1973, Demopoulos et al. 2003, Ingole and Koslow 2005). Ramiro-Sánchez et al. (2023) further highlighted the data paucity for deep-sea habitats in the southern Indian Ocean, with many studies limited to the northwest and central regions (e.g. Wafar et al. 2011, Saraswat et al. 2022). As a result, the Indian Ocean is lacking a definitive baseline for deep-sea biodiversity research to inform future exploration and achieve the goals of the United Nations (UN) Decade of Ocean Science for Sustainable Development (2021-2030).

Here we provide a review of the deep-sea ecosystems of the Indian Ocean greater than 1,000 m, contextualised with geological and oceanographic conditions that collectively form this 20% of the global ocean. We first provide background on the historical, socioeconomic, geological, oceanographic, and broad-scale biodiversity trends related to deep Indian Ocean research, then we present in-depth knowledge summaries for each major seafloor feature in terms of deep-sea ecosystems. Through this, we will identify knowledge gaps and present the outstanding problems that should be addressed for the advancement of deep-sea research in the Indian

Ocean. A map of all major Indian Ocean seafloor features is presented in Appendix 1 as a companion guide for this review. The aim is to provide a platform for future studies, to ensure this forgotten ocean is prioritised for deep-sea research during the Ocean Decade and beyond.

Geographical boundaries and data sources

The geographic extent of the Indian Ocean is defined here following Harris et al. (2014), modifying the boundaries set by 'The Limits of Oceans and Seas' (International Hydrography Organisation 1953) to include bounding marginal seas and exclude Southern Ocean waters south of 60°S. The Indian Ocean is bordered to the north by the northern limits of the Red and Arabian Seas, and the Bay of Bengal; to the east by the Andaman Sea and Java Trench, the easterly limits of the Timor and Arafura Seas, and down the 146°55' meridian from the southern point of Tasmania; to the south by the 60°S parallel of the Southern Ocean; and to the west by the 20°E meridian running from Cape Agulhas, South Africa. The Persian Gulf to the north of the Indian Ocean was excluded from this review as its maximum depth is < 100 m. Names of seafloor features were verified following the IHO-IOC GEBCO Gazetteer of Under Sea Feature Names (GEBCO). Where features are not listed by GEBCO, names were aligned with the US BGN Advisory Committee on Undersea Features (ACUF).

Table 1: Area (km²) and relative proportion (%) of different depth ranges in the Indian Ocean, calculated using the Mollweide map projection and GEBCO_2023 bathymetry data in ArcPro (v3.0). EEZ: Exclusive Economic Zone, ABNJ: Area Beyond National Jurisdiction.

	Area (km ²)		
	Indian Ocean	EEZs	ABNJ
Total area	70,884,251	24,023,171	46,861,080
Deep sea (> 1,000 m)	65,023,066 (91.73%)	18,275,565 (76.07%)	46,747,501 (99.76%)
Bathyal (1,000 – 3,000 m)	10,060,277 (14.19%)	6,343,043 (26.40%)	3,717,234 (7.93%)
Abyssal (3,000 – 6,000 m)	54,683,224 (77.14%)	11,842,225 (49.30%)	42,840,999 (91.42%)
Hadal (6,000 m +)	279,565 (0.39%)	90,297 (0.38%)	189,268 (0.40%)

Maps were produced in QGIS (v 3.32.2) using GEBCO_2023 bathymetric data and shapefiles downloaded from the following online repositories: shapefiles for the extent of the Indian Ocean and the national jurisdiction boundaries (EEZs) of Indian Ocean countries from [marineregions.org](https://marineregions.org/downloads.php) (Flanders Marine Institute 2023, <https://marineregions.org/downloads.php>, accessed 05/10/2023); marine protected areas from [protectedplanet.net](https://www.protectedplanet.net/en/thematic-areas/marine-protected-areas) (<https://www.protectedplanet.net/en/thematic-areas/marine-protected-areas>, accessed 01/11/2023); and deep-sea mining contracts for areas beyond national jurisdiction from the International Seabed Authority webpage (<https://www.isa.org.jm/exploration-contracts/maps/>, accessed 11/10/2023). Bathymetric contour layers were used to generate polygons of areas exceeding 6,000 m depth to determine the area and maximum depths of Indian Ocean hadal features, excluding polygons $< 2 \text{ km}^2$ for quality control. Note that these estimations are subject to variation based on the use of different map projections for GIS computations and are regularly updated with the availability of new, higher quality data (Stewart and Jamieson et al. 2018). Spatial analysis to determine the total area and proportion of different depth ranges across the Indian Ocean was conducted in ArcPro (v 3.0) using the Mollweide projection.

To study broad-scale sampling effort and biodiversity patterns in the Indian Ocean, data were downloaded from the open-access Ocean Biodiversity Information System (OBIS) mapper (<https://mapper.obis.org/>, accessed 15/11/2023). OBIS is the most complete and comprehensive open-access data repository on the biogeography of marine taxa, with access to 123 million records from over 5,000 datasets (<https://obis.org/>; Vanden Berghe et al. 2010). Data were filtered for records $> 1,000 \text{ m}$ depth and for the following areas: Arabian Sea, Red Sea, Bay of Bengal, Andaman Sea, Arafura Sea, Timor Sea, Great Australian Bight, Laccadive Sea, Gulf of Aden, Mozambique Channel, Persian Gulf, Gulf of Oman, Indian Ocean, ABNJ: Indian Ocean, Gulf of Aqaba, and Gulf of Suez. The resulting 65,604 records were imported to R Studio (v 2023.09.1) for data manipulation and visualisation. Data were cleaned to remove non-metazoans, blank records, and to check for erroneous depth records by removing those with

depths recorded < 1,000 or > 8,000 m (deeper than the maximum depth of the Indian Ocean) and bathymetry recorded < 1,000 m, leaving 32,257 records. Where data were plotted by taxonomic group, phyla with low numbers of records (< 300) were grouped as 'Other'. All figures were edited using Inkscape (v 1.3.2).

These OBIS data have been instrumental in identifying spatial, bathymetric, and ecosystem biases in marine exploration (Webb et al. 2010, Kennedy and Rotjan 2023). There are, however, significant discrepancies between the open-access OBIS records and available metadata for research cruises, as shown for the Central and South Atlantic (Bridges et al. 2023). This suggests that not all datasets collected during research cruises have been uploaded to OBIS.

Nevertheless, in the absence of data completeness, patterns can be derived and interpreted from the available data.

Historical background

The first scientific study of the Indian Ocean began with the 19th century European expeditions. The *HMS Challenger Expedition* (1872-1876) contributed relatively little to scientific knowledge of the deepwater ecosystems of the Indian Ocean as it surveyed circumpolar waters south of the polar front. The later German *Valdivia Expedition* (Deutsche Tiefsee-Expedition) (1898-1899) covered a large part of the Indian Ocean and sampled deep-sea benthic fauna from depths up to 2,490 m. The first half of the 20th century saw more European mounted expeditions surveying the Indian Ocean and providing some insights in deep-sea biodiversity, including the fourth Danish *Dana Expedition* (1928-1930), the Anglo-Egyptian *John Murray Expedition* (1930s) which yielded some samples of deep-sea bivalves (Knudsen 1967), and the Swedish *Albatross Expedition* (1947-1948), which collected deep benthic samples and sediment cores. The Danish *Galathea* (1950-1952) provided trawl samples from lower abyssal and hadal fauna in the west and central Indian Ocean. Trawling data from Soviet-led expeditions on the *R/V Vityaz* added to knowledge on bathyal and abyssal organisms (Bezrunov 1963).

Deep-sea sampling on these early cruises was relatively opportunistic and the International Indian Ocean Expedition (IIOE 1960-65) was the first comprehensive international effort dedicated to improving knowledge on the Indian Ocean including its deep ecosystems, with ship time provided by 40 vessels from over 20 nations (Zeitzschel and Gerlach 2012). Subsequently, and among others, the Royal Society Indian Ocean Deep Slope Fishing Expedition (1969) explored deep-slope resources in the Indian Ocean (Forster et al. 1970), Soviet/Ukrainian research institutes including YugNIRO led deep-sea fisheries explorations on plateaus, seamounts and ridges in the Indian Ocean (1970s-1980s; Budnichenko et al. 1997, Romanov 2003, Parin et al. 2008), French expeditions to the Indian Ocean involved some deep-sea studies in the 1970s (e.g. BENTHEDI Expedition, Thomassin 1977, Poupin 2015), and the Netherlands Indian Ocean Expeditions explored deep-sea environments in the northern Arabian Sea and western Indian Ocean (1992-1993; van Weering et al. 1997). Roberts et al. (2020) provide a more detailed summary of historical fisheries and seamount exploration in the Indian Ocean. The Second International Indian Ocean Expedition (IIOE-2) (Hood et al. 2016) was initiated approximately 60 years after the conclusion of IIOE with a running time between 2015-2025.

Further international drilling project collaborations including the Deep Sea Drilling Project (1968-1983) and IODP (2013-2023) frequented the Indian Ocean deep sea, but were mainly geologically focussed. On the ecological side, the Census of Marine Life (CoML) had five deep-sea projects which were largely focused on the Atlantic but included Indian Ocean research on abyssal plains (CeDAMar – Stuart et al. 2008, Ebbe et al. 2010), seamounts (CenSeam – Consalvey et al. 2010, Stocks et al. 2012), chemosynthetic environments (ChEss – Baker et al. 2010, German et al. 2011), continental margins (COMARGE – Menot et al. 2010) and mid-ocean ridges (MAR-ECO – Bergstad et al. 2008).

Of the contemporary research cruises in the Indian Ocean, those targeting depths > 1,000m remain the minority. However, important contributions to knowledge on Indian Ocean deep-sea

biodiversity has come from cruises on national and private research vessels (e.g. Ingle and Koslow 2005, Rogers and Taylor 2011, Trotter et al. 2021, Zhou et al. 2022, Bond et al. 2023).

Socioeconomics and management

Approximately one-third of the Indian Ocean lies within the Exclusive Economic Zones (EEZs) of 36 countries, 76% of which is deep sea greater than 1,000 m (Table 1, Figure 1). Thirty of these countries have deep sea > 1,000 m within their EEZ, 21 have abyssal depths > 3,000 m, and only Indonesia and Australia have hadal depths > 6,000 m (Table 2, Bond and Jamieson 2022). The governance of deep-sea environments within these EEZ is the responsibility of the countries that hold national jurisdiction over these areas (Table 2). However, access to assets such as vessels, research equipment, and funds for deep-sea exploration and research is low for Indian Ocean countries, despite the broad view that deep-sea environments are important across the region (Bell et al. 2022, Talma et al. 2022). Collaborations with the deep-water oil and gas industry have occasionally provided a platform for deep-sea exploration for countries with limited capacity (Gates et al. 2017a).

In Tanzania, ecological data collected as part of the SERPENT project (Scientific and Environmental ROV Partnership using Existing iNdustry Technology) provided the basis for recommendations on the sustainable use of deep-sea resources and marine spatial management (Gates et al. 2021). Such partnerships have also facilitated the discovery of new biodiversity and interesting ecological observations, highlighting the importance of deep-sea exploration in these areas (Moore and Gates 2015, Gates 2016, Gates et al. 2017b, Macreadie et al. 2018). Meanwhile, India has been working to expand its deep-sea research as part of its Deep Ocean Mission to explore and exploit the deep sea both within its EEZ and beyond (Atmanand 2019). National collaborations with independent research institutes also increasingly provide a window into deeper waters and support national research agendas for EEZs, though biological observations predominantly remain limited to depths < 1000 m (e.g. Monaco Explorations Indian Ocean Mission 2022, Nekton Maldives expedition 2022-2023, Ahusan et al. 2023, OceanX Seychelles

expedition 2024). However, given the narrow continental shelves of many countries bordering the Indian Ocean, deep sea > 1,000 m is part of many EEZs. Partnerships, knowledge exchange, and capacity building support a comprehensive understanding of deep-sea habitats and biodiversity within EEZs and achieve national priorities as well as the aims of international agreements (Woodall et al. 2021).

Areas beyond national jurisdiction (ABNJ) comprise the other two-thirds of the Indian Ocean, almost 100% of which is deep sea > 1,000 m (Table 1). This part of the seafloor, known as the 'Area', is governed by the International Seabed Authority (ISA) which was established under the United Nations' Convention on the Law of the Sea (UNCLOS) in 1994 (Thompson et al. 2018). Some factors impeding research effort for Indian Ocean ABNJ include the greater distance from shore, requiring significant funding and access to larger vessels, and a lack of incentive for research without direct national benefits. It is for these reasons that the southern Indian Ocean remains one of the least explored regions in the world (Ramiro-Sánchez et al. 2023). Yet the Area is also the target for emerging industries such as deep-sea mining. The ISA have granted four mining exploration contracts for polymetallic sulfides along the southwest, central, and southeast Indian ridges, as well as one contract and one reserved area for polymetallic nodules in the Mid-Indian Ocean Basin (ISA 2022, Figure 1). Two of these mining exploration contracts are held by the Government of India, yet the other three are held by countries that do not hold any national jurisdiction in the Indian Ocean, and no other ISA deep-sea mining contracts are held by Indian Ocean countries. Conversely, no established mining contracts exist for cobalt-rich ferromanganese crusts (CFC) in the Indian Ocean.

Marine conservation and sustainable management of marine resources is essential across the deep-sea environments of both EEZs and ABNJ. In EEZs, several marine protected areas (MPAs) have been designated in the Indian Ocean, the majority of which include depths > 1,000 m (Figure 1). However, these MPAs offer varying levels of protection from anthropogenic threats such as fishing and mining (Grorud-Colvert et al. 2021) and are not necessarily effective

conservation measures for deep-sea environments without full consideration of potential impacts (Thomas et al. 2021) and management capacity. The recent finalisation and adoption of Biodiversity Beyond National Jurisdiction (BBNJ), a treaty which also sits within the framework of UNCLOS, was a promising step towards the conservation and sustainable management of marine biodiversity in ABNJ, including deep-sea environments (Mendenhall et al. 2023). However, the BBNJ treaty now requires ratification and adoption by the 83 member states who have signed the agreement to date, including only eight Indian Ocean countries (UN 2023, Table 2), thus its effective implementation is not yet guaranteed.

The Indian Ocean is of high interest to commercial fisheries and fishing operations, some of which extend into waters > 1,000 m (Georgeson et al. 2020), though mainly concentrated on the ridges and seamounts in the southern part of the Indian Ocean (Clark et al. 2007). Most commercial deep-water fisheries have historically developed in these ABNJ without stock size and fishing effort assessments (Rogers et al. 2017), though some have been supplemented by research surveys and scientific monitoring (Romanov 2003, Marsac et al. 2020). Regional Fisheries Management Organizations (RFMOs) provide frameworks to manage fishing activity, including in ABNJ. The Southern Indian Ocean Fisheries Agreement (SIOFA) is an RFMO with 13 member entities and is responsible for the management of deep-water fishing activity in the Southern Indian Ocean including in ABNJs (Shotton 2006, SIOFA Secretariat 2023). The SIOFA has established 5 interim benthic protected areas (BPAs) - areas closed to bottom or midwater fishing due to their unique biodiversity value - and is actively undertaking research into the establishment of new protected areas (Ramiro-Sánchez et al. 2023, SIOFA 2023). In addition, the Southern Indian Ocean Deep-Water Fisheries Operators Association (SIODFA), an industry-led body representing the interests of deep-sea fishing operators and an observer to SIOFA, has delimited 11 voluntary BPAs (Shotton 2006, SIODFA Secretariat 2016).




Figure 1: Map of the Indian Ocean with exclusive economic zone (EEZ) boundaries of neighbouring countries (white lines), SIOFA management area (blue dashed lines), established marine protected areas (MPAs, green polygons), and International Seabed Authority (ISA) exploratory mining contracts (red polygons) and reserved areas (orange polygons). The blue area defines the limits of the Indian Ocean.

Table 2: Countries and their territories bordering the Indian Ocean and the depth classifications of their Exclusive Economic Zones. Bathyal: 1,000-3,000 m, abyssal: 3,000-6,000m, hadal: > 6,000 m.

Countries and territories	Bathyal	Abyssal	Hadal
Australia°			
- Christmas Island	X	X	X
- Cocos (Keeling) Island			
- Heard and McDonald Islands			
Bahrain#	-	-	-
Bangladesh°	X	-	-
Chagos Archipelago*	X	X	
Comoros	X	X	-
Democratic Republic of Timor-Leste°	X	X	-
Djibouti	X	-	-
Egypt#	X	-	-
Eritrea#	X	-	-
Federal Republic of Somalia	X	X	-
France°			
- French Southern and Antarctic Lands*	X	X	-
- Mayotte			
- Réunion			
India			
- Andaman and Nicobar	X	X	-
Indonesia°	X	X	X
Iran	X	X	-
Iraq#	-	-	-
Israel#	-	-	-
Jordan#	-	-	-
Kenya	X	X	-
Kuwait#	-	-	-
Madagascar	X	X	-
Maldives	X	X	-
Mozambique	X	X	-
Myanmar	X	X	-
Oman	X	X	-
Pakistan	X	X	-
Papua New Guinea	-	-	-
Qatar#	-	-	-
Republic of Mauritius°	X	X	-
Saudi Arabia#	X	-	-
Seychelles°	X	X	-
South Africa			
- Prince Edward Islands	X	X	-
Sri Lanka	X	X	-
Sudan#	X	-	-
Tanzania°	X	X	-
Thailand	X	-	-
United Arab Emirates#	X	-	-
Yemen	X	X	-

*Disputed territory, #Red Sea or Persian Gulf only, °Signatories of BBNJ Treaty

Note that the French Southern and Antarctic Lands comprise the Crozet, Kerguelen, Saint Paul and Amsterdam, and Scattered Islands, and Adélie Land.

Geological history

The initial formation of the Indian Ocean is linked to the break-up of the Gondwana super-continent when Africa–South America (West Gondwana) separated from Madagascar–Greater India–Antarctica–Australia (East Gondwana) during the Middle Jurassic (~ 175 Ma) (Figure 2; Schettino and Scotese 2005, Müller et al. 2019). As India-Madagascar continued to rotate south from Africa, the early rifting between the eastern and western Gondwana landmasses had developed into seafloor spreading within the Mozambique and Somali basin systems and the Davie Fracture Zone acted as the major transform fault to separate these two basins. Therefore, these regions recorded the oldest ocean floor in the Indian Ocean (Davis et al. 2016). Seafloor spreading between Africa and Gondwanaland continued until the Early Cretaceous at ~120 Ma and the landmass continued to be broken into blocks of various sizes (Figure 2; Gibbons et al. 2013). As a result, the Eastern Gondwana fragmented into the India-Madagascar-Sri Lanka-Seychelles dominated plate in the northwest and the unified Australian–Antarctic Plate in the southeast. Then, Greater India began separating from Australia’s western margin, and subsequently unzipped from Antarctica at around ~ 120 Ma. This northward motion and anticlockwise rotation of Greater India created the Naturaliste, Wallaby–Zenith, and Exmouth plateaus and associated fracture zones during the drift. In the mid-Cretaceous period, there was increasing magmatic activity in the region, related to the development of the Kerguelen Large Igneous Province from ~ 120 to 110 Ma, and formed the Kerguelen Plateau and the Broken Ridge (Figure 2; Frey et al. 2000). The continued northward movement of the Indian plate caused the separation of Madagascar from the Indian continent during the Late Cretaceous (~90–85 Ma) and Neotethys became narrower. At the same time, there was a significant plate reorganization and an increase in the spreading rates along the Central and Southeast Indian Ridges (Scotese et al. 1988). Rifting was also initiated in the southwestern boundary between the Australia and Antarctica plates.

The first collision of the Indian continent with the islands of the south Asian subduction zones occurred, and therefore the northward movement of the Indian continent slowed down in the Early Eocene at ~53 Ma (Figure 2; Scotese 2021). Meanwhile, relatively slow seafloor spreading continued and separated Australia from Antarctica, marking the formation of the Indian, Australian, and Antarctic continents in the east Indian Ocean. The relative movement of the Indian and Antarctic plates continues along the transform fault, which is expressed as a deep trough located east of the Ninetyeast Ridge (Levchenko et al. 2021). During the Mid-Eocene (~40 M) there was the second phase of plate reorganization due to the hard collision of the Indian and Eurasian plates. As the relative movement of the Indian and Australian plates along the transform boundary ceased, they merged into the single Indo-Australian Plate. When the Australian plate began to move rapidly northward during the late Eocene (40–35 Ma), it separated from the Antarctic Plate, and allowed the deep waters from the Indian Ocean to mix with the south Pacific Ocean for the first time through the Tasman Strait (Figure 2; Hodel et al. 2021).




Figure 2: Oblique view of the oceanic crustal age of the Indian Ocean region since 200 Ma to present day, showing the break-up of Eastern Gondwana continent and opening of the Indian Ocean. Images created using gPlates software (<https://www.earthbyte.org/category/resources/software-workflows/pygplates/>). Base maps are from Müller et al. 2019.

Geomorphology

The ocean can be delimited into different depth zones: the bathyal zone (1,000-3,000 m), the abyssal zone (3,000-6,000 m) and the hadal zone (> 6,000 m). Table 1 summarises the proportion of the Indian Ocean that reaches these depth zones, with over 77% lying at abyssal depths. On the contrary, hadal depths represent < 1% of the total Indian Ocean area (0.35%: Harris et al. 2014, 0.39%: this study, Table 1). Excluding polar regions, which exhibit much lower levels of seafloor roughness, this is the lowest proportion of oceanic basin classified as hadal zone, globally (Harris et al. 2014, Jamieson 2015). The geomorphological diversity of the Indian Ocean basin is important from an ecological and biological perspective, as different seafloor features are commonly associated with particular habitats (Harris and Baker 2020).

The Indian Ocean is geomorphologically complex: it comprises 20 major seas and basins intersected by several active ridge systems (Appendix 1, Figure 3); most notably, the Central Indian Ridge (CIR), Southwest Indian Ridge (SWIR) and Southeast Indian Ridge (SEIR), which meet at the central point of the Rodriguez Triple Junction (RTJ) at ~25.5°S, 70°E (Munsch and Schlich 1988), dividing the western and eastern Indian Ocean. The topographic complexity of the western Indian Ocean is encapsulated by its diverse range of tectonic boundaries and mid-ocean ridge/fracture zone configurations (Parson and Evans 2005). The CIR extends northwards from the RTJ, punctuated by several fracture zones including the Flinders, Mary Celeste, Argo, Vema Trench, and Vityaz Fracture Zones (Fisher et al. 1971). The Carlsberg Ridge runs northwest from the CIR to the Owen Fracture Zone (Tao et al. 2013), forming a barrier between the Arabian Sea and the Somali Basin (Figure 3, Appendix 1). The Arabian Sea hosts the Indus Fan which extends 1,500 km into the Indian Ocean from the Indus Canyon at the Pakistan continental shelf break (Clift et al. 2001). Into the Gulf of Aden, the East Sheba Ridge and West Sheba Ridge are offset by the Alula-Fartak Trough (Fournier et al. 2010), with the Red Sea to the northeast.

To the west of the CIR lies the Mascarene Basin, bounded to the east by the Mascarene Plateau, and Madagascar Basin, bounded by the Madagascar Plateau to the west and SWIR to the south (Figure 3, Appendix 1). The Mozambique Basin is situated south of the Mozambique Channel, with its associated homonymous plateau and escarpment to the west. The ultraslow spreading SWIR extends southwest from the RTJ intersected by numerous fracture zones including the Melville, Discovery II, and Andrew Bain Fracture Zones (Sauter and Cannat 2010). The Elcano Rise extends west-east between the SWIR and the Crozet Islands, with the Crozet Basin lying to the northeast, bounded by the SWIR, SEIR, RTJ, and the Kerguelen Plateau to the south (Figure 3, Appendix 1). At the western boundary of the Indian Ocean, the SWIR divides the Agulhas Basin to the north (with its homonymous plateau) from the Weddell Basin to the south, both of which extend beyond the limits of the Indian Ocean, into the Atlantic and Southern Oceans, respectively.

East of the CIR, the aseismic Chagos-Laccadive Ridge divides the Arabian Sea from the Mid-Indian Ocean Basin, running ~2,000 km south from the western continental margin of India to ~9°S adjacent to the Vema Trench (Figure 3, Appendix 1; Parson and Evans 2005). To the northeast, the largest submarine fan globally, the Bengal Fan, extends 3,000 km from the Swatch of No Ground Canyon (ACUF) at its head, along the length of the Bay of Bengal to the northern limits of the Mid-Indian Ocean Basin (Curry et al. 2003). East of this, beyond the Andaman-Nicobar Ridge, lies the marginal Andaman Sea, also known as the Burma Sea. In the Cocos Basin, the smaller Nicobar Fan extends southwards from between the Andaman-Nicobar Ridge and Ninetyeast Ridge (Pickering et al. 2019). The aseismic Ninetyeast Ridge intersects the east Indian Ocean, running almost meridionally for more than 4,500 km from ~7°N, 90.4°E to ~34°S, 87°E and dividing the Mid-Indian Ocean Basin from the Wharton Basin to the east (Figure 3, Appendix 1; Sclater and Fisher 1974). At ~15°S, the Osborn Plateau juts out from the western side of the Ninetyeast Ridge, while the Broken Ridge extends to the east at 32.25°S. The eastern Indian Ocean basins are generally much deeper than western Indian Ocean (Figure 3). The Wharton Basin, for example, is intersected by several north-south directional fracture zones

including the so-called Investigator Ridge, resulting in a large number of discrete depressions within the basin reaching hadal depths ($> 6,000$ m; Weston et al. 2021, Jamieson et al. *in review*).

Extending the length of Indonesia's continental margin, the Java, or Sunda, Trench is the only subduction trench in the Indian Ocean, formed by the subduction of the Indo-Australian Plate beneath the Eurasian Plate (Appendix 1; Jamieson et al. 2022). Its deepest point, and the deepest point in the Indian Ocean, is located in its eastern sector, south of Bali at 11.129°S , 114.942°E (Bongiovanni et al. 2022). At its eastern end lies the North Australian Basin and associated Argo Abyssal Plain, and the Timor Sea and associated Timor Trough (Wei et al. 2023), with the Arafura Sea denoting the eastern limits of the Indian Ocean north of Australia. Southeast of the Wharton Basin is a geomorphologically complex area of seafloor encompassing the Cuvier and Zenith Plateaus, Cuvier Abyssal Plain, Wallaby-Cuvier Escarpment, Lost Dutchmen and East Indiaman Ridges, Brouwer Trough, and the Wallaby-Zenith Fracture Zone (Appendix 1; Weston et al. 2021, Bond et al. 2023). The abyssal Perth Basin is bounded by the above features to the northwest and by the Naturaliste Plateau and Diamantina Fracture Zone to the south. The Diamantina Fracture Zone runs along the southern aspect of Broken Ridge, past Cape Leeuwin to the South Australian Basin, with its deepest point located at Dordrecht Deep (Bongiovanni et al. 2022). The South Australian Basin is enclosed to the north by the Great Australian Bight and separated by the SEIR from the more southerly Australian-Antarctic Basin, which extends from the Kerguelen Plateau to the southeastern boundary of the Indian Ocean (Figure 3, Appendix 1). The SEIR has fewer named fracture zones than the SWIR and CIR – its most notable are the Geelvinck, Nieuw Amsterdam and Zeewolf Fracture Zones (Ma and Cochran 1996). The SEIR is however interrupted by a region of anomalous geophysical and geochemical properties between 120° and 128° E, named the Australian-Antarctic Discordance (Sempéré et al. 1991).




Figure 3: Bathymetric map of the Indian Ocean with major basins and seas labelled. Note that the Persian Gulf to the north is not included in this study as it is shallower than 1,000 m.

Oceanographic setting

The Indian Ocean is characterised by unique oceanographic conditions with a profound influence on deep-sea habitats (Demopoulos et al. 2003). From a physical oceanographic perspective, the Indian Ocean has distinct features in the upper ocean layers, setting it apart from other oceans (Phillips et al. 2021). The primary flow regimes in the upper layers are monsoonal-forced circulation of the tropics and northern hemisphere alongside the subtropical gyre of the south Indian Ocean (Talley et al. 2011). In the deep and bottom waters, the paths of bifurcation are controlled by the complex topography such as deep ridges and mixing with overlying waters.

From a global circulation perspective, the Indian Ocean functions as an upwelling region, much like the Pacific Ocean, and has a greater impact on global climate variability than historically

thought (Schott et al. 2009). Over the past decade, higher resolution observations encapsulating broader spatial coverage have improved numeric simulations and provided a clearer picture of the oceanographic processes of the Indian Ocean (Phillips et al. 2021). Additionally, programs such as the second Indian Ocean Expedition (IIOE-2, <https://iioe-2.incois.gov.in/IIOE-2/index.jsp>), the Global Ocean Observing System for Indian Ocean (IOGOOS, <https://incois.gov.in/iogoos/home.jsp>) and the Global Ocean Ship-Based Hydrographic Investigations Program (GO-SHIP, <https://www.go-ship.org/index.html>) have improved our physical and biogeochemical understanding of this poorly understood basin (Hood et al. 2022).

There are major modes of variability which influence the upper layers of the equatorial Indian Ocean on seasonal and interannual timescales. The Madden-Julian Oscillation (MJO) drives thermodynamic responses in the upper Indian Ocean through atmospheric variability causing wind, surface heat flux and rainfall anomalies (Madden and Julian 1971, Hendon and Salby 1994, McPhaden and Foltz 2013). The Indian Ocean Dipole (IOD) can be categorized as a positive or negative event, altering the monsoon season, and influencing global climate (Cai et al. 2011; Ihara et al. 2008). When positive, there is a cold sea surface temperature (SST) anomaly and anomalous upwelling in the southeastern tropical Indian Ocean (Saij et al. 1999, Webster et al. 1999, Horii et al. 2013). When negative, there is a warm SST anomaly in the southeastern tropical Indian Ocean and a cold SST anomaly in the central-western Indian Ocean (Horii et al. 2013). Wyrтки jets are semi-annually occurring narrow currents that flow eastward along the equator during inter-monsoonal periods (Wyrтки 1973, Schott and McCreary 2001, Nagura and McPhaden 2010). Their structure causes a thermocline depression on the eastern side of the basin and uplift in the western basin (Wyrтки 1973, Wiggert et al. 2006). Combined, these physical mechanisms impact biogeochemical processes at basin-wide scales in the upper Indian Ocean by impacting primary production and chlorophyll maximum depths, altering biological productivity, and therefore impacting higher trophic levels (Wiggert et al. 2009, Prasanna Kumar et al. 2014, Strutton et al. 2015). These changes in the biological productivity

of the upper Indian Ocean in turn affect the influx of particles to the deep sea and as such the food availability at the deep seafloor (McCreary et al. 2013, Hood et al. 2017).

There is no high-latitude ocean cooling due to the Asian continental border at 30°N, driving monsoon dynamics to play a key role in shaping ocean circulation. Strong and seasonally reversing monsoon winds (northeast in boreal summer and southwest in boreal winter), drive many changes in the Indian Ocean (Vinayachandran et al. 2018). For example, reversal of western boundary currents (Zang et al. 2021), enhanced biological productivity (Thushara and Vinayachandran 2016), and changes in the mixed layer depth (Kara et al. 2003). The Indian Ocean connects to the Pacific Ocean through the Indonesian Archipelago, forming the Indonesian Throughflow. Excess heat is transferred from the western Pacific Ocean to the eastern Indian Ocean (Schneider 1998, De Deckker 2016), creating a north-south density gradient that drives eastward surface currents between Australia and Madagascar and generates a uniquely southward-flowing eastern boundary current along Western Australia, the Leeuwin Current (Feng et al. 2015).

Off the continental shelf, complex topographic ridges separate the Indian Ocean into basins (Figure 3, Appendix 1), generating intense mixing at the basin boundaries, contributing to internal wave energy generation (Scott et al. 2011, Huussen et al. 2012), and ultimately supporting meridional overturning circulation (Sloyan 2006). The overturning of water mass in the Indian Ocean is driven by the lack of a northern deep-water source, resulting in net northward inflow in the cool deep water from the south, and outflow in the upper ocean, evident in the water mass properties at depth (Figure 4). Circumpolar Deep Water (CDW) fills into the greater deep Indian Ocean as deep western boundary currents in the Crozet Basin, Central Indian Ocean Basin, and Perth Basin (Toole and Warren 1993). North Atlantic Deep Water (NADW) enters the south-western Indian Ocean from the South Atlantic, driven by the topographic of the Madagascar Ridge and Mozambique Channel (Toole and Warren 1993). Principle flows below 3,500 m are northward along the deep channels leading to the Arabian

Sea and the Bay of Bengal. In the Cape and Agulhas basins, there are isolated cyclonic flows around 4,500 m (Reid 2003). These water mass movements enable upwelling of the lower meridional overturning circulation (MOC) from abyssal waters through diffusive mixing and export the largest poleward heat flux of all Southern Hemisphere basins, contributing to global climate variability (Schott et al. 2009, Roxy et al. 2014).




Figure 4: Deep and bottom water properties of the Indian Ocean from the World Ocean Circulation Experiment Indian Ocean Atlas (Talley 2013). (a) Potential Temperature (°C) and (b) Practical Salinity at 2,500 m, (c) Potential Temperature (°C) and (d) Practical Salinity at 4,000 m.

Broad-scale biodiversity patterns

Mapping the available, cleaned OBIS data for metazoan occurrence records > 1,000 m in the Indian Ocean (32,257 records) highlights the regions that have been well versus poorly sampled (Figure 5). There is a clear pattern of greater sampling effort along continental margins, particularly around India, within the Bay of Bengal, and south of Australia (Figure 5a). There are also clusters of records for ABNJ that are of particular interest and have been targeted by research cruises such as in the Mid-Indian Ocean Basin, around the Crozet Plateau, at the Madagascar Plateau, and along the Southwest Indian Ridge (see subsequent knowledge summaries for abyssal plains and seamounts). Equally, there are large areas of the Indian Ocean that are seemingly void of sampling, especially in the southeast, however, this may simply be a symptom of collected data not being uploaded to OBIS. It is possible to see specific sampling transects from the data (Figure 5b), for example the majority of Chaetognatha records appear to follow straight lines from north to south that can be attributed to the sampling transects of the first International Indian Ocean Expedition (IIOE). Similarly, clusters of annelid, arthropod, echinoderm records in the Mid-Indian Ocean Basin can be attributed to India's research program to explore the basin's polymetallic nodule field (see IndOBIS database, <https://indobis.in/>). Arthropods make up a large proportion of the available records, especially in the northern Indian Ocean and Red Sea, while a high number of chordates have been recorded off southeast Australia, primarily by Australia's Commonwealth Scientific and Industrial Research Organisation (CSIRO) (Figure 5b). These records off southern Australia can be attributed to the Great Australian Bight Research and Deepwater Marine programs, comprising six research cruises to the area between 2013 and 2017 (see subsequent summary for continental margins).

There is a clear bias towards shallower sampling, with the vast majority (> 24,000) of deep Indian Ocean metazoan records from bathyal depths between 1,000 and 3,000 m (Figure 6), compared to almost 8,000 records from abyssal depths and only 66 records from hadal depths.

Furthermore, there are no Porifera records from > 6,000 m, nor hadal records for other phyla such as Ctenophora (Figure 6), despite the known existence of sponge and ctenophore species at these depths (Jamieson et al. 2022, Jamieson et al. 2023; see subsequent summary for subduction trenches). While this may be due to the lack of open-access data, the decrease in sampling effort with increasing depth is a well-documented trend, likely owing to the technical challenges of sampling at abyssal and hadal depths (Jamieson et al. 2013, Bridges et al. 2023).




Figure 5 (previous page): Heat maps of metazoan biodiversity records, for all phyla (A) and across different phyla (B), in the deep Indian Ocean (> 1,000 m) using data from the Ocean Biodiversity Information System (OBIS). Other category includes phyla with low record numbers (< 300): Brachiopoda, Bryozoa, Ctenophora, Hemichordata, Kinorhynca, Nematoda, Nemertea, Platyhelminthes, Priapulida, and Tardigrada.




Figure 6: Metazoan biodiversity records across different depth ranges in the Indian Ocean using data from the Ocean Biodiversity Information System (OBIS). Depth ranges – bathyal: 1,000-3,000 m, abyssal: 3,000-6,000 m, hadal: > 6,000 m. Other category includes phyla with low record numbers (< 300): Brachiopoda, Bryozoa, Ctenophora, Hemichordata, Kinorhynca, Nematoda, Nemertea, Platyhelminthes, Priapulida, and Tardigrada.

Knowledge summaries by seafloor feature

Continental margins

Beyond the continental shelf, the ocean deepens to bathyal depths along high gradient slopes, the size and steepness of which vary along the continental margins of the Indian Ocean. These margins are geomorphologically complex, encompassing a variety of heterogenous deep-sea habitats, including canyons, oxygen minimum zones, fans, cold seeps, and cold-water coral reefs (Menot et al. 2010, Levin and Sibuet 2012). The continental shelf is generally much narrower along the east coast of Africa compared to the Australian and northern continental margins; the Tanzanian shelf (200 m contour), for example, extends less than 10 km from shore in places (Gates et al. 2021). The mean Indian Ocean continental shelf width is only 37 km, while the slope has a mean width of 51.9 km (Harris et al. 2014). Consequently, the deep sea is very close to the coast in certain areas of the Indian Ocean, presenting an excellent opportunity to study deep continental margin habitats and faunal communities, as demonstrated for the oxygenated slopes of the Kenyan margin (Demopoulos et al. 2003) and the Tanzanian continental slope (Gates 2016). The Census of Marine Life's (CoML) Continental Margin Ecosystems on a Worldwide Scale (COMARGE) project further investigated the various complex ecosystems of the global continental margins, summarised by Menot et al. (2010), though most of the research effort was focussed on the Atlantic Ocean.

Of the countries bordering the Indian Ocean, the Australian continental margin is perhaps the best understood (McEnnulty et al. 2011). For example, the western continental margin of Australia was surveyed during 2005 and 2007 as a contribution to the COMARGE project (Williams et al. 2010, McEnnulty et al. 2011, Poore et al. 2015). Geoscience Australia's 2008-2009 survey provided further observations of deep-water benthic habitats and biota from the western margin such as 38 different types of Lebensspuren (Daniell et al. 2009, Przeslawski et al. 2012). Most notably, the southern Australian continental margin has been subject to intense research interest, especially areas of the Great Australian Bight that have been earmarked as

frontiers for oil and gas exploration (Rogers et al. 2013). The recent CSIRO-led Great Australian Bight Research Program and Great Australian Bight Deepwater Marine Program provided the deepest systematic biological sampling in Australian waters to date (Beg 2018, Ross et al. 2018, Smith et al. 2018). These programs collectively recorded 1,267 benthic invertebrate species from the Great Australian Bight's continental margin, including 401 new to science and numerous species records for the region (MacIntosh et al. 2018), and culminated in the publication of several studies of the margin's deeper ecosystems (Ross et al. 2017, CSIRO 2018, Fulton et al. 2018, Hook et al. 2018, Ross et al. 2018, Williams et al. 2018a, Williams et al. 2018b, Farrelly and Ah Yong 2019, Barrett et al. 2020). As with Gates et al. (2017), these examples benefitted from collaborations with the oil and gas industry, highlighting how hydrocarbon exploration can lead to and incentivise deep-sea research. Similarly, fisheries surveys, such as the Royal Society Indian Ocean Deep Slope Fishing Expedition and YugNIRO expeditions in the 20th century, produced valuable biological datasets for continental slope communities, though many are restricted to the upper 1,000 m (e.g. Forster et al. 1970).

Canyons

The continental slopes bounding the Indian Ocean are incised by numerous canyons that can reach several thousand meters deep. There are an estimated 1,590 canyons in the Indian Ocean, with a cumulative total area of 760,420 km², accounting for 11.2% of the continental slope (Harris et al. 2014). While some canyons are sub-sea extensions of large river systems (downwards formation, e.g., Indus and Swath of No Ground Canyons), others have no direct connection with land, having either been isolated from their terrestrial inputs by sea level fluctuations (e.g., Perth Canyon) or forming via repetitive landslides along steeper margins of the continental shelf (retrogressive slope failures, e.g., Bremer Canyon). Linking the continental shelf and abyssal seafloor, canyons have been recognised as key areas of localised productivity resulting in hotspots of benthic biodiversity in the deep sea (De Leo et al. 2010, Robertson et al. 2020). However, most large studies to date have excluded the Indian Ocean's canyon systems.

For example, a special issue from 2014 which presented 24 new studies on global submarine canyons did not include any research from the Indian Ocean (Huvenne and Davies 2014).

A recent increase in research interest in Australia's submarine canyons has resulted in detailed surveys for several canyons along the Western and Southern Australian coast (Heap et al. 2008, Daniell et al. 2009, Currie and Sorokin 2014, Conlan et al. 2015, Trotter et al. 2019, Post et al. 2021, Trotter et al. 2022a, Trotter et al. 2022b, Taviani et al. 2023). Geoscience Australia reported mapping 109 canyons along the western Australian continental margin in 2008-2009 (Daniell et al. 2009). The largest of these, the Perth Canyon, was first explored using an ROV during 2015, revealing a previously unknown deep-sea faunal community (Trotter et al. 2019). Perth Canyon is quiescent and described as a 'fossil canyon system', hosting extensive fossil coral graveyards that chronicle a history of thriving deep-water coral reefs during the last ice age (Trotter et al. 2019). In the modern day, the canyon is characterised by sporadic, localised hotspots of mega- and macrofauna that form relatively diverse communities along the canyon walls and rim, including sponges, corals, echinoderms, crustaceans, annelids, brachiopods, molluscs, and fish (Trotter et al. 2019). However, the Perth Canyon is relatively depauperate of life compared to the Bremer Canyon which lies further south, facing the Southern Ocean (Trotter et al. 2022a).

The dendritic Bremer Canyon system is comprised of several deeply incised submarine canyons extending over 4,380 km², including the Bremer, Hood, Henry, and Knob Canyons (Trotter et al. 2022a). Both the Perth and Bremer canyon systems are disconnected from continental river systems so receive very little detrital input; however, the Bremer system benefits from a greater prevalence of hard substrates, relatively high dissolved oxygen concentrations, and enhanced upper ocean mixing (Trotter et al. 2022a). Consequently, the 2020 *RV Falkor* FK200126 cruise to the Bremer Canyon recorded flourishing benthic communities with bryozoan and sponge-dominated forests and extensive coral gardens (scleractinians, octocorals, antipatharians), which form important community centres for echinoderms (cactus urchins, basket stars,

ophiuroids, crinoids), barnacles, anemones, crustaceans, and other taxa (Trotter et al. 2022a). Crucially, these studies demonstrate that the benthic communities of Australia's southwest canyons are comparable in diversity and density to the better-known nutrient-rich canyons of the North Atlantic and Northeast Pacific.

Similarly, the more northerly Cape Range and Cloates canyons off Exmouth are host to localised high densities of glass sponges and echinoderms, driven by the current-regulated sedimentation and nutrient availability (Post et al. 2020, Post et al. 2021). Another shallower study from a tributary of the Cape Range canyon system determined that these canyon habitats support richer fish assemblages than non-canyon habitats, with observed differences in the assemblages between the deeper and shallower areas of the canyon feature (Saunders et al. 2021). A comparison of macro- and megafaunal patterns at two topographically contrasting canyons at the southern Australian margin (the shelf-incising du Couedic and slope-limited Bonney canyon) further revealed the need to explore different canyon types along the Indian Ocean's continental margin, as well as the effects of different water masses and bioregions, to fully comprehend localised biodiversity patterns (Currie and Sorokin 2014, Conlan et al. 2015).

Several canyons have also been morphologically described along the African margin of the Indian Ocean, such as the Tugela Canyon off South Africa (Wiles et al. 2013) and the submarine canyons of northwest Madagascar (Wiles et al. 2019); however, there are no published data on their biological communities. Similarly, the benthic biodiversity of the large Swatch of No Ground and Indus canyons at the northern margins of the Indian Ocean have not been studied. Despite its name, the Zambezi Canyon is a north-south orientated submarine channel with no direct connection to the Zambezi River nor any other significant fluvial source (Wiles et al. 2017).

Owing to their proximity to terrestrial and coastal activities, submarine canyons are also of interest for the study of anthropogenic impacts to deep-sea environments. Canyons are considered potential sinks for marine litter (Hernandez et al. 2022) however, Australian

canyons appear to be relatively devoid of human traces (Trotter et al. 2019, Taviani et al. 2023), possibly owing to their predominately upward formation. Comparatively, Perth Canyon has more macro-litter than other southwestern Australian canyons, presumably due to its proximity to urban developments and commercial shipping operations (Taviani et al. 2023). Nonetheless, the physical imprint of human activities is remarkably absent relative to global patterns (Trotter et al. 2019). Conversely, demersal trawling is an increasingly concentrated threat to Australia's southern canyons, with large portions of the upper and mid slope trawled over the last two decades (Currie and Sorokin 2014, Conlan et al. 2015). Monitoring of continental shelf and slope environments is essential to understand the potential anthropogenic impacts to their related ecosystems, including canyons, yet the anthropogenic impacts faced by other Indian Ocean canyons remains understudied.

Fans

The northern Indian Ocean is host to the largest submarine fans in the world, accounting for more than half of the global fan area (Harris et al. 2014). Fans are depositional features that continuously deepen from their sediment source, usually at the mouth of a canyon (IHO 2019). The largest, the Bengal Fan, extends 3,000 km from the Swatch of No Ground Canyon to ~5,000 m depth in the Mid-Indian Ocean Basin, spanning 1,430 km at its widest point (Curray et al. 2003, Subrahmanyam et al. 2008). Sediments of the Bengal Fan have a maximum sub-seafloor thickness of 16.5 km, primarily derived from the Ganges-Brahmaputra River basin. To its east lies the smaller Nicobar Fan – an inactive sub-fan that was divided from the sediment supply of the Bengal Fan by the convergence of the Ninetyeast Ridge with the Java subduction zone in the late Pleistocene (Pickering et al. 2020). The Arabian Sea hosts the second largest submarine fan, the Indus Fan, which extends 1,500 km from the Indus Canyon at the Pakistan continental shelf break and is > 9 km thick at its thickest point (Clift et al. 2001). The natural significance of these fans lies in their transport of vast quantities of terrigenous sediments to the deep sea, serving as deep reservoirs of particulate organic carbon. As a result, the abyssal plain of the southern Bay

of Bengal is among the flattest large areas of the Earth's surface (Demopoulos et al. 2003). However, modern sea-level conditions are less conducive for sediment transport along the Bengal Fan via its turbidity channels than past states (Curry et al. 2003). The Netherlands Indian Ocean Expeditions further provided insights to the depositional architecture and sediment facies of the middle and lower Indus fan (van Weering et al. 1997). The direct influence of the submarine fans on the deep-sea biodiversity of the region has never been studied.

Oxygen minimum zones

The enclosed nature of the northern Indian Ocean is unique among global oceans, resulting in a high productivity-low ventilation system with strong stratification (McCreary Jr. et al. 2013, Hermes et al. 2019). Consequently, major oxygen minimum zones (OMZs), with dissolved oxygen concentrations < 0.5 ml/L, are located both in the Arabian Sea and Bay of Bengal at upper bathyal depths (Demopoulos et al. 2003, Rixen et al. 2020). These areas host ~21% of the total global volume of oxygen-depleted waters and ~59% of the Earth's marine sediments exposed to severe oxygen depletion (Rixen et al. 2020). Thus, despite the core OMZ (≤ 0.1 ml/L DO) occurring at depths shallower than 1,000 m (~250-750 m at the Indus margin), these OMZ strongly influence the community dynamics of the deep benthos across the continental margins of both basins (Rogers 2000, Gooday et al. 2010, Levin et al. 2013, Enge et al. 2014, Rixen et al. 2020). For example, strong edge effects of faunal diversity and density shifts with oxygen gradients have been observed across the OMZ boundaries at the Pakistan (Gooday et al. 2009, Hughes et al. 2009, Levin et al. 2009, Murty et al. 2009), Oman (Levin et al. 2000), and western Indian (Ingole et al. 2010, Levin et al. 2013, Enge et al. 2014) margins of the Arabian Sea, and in the Bay of Bengal (Gooday et al. 2010).

As well as shifts in oxygen concentrations, peaks in the abundance and diversity of deep-sea invertebrates at the lower boundaries of OMZs are thought to be strongly tied to the plentiful food supply available immediately beneath, resulting from the reduced utilisation of organic

matter during its downwards flux through the anoxic water column (Rogers 2000). At the Indian margin, foraminifera have been shown to rapidly process phytodetritus under almost anoxic conditions, suggesting they play an important role in short-term carbon cycling in this OMZ region (Enge et al. 2014). Furthermore, megafaunal aggregations at the lower OMZ boundary, including ophiuroids and *Encephaloides* spider crabs, have been shown to contribute substantially to the reworking and redistribution of organic matter at continental margins (Smallwood et al. 1999). There are however marked differences between the OMZs of the Arabian Sea and Bay of Bengal, primarily the occurrence of denitrification in the Arabian Sea and lack thereof in the Bay of Bengal (McCreary Jr. et al. 2013, Bristow et al. 2017). Consequently, differences in benthic faunal assemblages have also been observed between the two basins, as shown for foraminifera by Suokhrie et al. (2020). Demopoulos et al. (2003) provide a further summary of Indian Ocean OMZ research from the late 20th century.

Cold seeps and brine pools

Cold seeps are characterised by the expulsion of methane or other hydrocarbon-rich fluids from the seafloor and are associated with a diverse range of seafloor geomorphologies including mud volcanoes, pockmarks, diapirs, and brine pools (Foucher et al. 2009, Ceramicola et al. 2018). Various confirmed and inferred cold seeps occur along the northern continental margins of the Indian Ocean (Milkov 2000, Ceramicola et al. 2018), in areas such as the Makran Accretionary Prism off Pakistan (von Rad et al. 2000, Fischer et al. 2012), the Cauvery-Mannar and Krishna-Godavari Basins off India (e.g. *methane hydrates*: Mazumdar et al. 2019, *pockmarks*: Mazumdar et al. 2021), and within the Red Sea (e.g. *brine pools*: Purkis et al. 2022).

Of the abundant brine pools in the Red Sea, there are at least 25 pool complexes located in deep water (> 1,000 m) along the basin's axial trough (Purkis et al. 2022), the deepest of which lies at 2,850 m in Suakin Deep (Baumann et al. 1973). A further deep brine pool complex, named NEOM, was discovered in 2020 which extended the known range of Red Sea brine pools into the Gulf of Aqaba (Purkis et al. 2022). Despite their extreme conditions, cold seeps, including brine

pools, are often host to distinct bacterial and faunal communities driven by chemosynthesis of abundant hydrocarbon seepage (Sibuet and Olu 1998). The discovery of extremophile microbial diversity inhabiting these environments has provided a promising source of pharmaceuticals, with evidence that the Red Sea brine pool communities may yield bioactive molecules with potential for biomedical applications including anticancer and antibacterial properties (Ziko et al. 2019, Renn et al. 2021).

Within the Red Sea, large assemblages of bivalves, tubeworms and sponges, as well as dense bacterial mats, have been associated with the fringes of brine pools (Antunes et al. 2011, Batang et al. 2012). Predators will also make use of the brine as part of their feeding strategies, feeding on animals shocked or killed by the anoxic, hypersaline pools (Purkis et al. 2022). A new species of Corbulidae clam, *Apachecorbula muriatica*, observed in narrow bands along the margins of Red Sea brine pools is suggested to be both tolerant of and endemic to the hypersaline, hypoxic conditions of this extreme environment (Oliver et al. 2015). Corbulid clam shells are adapted to withstand such toxicity, for example, their unequal and globose valves form a tight seal when closed while conchiolin layers afford protection from carbonate dissolution (Batang et al. 2012). While cold seep bivalves have been shown to harbour chemosynthetic bacterial endosymbionts in other regions (Duperron et al. 2013), there is currently no available evidence that *A. muricata*, nor the wider bivalve assemblages recorded along the perimeter of Red Sea brine pools, are chemosymbiotic (Oliver et al. 2015, Purkis et al. 2022).

On the Indian margin of the Bay of Bengal, several active cold seeps were discovered in 2018 at the Krishna-Godavari Basin between 900-1,800 m deep, in association with shallow methane hydrates and supporting a rich biodiversity (Mazumdar et al. 2019, Dewangan et al. 2021, Sangodkar et al. 2023). Cold seep ecosystems have a long history at the Krishna-Godavari Basin with the discovery of cold-seep carbonate layers, including trace fossils of faecal pellet-filled burrow casts, deposited at least 46-58 kya (Mazumdar et al. 2009, Mazumdar et al. 2011). The modern faunal communities associated with the Krishna-Godavari cold seeps are diverse, with

over 2,300 individuals representing 18 macrofaunal families recorded (Sangodkar et al. 2023), and characteristic of global cold seep environments (Sibuet and Olu 1998).

Unlike the Red Sea brine pools, the Krishna-Godavari cold seeps support chemo-endosymbiont bearing bivalves of the genera *Bathymodiolus*, *Calyptogena*, *Conchocele*, and *Acharax*, as well as other chemosymbiotic species such as provannid gastropods and *Sclerolinum* siboglinid polychaetes (Mazumdar et al. 2019). They also host heterotrophs including munidopsid and galatheid squat lobsters, *Amphiodia* ophiuroids, and *Neolepas* goose barnacles (Mazumdar et al. 2019, Sangodkar et al. 2023), some of which represent new distribution records for species that were not previously known from the Indian Ocean (e.g., *Shinkaia crosnieri*, Gonsalves et al. 2022). Isotopic analyses provided a glimpse into the trophic structure of the seep ecosystems, revealing that *Bathymodiolus* are dependent on methanotrophic symbionts while *Sclerolinum*, *Conchocele*, and *Acharax* host thiotrophic symbionts (Peketi et al. 2022). A parasitic relationship was further observed between *Bathymodiolus* and the polychaete *Branchiopolynoe seepensis*, whereby the latter lives inside the mussel's mantle cavity and derives its food from its host's tissues (Mazumdar et al. 2019, Peketi et al. 2022). Meanwhile, the broad range of carbon and nitrogen isotope ratios are indicative of the variety of different food sources utilised by the heterotrophic fauna, from the soft tissues of the chemosynthetic organisms to suspended particulates to microbial mats and epibionts (Peketi et al. 2022).

Chemosynthetic organisms have similarly been recorded at cold seeps in the Cauvery-Mannar Basin between India and Sri Lanka, including *Lamellibrachia* tubeworms, *Munidopsis* squat lobsters, *Calyptogena* clams, and Buccinidae gastropods (Mazumdar et al. 2021). To date, the Krishna-Godavari and Cauvery-Mannar cold seeps are the only environments of their kind to be recorded within the Indian EEZ. Furthermore, as with the squat lobster *S. crosnieri* (Gonsalves et al. 2022), prior to their recent discovery, *Lamellibrachia* tubeworms had never been recorded from the Indian Ocean (Mazumdar et al. 2021), highlighting the need for further exploration and investigation of cold seeps and other reducing environments within the region.

Abyssal plains

Abyssal plains are extensive flat or gently sloping seafloor areas that occur at depths around 3,000-6,000 m (IHO 2019). Commonly considered the largest environment on Earth, abyssal plains dominate the abyssal depth zone and comprise ~35% of the Indian Ocean seafloor (Ramirez-Llodra et al. 2010, Harris et al. 2014). Abyssal plain habitats form the vast proportion of the seafloor environment of all major Indian Ocean basins, namely the Argo, Cuvier, Perth, Enderby, and Somali abyssal plains. Abyssal plains are generally characterised by vast areas of fine sediment tens to hundreds of metres thick, formed by the deposition of organic material from surface waters (Ebbe et al. 2010). However, recent research has revealed the heterogeneity of the abyssal seafloor with extensive hard substrate availability, including nodules and rocky outcrops (Figure 7B,G,H), especially when associated with more complex features such as fracture zones (Riehl et al. 2020). Similarly, abyssal hills (<1,000 m elevation) have also been recognised as sources of increased habitat heterogeneity and thus increased faunal biomass and diversity compared to surrounding plains (Durden et al. 2015). A boulder harbouring a diverse community of sponges and cnidarians observed in the North Australian Basin exemplifies this paradigm (Figure 7H), and given the geomorphological complexity of the Wharton Basin, this region of the eastern Indian Ocean may demonstrate comparable abyssal habitat heterogeneity (Figure 3).

The biodiversity and biogeography of abyssal plain communities are poorly known (Lörz et al. 2023). Low food availability and oligotrophic conditions in the abyss often result in low faunal abundance and biomass across the plains, while abyssal species richness has also been recorded as paradoxically high (Snelgrove and Smith 2002, Ebbe et al. 2010). In contrast, Parulekar et al. (1982, 1992) report rich fauna and high biomass for the abyssal benthos of the western and central Indian Ocean, linked to high organic production in the overlying euphotic zone. There are two major components to benthic abyssal plain communities, the epifauna and infauna, both of which play an important role in ecosystem functioning (e.g., bioturbation). Abyssal plain

epifaunal communities tend to be dominated by deposit feeders like holothurians (Figure 8A-D; Thurston et al. 1994, Iken et al. 2001). A 2005 cruise to the Crozet Plateau and surrounding plains resulted in the discovery of several new species of abyssal elasipodid holothurians (Cross et al. 2009, Rogacheva et al. 2009), demonstrating that even the most ubiquitous and conspicuous deep-sea taxa remain unknown from Indian Ocean abyssal plain habitats.

While smaller holothurians are also characteristic of abyssal infaunal communities, nematodes and copepods dominated the abyssal meiofaunal assemblage in the Arabian Sea, with an average total meiofaunal density of 234.99 ± 61.60 individuals per 10 cm^2 (Sautya et al. 2021). Meiofauna hold broad importance as bioindicators of ecosystem health that are capable of predicting global change (Zeppilli et al. 2015); however, further research is required to determine how the community patterns across abyssal plains influence the ecosystem's functioning (Sautya et al. 2021).

Another key feature of abyssal plains is the abundance of lebensspuren or 'life traces', which can often be more conspicuous than the organisms that make them (Jamieson and Onda 2022). For example, trails, burrows, spiral loops, and faecal casts were recorded at abyssal habitats at the western flanks of the Carlsberg Ridge in the Somali Basin (Glasby 1973), as well as star-shaped feeding traces of echiuran worms in the Bay of Bengal (Ohta 1984). Abyssal plain ecosystems are also host to benthopelagic mobile scavenger communities, including fish and crustaceans, which can be studied using baited deployment systems (see subsequent section on organic food falls; Bailey et al. 2007).

Indian Ocean abyssal plains are rarely prioritised for biological research. The little research effort that has occurred is spatially patchy such that not all basins are represented equally in the literature. Perhaps the best-known abyssal region in the Indian Ocean is the Mid-Indian Ocean Basin, also known as the Central Indian Ocean Basin (Ingole and Koslow 2005, Wafar et al. 2011). Seafloor surveys have been conducted here by Indian researchers since the 1980s to explore resource potential and inform environmental impact assessments for deep-sea mining

of the central Indian Ocean nodule field (Ansari 2000, Mukhopadhyay et al. 2002, Wafar et al. 2011). Data on the type and distribution of the area's meio-, macro-, and megabenthic fauna are suggestive of an extremely rich and diverse abyssal community (Ingole and Koslow 2005). However, this research represents only a limited area within the Mid-Indian Ocean Basin and given its driving objective to understand mining impacts (e.g., Ingole et al. 2005), these patterns may not reflect the faunal trends for the entire region (Ingole and Koslow 2005). Other abyssal plain research programmes in the Indian Ocean have focussed on the influences of surface productivity on particulate mass fluxes and the biomass and diversity of seafloor communities within the Arabian Sea (BIGSET project: Lochte and Pfannkuche 2000, Pfannkuche and Lochte 2000, Turnewitsch et al. 2000, Witte 2000, Witte and Pfannkuche 2000, Treude et al. 2002), Somali Basin (Netherlands Indian Ocean Expeditions 1992-93: van Weering et al. 1997), and Crozet Basin (CROZEX project: Hughes et al. 2007, Ebbe et al. 2010). In these basins, seafloor activity and community composition were found to be closely tied to the surface productivity patterns, though these interactions were complex and varied seasonally, often influenced by monsoonal dynamics (van Weering et al. 1997, Lochte and Pfannkuche 2000, Ebbe et al. 2010).

In contrast, there is a paucity of abyssal plain data elsewhere in the Indian Ocean. Bond and Jamieson (2022) highlighted the lack of research effort to understand abyssal plain ecosystems in Australia's EEZ, despite abyssal depths accounting for 46% of its jurisdiction. A recent report on Australia's Gascoyne Marine Park noted only that the Cuvier Abyssal Plain appeared to be typical of the habitat type in its vast homogenous sedimented plains interspersed with holothurians, cusk eels, xenophyophores, and glass sponges (Post et al. 2021). Furthermore, data were found to be sparse for deep-sea habitats across the southern Indian Ocean, particularly in the south-eastern region (Ramiro-Sánchez et al. 2023). Thus, clear knowledge gaps remain regarding the spatial and temporal biodiversity patterns, connectivity, productivity, and ecosystem functioning of abyssal plains in the Indian Ocean.




Figure 7: Examples of substrate and habitats of the deep Indian Ocean, where A = debris fields within a fracture zone, B = manganese nodules, C = volcanic crust of seamount, D-E = putative bacterial mats on trench forearc, F = rock face of escarpment, G-H = large boulder puncturing abyssal plain sediments, I = cobble and soft sediment mix on fracture zone floor, and J = soft sediment on subduction trench floor. All images from between 3000 and 7300 m in the East Indian Ocean.

Nodule fields

Polymetallic nodule fields, also known as (ferro)manganese nodules, are common across the abyssal seafloor and are especially abundant in areas with oxygenated bottom waters and low sedimentation rates (Figure 7B; Hein et al. 2013). Nodules are derived from the precipitation of metals from seawater forming concentric micro-layers around a nucleus, resulting in potato-sized (1-12 cm) deposits that can be rich in manganese, cobalt, nickel, and copper (Kuhn et al. 2017). These seafloor features provide important habitat heterogeneity through the availability of hard substrate for the benthic community (Figure 8J,S,X), with significantly higher densities of sessile and mobile epifauna recorded at areas with dense nodule coverage compared to nodule-free areas of the abyssal plain (Vanreusel et al. 2016).

In the Indian Ocean, nodule fields are primarily known from the Mid-Indian Ocean Basin (central Indian Ocean nodule field), Diamantina Fracture Zone (Cape Leeuwin nodule field), and Mozambique Basin. Of these, the central Indian Ocean nodule field (IONF) is the best characterised (see Mukhopadhyay et al. 2018 for a summary of IONF biota), owing to the deep-sea mining interest in the region (Hein et al. 2013, Mukhopadhyay et al. 2019). The IONF is comprised of two $\sim 150,000$ km² areas that have a combined estimated nodule resource of $\sim 1,300$ million metric tonnes, making this the world's second greatest nodule deposit in extent, abundance, and metal content (Mukhopadhyay et al. 2002, Mukhopadhyay and Ghosh 2010). As such, the Government of India holds an exploratory mining contract for a 75,000 km² area of the IONF, and a further 150,000 km² is reserved by the ISA for developing states to access deep-sea mineral resources (ISA 2019, Mukhopadhyay et al. 2019; Figure 1).

The Indian Deep-sea Environment Experiment (INDEX) was launched in 1995 to simulate and predict the environmental impacts of nodule mining in the IONF (Desa 1997). Meio- and macrofaunal communities were sampled before, 1 month after, and ~ 3.7 years after a simulated mining disturbance (Ansari 2000, Ingole et al. 2000, Ingole et al. 2001, Rodrigues et al. 2001, Ingole et al. 2005). Benthic communities experienced a significant decrease in both meiofaunal

and macrofaunal densities post-disturbance, with the top 2 cm of sediment experiencing the greatest physical disturbance (Ingole et al. 2000, Ingole et al. 2001). Similarly, benthic megafaunal populations, characterised by xenophyophores, holothurians, and sponges, experienced a 32% reduction post-disturbance (Rodrigues et al. 2001). While there was some recovery in nematode and oligochaete populations after 3.7 years, polychaete and crustacean populations did not recover to their pre-disturbance baseline (Ingole et al. 2005). Furthermore, a study integrating these results with 10 other simulated mining disturbance experiments from global sites concluded that the effects of nodule mining are likely to be long lasting (Jones et al. 2017).

Despite not appearing on recent maps of global nodule distribution (Hein et al. 2013, Miller et al. 2018), the Cape Leeuwin nodule field has been known since the 1970s and forms an extensive 1 million km² nodule pavement along the southern edge of the Diamantina Fracture Zone, from the Broken Ridge into the South Australian Basin (Kennett and Watkins 1975, Frakes et al. 1977, Bond and Jamieson 2022). However, the Cape Leeuwin deposits are of a lower grade and therefore are not currently of interest for deep-sea mining (Pettis and de Forest 1979, Frakes 1982). As such, there are no published studies of the benthic faunal communities associated with the Cape Leeuwin nodule field. Likewise, the Mozambique Basin nodules also have low-grade combined nickel, copper, and cobalt contents (Rogers 1995); therefore there is a lack of interest in the region despite a relatively high nodule abundance of 37.1 kg/m² and growth rate up to 248.3 mm/Ma (Geodekyan and Rozanov 1984, Ovechkina et al. 2021).

Nodules have also been recorded in the Wharton Basin at the Zenith Plateau (Weston et al. 2021), Wallaby-Zenith Fracture Zone (Niyazi et al. 2024), south of Christmas Island (Exon et al. 2002), at the western flank of the Carlsberg Ridge (Glasby 1973), and south of Tasmania (Watkins and Kennett 1971, Watkins and Kennett 1977, Bolton et al. 1988). Similarly, Nagender Nath et al. (1992) report nodules from the Mascarene and Somali Basins. Indeed, manganese nodule deposits appear to occur across the extent of the abyssal Indian Ocean, with varying

degrees of nodule density and grade (Mukhopadhyay et al. 2018). Despite this, very little work has been done to understand the importance of these areas for seafloor faunal communities in the Indian Ocean, with a focus on the economic value of the nodule fields instead and most polymetallic nodule research globally is concentrated in the high-grade Clarion-Clipperton Zone in the Pacific.

Journal Pre-proof




Figure 8 (previous page): Benthic biodiversity of the deep Indian Ocean, where A-D = Holothuroidae, E = Asteroidea, F = Brisingida, G = Echinoida, H-J = Polychaeta, K = Enteropneusta, L = Sipuncula, M = Gastropoda, N-O = Isopoda, P-R = Decapoda, S = Pantopoda, T-U = cf. *Caulophacus*, V-W = cf. Hyalonematidae, X-Y = Actiniaria, Z = Antipatharia, Aa = Phlebobranchia. All images from between 1000 and 7300 m in the East Indian Ocean.

Organic food falls

In an environment largely dependent on the transfer of food sources from surface waters, the deposition of animal carcasses and large plant matter to the seafloor is considered to play an important role in the nutritional needs of deep-sea species, including scavengers, predators and chemosymbiotic specialists (Smith et al. 2015, Jamieson 2016). Such events are termed 'food falls' or 'organic falls' and can include sunken whales (Smith and Baco 2003), large fish (Higgs et al. 2014), masses of gelatinous invertebrates (Billett et al. 2006), pelagic crustaceans (Christiansen and Boetius 2000), macroalgal rafts like kelp (Bernardino et al. 2010), and wood (Bienhold et al. 2013).

As naturally occurring food falls are ephemeral and extremely difficult to locate on the deep seafloor (Jamieson 2016), several studies have used experimental deployments to study the faunal communities associated with these environments, using different types of bait to mimic the various food fall types (e.g., Hilario et al. 2015, Amon et al. 2017, McClain et al. 2019). Baited camera and trap systems have been used in the deep Indian Ocean since the 1970s (Bailey et al. 2007) and have been successfully employed to study the assemblage composition and distribution of deep-sea scavengers (e.g. amphipods, macrourid fish, penaeid decapods) and other mobile bait-associated species (e.g. *Bassozetus*) across the Indian Ocean (Figure 9D-E & G-O; *Arabian Sea*: Witte 1999, Janßen et al. 2000, *Crozet Plateau*: Cousins et al. 2013, *Wallaby-Zenith Fracture Zone*: Weston et al. 2020, Weston et al. 2021, *Red Sea*: Pearce et al. 2022).

Other food fall experiments have focussed on the species colonising the bait itself, rather than mobile scavengers and opportunists captured on camera. These experiments often utilise packages of fresh mammal bones and are left for longer periods to simulate the latter

enrichment-opportunist and sulfophilic stages of food fall communities (Smith and Baco 2003). For example, the deployment of pig bones for two years on a mooring system on the Southwest Indian Ridge resulted in the first Indian Ocean records of the famous bone-eating zombie worms, *Osedax* (Zhou et al. 2020). A further two experiments, using whale bone and wood packages, were deployed at the Coral and Atlantis Bank seamounts on the Southwest Indian Ridge and left for two years (Amon et al. 2017); however, these deployments occurred at depths shallower than the remit of this review (≤ 750 m).

While several natural whale falls have been observed in the Atlantic, Pacific and Southern Oceans (Smith et al. 2015, Li et al. 2022), to date only one has been recorded in the Indian Ocean (Trotter et al. 2021, Trotter et al. 2022). This is surprising given the numerous whale populations and migrations that occur in the Indian Ocean (e.g. *blue whales in the northern Indian Ocean*: Anderson et al. 2012; *breeding and migrating humpback whales*: Jenner et al. 2001, Fossette et al. 2014; *pygmy blue whales in the eastern Indian Ocean*: Thums et al. 2022; *southern right whales in the southeast*: Carroll et al. 2011; *widespread occurrence of sperm whales*: Laran et al. 2017, Chambault et al. 2021), and perhaps highlights the paucity of exploratory deep-sea expeditions and reduced research capacity (e.g. ROV and submersible availability) in the region. Similarly, there are no studies of natural wood falls from the Indian Ocean, despite the likelihood of wood debris transported into the Bay of Bengal during monsoon season (Lee et al. 2019, Wohl and Iskin 2021) and observations of sunken wood in the Java Trench and off Tanzania (Gates et al. 2021, Jamieson et al. 2022). The only other records of natural organic falls in the Indian Ocean include observations of mass depositions of *Charybdis smithii* swimming crabs and *Crambionella* jellyfish in the Arabian Sea, in 1995 and 2002 respectively, both of which resulted in significant organic carbon fluxes that exceeded annual records by more than an order of magnitude (Christiansen and Boetius 2000, Billett et al. 2006).




Figure 9: Pelagic and demersal biodiversity of the deep Indian Ocean, where A = lobate Ctenophore, B = Trachymedusae, C = remosiid siphonophore, D-E = penaeid decapods, F = cirrate octopus, G = *Pseudotriakis microdon* (false catshark), H-I = Synphobranchid eels, J = Morid (blue hake), K = Macrourid (grenadier), L-M = Ophidiid (cusk eel) and N-O = Liparid (snailfish). All images from between 1000 and 7300 m in the East Indian Ocean.

Plateaus

Plateaus are large, relatively flat areas elevated from the surrounding relief with at least one steep side or escarpment (IHO 2019). They most often present as single features and can present from continental or oceanic affinities formed through volcanic activity, tectonic uplift of ocean crust, or rifting of continents where large fragments are left offshore. Several plateaus in the Indian Ocean abut continental margins and slope down on all remaining sides (e.g., Exmouth Plateau and Madagascar Plateau) whilst others are isolated offshore features, steep on all sides (e.g., Zenith Plateau and Mascarene Plateau). It is important to note that plateaus formed from

isolated volcanic activity create large igneous provinces which are geochemically different from mid-ocean ridges created at divergent plate boundaries (Frey et al. 2000). Occasionally, these two processes interact, and a plateau can break apart, form two isolated structures with escarpments where they once adjoined. Such is the case with Kerguelen Plateau and Broken Plateau which rifted apart at Broken Ridge.

The Kerguelen Plateau is the largest plateau (1,226,230 km²) in the Indian Ocean and third largest in the world (Harris et al. 2014). It rises more than 2,000 m from its surrounding basins and has been the focus of many geological surveys due to its enigmatic and geodynamic complexities (see Bénard et al. 2010). Clasts of continental crust have been found within the Kerguelen Plateau's basaltic flows which represent slithers of the Indian plate, left behind as India broke away from Antarctica and moved north (Frey et al. 2000). The Kerguelen Plateau has long been an important commercial fishing ground (Duhamel and Williams 2011). From the 1970s, Soviet trawlers targeted marbled rockcod, mackerel icefish and grey notothen. Since the establishment of French and Australian EEZs in 1978 and 1979, unrestricted trawling activity has gradually been replaced with longliners targeting Patagonian Toothfish (*Dissostichus eleginoides*) in 500-2,300 m water depth (Lord et al. 2006, Park et al. 2008, Palomares and Pauly 2011). Whilst fishing activity in these Kerguelen plateau EEZs is managed and regulated, the area did see a period of unregulated illegal longline fishing activity between 1997-2004 (Duhamel and Williams 2011).

Fortuitously, this commercial fishing activity has broadly facilitated observer contributions (Gasco 2011) and scientific research on a broad range of topics in an otherwise remote, isolated and hostile location, including: Southern Indian Ocean fronts (Belkin and Gordon 1996), slope currents (Park et al. 2008), cephalopod fauna and predator feeding behaviour (Cherel et al. 2004), the feeding ecology of two deep-sea skates (Faure et al. 2023), the macroinvertebrate assemblage (Martin et al. 2023), and life history and spatial distribution of fish (Peron et al.

2016). Less isolated plateaus such as the Exmouth Plateau off Western Australia are easier to access, however research there is also driven by extraction operations.

The Exmouth Plateau abuts the mid-west coastline of Western Australia and is rich in subsea hydrocarbon deposits. Permission to extract these deposits has required facility operators to undertake environmental assessments and monitoring, resulting in biological and ecological knowledge at depths and locations often forgotten (Macreadie et al. 2018, McLean et al. 2022). Although most operations exist on the top of the plateau, advances in technology have resulted in extraction and associated subsea infrastructure installed beyond 1,000 m. The marine community on the slope of Exmouth Plateau beyond 1,000 m loosely reflects that of the shelf break off Western Australia (Williams et al. 2001, Douglas 2021). Future research at Exmouth Plateau beyond 1,000 m will likely be driven by the exploration and extraction of hydrocarbons, however areas without extraction – oil and gas, fisheries or otherwise – may remain largely unexplored. One such area is the Naturaliste Plateau.

The Naturaliste Plateau is a large (90,000 km²) marginal plateau off the southwestern coast of Western Australia and, like the Kerguelen Plateau, has geological ties to the greater India and Antarctica plates after the break-up of Gondwana (Halpin et al. 2008). Three of its steep sides extend from 2,000 m to beyond 5,000 m, and its southern face edges to hadal depths at the eastern margin of the Diamantina Fracture Zone. Despite its proximity and physical presence, a lack of commercial interest in this plateau has not compelled any biological or ecological research beyond 2,000 m.

The Zenith Plateau and Cuvier Plateau (also known as the Wallaby Plateau) are two more plateaus that rise in the eastern Indian Ocean, situated in a geologically complex area surrounded by ridges and deep fracture zones. Like other plateaus in the Indian Ocean, their geology, formation and evolution are relatively well-described (see Symonds et al. 1998, Sayers et al. 2002, Daniell et al. 2010, Gibbons et al. 2012) compared to their biological community. Bond et al. (2023) classified the broad scale geomorphology of the Wallaby-Cuvier Escarpment

on the southern margin of the plateau (Figure 7F) and linked this to ecology, finding the highest diversity of organisms in the rocky debris field interlaced with soft sediment. The assemblage of macrobenthic organisms on the plateau and its slopes and escarpments are described as diverse but with exceptionally low abundance of animals (Daniell et al. 2010, Przeslawski et al. 2012, Bond et al. 2023). Low abundance of organisms could be a result of the oligotrophic conditions in the eastern Indian Ocean (Johannes et al. 1994, Lourey et al. 2006).

Conditions differ in the southeast Indian Ocean where iron-rich plateaus fertilise surrounding waters. Like the Kerguelen Plateau, the Crozet Plateau is located on the northern boundary of the Southern Ocean and within a larger region known to have high-nutrient, but low-chlorophyll, primarily due to low iron (Blain et al. 2007, Pollard et al. 2009). However, waters surrounding the plateaus like Crozet and Kerguelen are isolated hotspots of productivity caused by dissolved iron leaching from the shelf (Pollard et al. 2007, Wolff et al. 2011). Pulses of chlorophyll occur once irradiance increases during the austral spring and summer which sinks to the deep sea and increases the amount of organic matter available to organisms (Pollard et al. 2009, Wolff et al. 2011). Areas with high surface productivity induced by iron fertilisation have a different assemblage of macrobenthic organisms and fish compared to low iron areas nearby (Wolff et al. 2011, Cousins and Priede 2012). Plateaus like Crozet and Kerguelen play a nuanced role in shaping the biological community of their surrounding deep sea and understanding this role better, warrants further research.

The Indian Ocean hosts several other plateaus of note as listed in Appendix 1, including the Mascarene Plateau, the second largest in the Indian Ocean (Fisher et al. 1967), and the Agulhas, Mozambique and Madagascar Plateaus that lie off southeast Africa (Goslin et al. 1980, Parsieglia et al. 2008, Jacques et al. 2019). These plateaus have also been subject to commercial interest from fisheries, and consequently research interest (see Roberts et al. 2020), especially at the Madagascar (Clark et al. 2007, Rogers 2012) and Mozambique (Budnichenko et al. 1997, Parin

et al. 2008) Plateaus. However, there remains a paucity of biological data at depths greater than 1,000 m for these features, especially for benthic environments.

Ridges

A ridge is an elongated elevation with steep sides of varying complexity and size (IHO 2019), some of which may be seismically active while others are aseismic. The Indian Ocean has four main active spreading ridges: the Central Indian Ridge (CIR) which extends into the Carlsberg Ridge (CR) to the north, the Southwest Indian Ridge (SWIR) which is connected to the Mid-Atlantic Ridge system, and the Southeast Indian Ridge (SEIR) which is connected to the Pacific-Antarctic Ridge. Each of these ridge systems have differing characteristics and spreading rates, though their crusts are generally considered to be thinner than normal because of their slow spreading rates (Suo et al. 2016). They also host a diverse range of geological formations including guyots, seamounts and hydrothermal vents; the biodiversity of which are discussed in subsequent sections of this review.

The SWIR is the second slowest spreading ridge in the world at an ultraslow rate of ~15 mm/year (Patriat et al. 1997, van der Most et al. 2023). Investigations of the SWIR were instrumental in the discovery of ultraslow spreading ridges globally (Dick et al. 2003). One key artifact of the SWIR is its hypothesised role as a corridor for gene flow between the world's oceans (e.g., for hydrothermal vent fauna, German et al. 1998). Exploration of the ridge has resulted in the discovery of new species and new geographic records such as the sponge-associated polychaete *Bathymoorea*, a genus previously only known from the northeast Pacific (Periasamy et al. 2023). Conversely, the SEIR has a more uniform and intermediate spreading rate of 69-75 mm/year while the CIR has a slow to moderate spreading rate of ~48 mm/year (Sempéré and Cochran 1997, Kumagai et al. 2008, van der Most et al. 2023). Variation in spreading rates and the resulting stability of ridges often leads to differences in their biology (Copley et al. 2016). The CR to the north has a slow spreading rate of 22-32 mm/year (Murton and Rona 2015, van der Most et al. 2023). Sautya et al. (2017) provided the first quantitative

exploration of the benthic megafaunal assemblages on the CR, recording eight different substratum types and 90 megafaunal taxa from seven phyla, including cnidarians, echinoderms, poriferans and chordates. As with the SWIR, investigations of the CR have resulted in the discovery of new taxa including a new genus and species of glass sponge, *Indiella ridgenensis* (Sautya et al. 2011). In contrast, the SEIR remains one of the least biologically studied regions of the Indian Ocean, including for seamount and hydrothermal vent ecology (see subsequent sections), despite being a major active spreading centre. Indian Ocean ridges have also been a focal area for fisheries operations, with fishery activity historically covering most ridges of the Indian Ocean (Budnichenko et al. 1997, Romanov 2003; see subsequent section on seamounts). Other Indian Ocean ridges of note include the Ninetyeast Ridge, the longest linear intraplate rise globally which formed via hotspot trace of the Kerguelen mantle plume (Levchenko et al. 2021), the associated Broken Ridge (Frey et al. 2000), and the similarly formed Chagos-Laccadive Ridge via the Réunion hotspot trace (Sreejith et al. 2019); all of which are examples of aseismic ridges driven by intraplate volcanism.

Seamounts

The seafloor is scattered with volcanically derived features of discrete elevation that form at ridge-transform fault intersections, overlapping spreading centres, in intraplate regions and at hotspots, and that can occur as chains, clusters or in isolation (Figure 7C; Iyer et al. 2012). According to the IHO (2019), there are four distinct features that fall under the general umbrella of this definition, though applied definitions vary between studies (Rogers 2018). *Seamounts* are conically shaped features with elevations greater than 1,000 m from the surrounding seafloor, although the term 'seamount' is also readily used for features of lower elevations (> 100 m) in ecology (Wessel 2007), *guyots* are flat-topped seamounts, *knolls* have a rounded profile but with less than 1,000 m elevation, and *hills* are irregularly shaped knolls. The number of seamounts that occur globally varies depending on the estimation method and definitions used (Kitchingman et al. 2007, Clark et al. 2010, Yesson et al. 2021), with recent estimates for

the Indian Ocean ranging from 1,082 seamounts and 28 guyots (Harris et al. 2014) to 4,023 seamounts and 19,909 knolls (Yesson et al. 2011). Despite improvements in modern datasets, 'phantom seamounts' derived from misinterpretations of no-bottom-sounding data persist in updated predictions, with the majority of those identified concentrated in the Indian Ocean (Yesson et al. 2021). However, even by Harris et al.'s (2014) more conservative estimate, the Indian Ocean's seamounts occupy a considerable proportion of the ocean basin, with a collective area almost twice the size of Madagascar. For the purpose of this review, the term seamounts will encompass all these features hereafter (Clark et al. 2010).

Seamounts are globally recognised as areas of biological importance in the deep sea, supporting abundant benthic communities and forming complex habitats at their summit and flanks (Clark et al. 2010, Rogers 2018). Seamounts are also known to attract mobile megafauna such as cetaceans, sharks and large fish (Rogers 2018), the physical processes associated with their topography enhancing productivity or food supply within their vicinity (Rogers 2012). In the Indian Ocean, seamounts are mostly associated with ridge systems and amongst others found in abundance in the Mid-Indian Ocean Basin and along the Southwest Indian Ridge. Iyer et al. (2012) further provided a summary of the differing formation and emplacement of Indian Ocean seamounts with descriptions of several known seamounts, primarily in the northern and central Indian Ocean. However, research effort to explore the faunal communities and ecology of Indian Ocean seamounts is particularly poor compared to other regions (Clark et al. 2010).

Demopoulos et al. (2003) called for complete benthic habitat descriptions for Indian Ocean seamounts, yet scientific knowledge 20 years later is still lacking. Roberts et al. (2020) summarise the history of seamount exploration and fisheries research in the Indian Ocean. Species records collated by the SeamountsOnline initiative up to 2010 are limited for the Indian Ocean, with only 1,148 records for 279 species compared to 12,010 records for 1,953 species in the Pacific and 6,588 records for 1,141 species in the Atlantic (<http://seamounts.sdsc.edu/>). Seamounts in the southern Indian Ocean support aggregations of commercially important fish

species, and biological datasets predominantly come from the deep-sea fishing industry or from national fisheries research programmes (Romanov 2003). Furthermore, the majority of these Indian Ocean records are from Walters Shoal, a collection of shallow (18 m) seamounts at the Madagascar Plateau (Rogers 2012), thus are not representative of the biodiversity of deeper seamounts in the region. Similarly, extensive seamount chains and abundant isolated seamounts have been mapped in the Mid-Indian Ocean Basin (Das et al. 2007), yet their faunal communities remain unknown.

Most seamounts in the Indian Ocean have summits > 1,000 m, with a mean summit depth of ~2,400 m (Yesson et al. 2011), yet these deeper seamounts are rarely explored (Clark et al. 2010). Instead, the little benthic research that has been conducted for Indian Ocean seamounts has generally been restricted to those with shallower summits (< 1,000 m depth), primarily at the SWIR. Two major research cruises surveyed the area in 2009 and 2011 as part of the NERC/IUCN Seamounts Project, specifically targeting the Coral, Melville Bank, Middle of What, Sapmer Bank, and Atlantis Bank seamounts (Rogers and Taylor 2011, Rogers et al. 2017). The SWIR seamounts are of biogeographical interest owing to the Agulhas, Subtropical and Subantarctic Fronts that demarcate the region (Pollard and Read 2017, Read and Pollard 2017, Rogers et al. 2017). These surveys resulted in the discovery of several new species including octocorals (Taylor and Rogers 2017), carnivorous sponges (Hestetun et al. 2017) and holothurians (O'Loughlin et al. 2013). Diverse faunal communities were also discovered from within coral rubble and carbonate material atop the seamounts, including 34 different polychaete families (Narayanaswamy et al. 2017). Furthermore, given the commercial interest in seamount-based fisheries in the region, a strong focus was placed on understanding the productivity dynamics and pelagic communities of the SWIR seamounts (Budnichenko et al. 1997, Djurhuus et al. 2017, Laptikhovsky et al. 2017, Letessier et al. 2017, Rogers 2017, Rogers et al. 2017, Sonnekus et al. 2017, Rogers 2018, Cherel et al. 2020).

The megabenthic communities of seamounts have also been explored in the Andaman Sea, whereby two seamounts with different summit depths were surveyed and compared to the biodiversity of the adjacent basin floor. Faunal abundances were found to be greatest at the seamounts compared to the surrounding basin, with the highest number of taxa observed at their flanks (Sautya et al. 2011). Distinct faunal communities were recorded between the shallow- (434 m) and deep-summited (1,336 m) seamounts, the deeper of the two dominated by porifera and cnidaria at its summit, as well as arthropods and other taxa at its more biodiverse flank (Sautya et al. 2011). Six ophiuroid species were also collected at the Andaman seamounts, all of which were new records for the Andaman Sea including two new species (Stöhr et al. 2012).

The research conducted at the SWIR and Andaman Sea highlights the biological importance of seamounts and emphasises the need to pursue similar exploratory studies for deeper seamounts in the Indian Ocean (Clark et al. 2010). Further, seamounts are highly variable in morphology and depth ranges, with implications for seamount ecology. For example, five seamounts mapped at the Southwest Indian Ridge (SWIR) differed in their seascape composition and configuration despite belonging to the same geological feature (Swanborn et al. 2023a). The varied seascapes and spatial heterogeneity of seamount habitats influenced the biodiversity supported by these seamounts and likely their ecological functioning (Swanborn et al. 2023ab). However, such interactions have not yet been explored for the majority of Indian Ocean seamounts, compounding the need for increased ecological research.

Recent surveys as part of baseline environmental and benthic ecological assessments for deep-sea mining in the Indian Ocean have been a catalyst for new biodiversity discoveries for deep-summited seamounts. Such exploration has resulted in the description of several new species and new species distribution records for seamounts on the Central Indian Ridge and SWIR, including bamboo corals (Periasamy et al. 2023a), black corals (Periasamy et al. 2023b), octocorals (Periasamy et al. 2023c), carnivorous sponges (Periasamy and Ingole 2022),

Spongicoloides shrimp (Periasamy et al. 2022), and squat lobsters (Periasamy et al. 2023d). As well as improved research effort for deeper seamounts and for Indian Ocean seamounts in general, future research should also include more detailed study of the flanks and bases of seamounts given the indications that these areas support distinct assemblages (Baco 2007, Sautya et al. 2011). For example, autonomous landers deployed to the base of the Afanasij Nikitin Seamount (~ 4,700 m) in the Mid-Indian Ocean Basin in 2017 resulted in the collection of seven amphipod species including two potentially new abyssal species of *Hirondellea* and *Cyclocaris* (Weston et al. 2021).

The anthropogenic threats to Indian Ocean seamounts are stark. Demersal and benthopelagic fish have been trawled at SWIR seamounts since the 1970s (Marsac et al. 2020). Today, seamount fisheries are prominent in the southern Indian Ocean including seamounts at the Southwest Indian Ridge, Conrad Rise, Mozambique and Madagascar Plateaus, and Ninetyeast and Broken Ridges (Clark et al. 2007: Table 17.9). Five SWIR seamounts were classified as SIOFA BPAs (Walters Shoal, Coral, Middle of What, Atlantis Bank and Fools Flat) in 2018. While these seamounts were part of the voluntary SIODFA BPAs prior to this (Shotton 2006), the SIOFA designation provides active conservation and management measures prohibiting bottom fishing in the BPAs (Marsac et al. 2020, SIOFA 2023, SIOFA Secretariat 2023). As with other deep-sea environments, marine litter often pollutes seamounts, the majority associated with lost fishing gear (Woodall et al. 2015, Rogers 2018), and microplastic fibres have also been found in the sediments and biota of SWIR seamounts (Woodall et al. 2014, Taylor et al. 2016). Finally, although no mining contracts have yet been agreed for Indian Ocean seamounts, they may nonetheless be future deep-sea mining targets for the cobalt-rich ferromanganese crusts (CFC) that form at their summits (Halbach et al. 2017, Miller et al. 2018), thus improved research effort is critical for understanding potential CFC mining impacts in the region.

Like seamounts, oceanic islands are topographic elevations rising from the seafloor that penetrate the sea surface instead. They often form part of larger chains but have never been

connected to the mainland (Dawson 2016). Several nations bordering the Indian Ocean have oceanic islands in their territory, such as the Seychelles and Maldives in the western Indian Ocean, India's Lakshadweep islands, and Australia's Cocos-Keeling archipelago in the eastern Indian Ocean. Deeper slopes of oceanic islands in the Indian Ocean equally exhibit complex geomorphology hosting rich biodiversity. For example, a 2022 expedition to the Cocos-Keeling archipelago revealed variety of deepwater habitats and species potentially new to science along atoll slopes, providing data that will also assist management plans (O'Hara 2022). However, atoll slopes > 1,000 m typically remain under-sampled in the Indian Ocean, despite the importance for marine management in national waters.

Hydrothermal vents

Several Indian Ocean ridge systems are associated with hydrothermal vents. Hydrothermal vents are insular, chemosynthetic environments that occur in geothermally active areas where seawater percolates into the oceanic crust and is reemitted as superheated, mineral-rich fluid, which precipitates to form large chimney structures and supports a plethora of life via chemosynthesis (Van Dover 2000). Active hydrothermal vent communities were first discovered in the Indian Ocean in 2000, with the description of faunal assemblages at the Kairei vent field just north of the Rodriguez Triple Junction on the Central Indian Ridge (CIR) (Hashimoto et al. 2001). Earlier evidence of a potential active vent community was recorded from the Southeast Indian Ridge (SEIR), with the retrieval of a vent-associated *Neolepas* barnacle from a dredge at the Site 21 vent field (Scheirer et al. 1998, Watanabe et al. 2018); however, this vent field has never been directly observed and no further publications have confirmed faunal activity at this site. Including Site 21, a total of 15 active vent fields have been observed: three along the Carlsberg Ridge (CR), seven on the Central Indian Ridge (CIR) (including two newly discovered sites), three on the Southwest Indian Ridge (SWIR), and two on the Southeast Indian Ridge (SEIR) (Table 3; van der Most et al. 2023, Jang et al. 2023).

Despite falling short compared to the research effort in the North Pacific and at the Mid-Atlantic Ridge (Thaler and Amon 2019), hydrothermal vents are undoubtedly the best biologically characterised deep-sea ecosystem in the Indian Ocean, as summarised by Perez et al. (2021) and van der Most et al. (2023). This is likely owing, in part, to the deep-sea mining interest for the polymetallic sulfides that form the chimney structures of hydrothermal vents. Exploratory mining contracts have been granted by the International Seabed Authority (ISA) to the China Ocean Mineral Resources and Development Association (COMRA), Federal Institute for Geosciences and Natural Resources of Germany (BGR), and the Governments of India and Korea (Table 3, Figure 1). Within the Indian Ocean, the region most in need of increased hydrothermal vent study is the SEIR, with 50 inferred active vent fields listed on the InterRidge global database that have been identified by plume activity but never observed directly (Beaulieu and Szafranski 2020).

Indian Ocean vents form their own biogeographic province, hosting several unique species that have not been recorded elsewhere (Van Dover et al. 2001, Rogers et al. 2012, Copley et al. 2016, Perez et al. 2021), including, most notably, the scaly-foot snail *Chrysomallon squamiferum* (Chen et al. 2015). The Indian Ocean also hosts a high degree of locally endemic species known only from a single vent field, such as the gastropod *Dracogyra subfusca* (Chen et al. 2017, Perez et al. 2021, Thomas et al. 2021a). Further, four newly described species of *Anatoma* gastropods are considered endemic to vents in the central Indian Ocean (Hoffman et al. 2022). Such high levels of biodiversity uniqueness contributed to the evaluation that Indian Ocean vent species are under the greatest threat of extinction from deep-sea mining of the nine global vent biogeographic provinces (Thomas et al. 2021b). Using vent-endemic molluscs as representative taxa and given the incidence of several deep-sea mining contracts within the region (Figure 1), 60% of species assessed were found to be Critically Endangered with 100% considered threatened (Thomas et al. 2021b). Furthermore, van der Most et al. (2023) determined that all active vent fields in the Indian Ocean meet the scientific criteria for protection under CBD, IMO and FAO guidelines. At a higher taxonomic level, however, there are broad similarities between

the faunal communities of vent fields in the Indian Ocean and those in the Atlantic, Pacific and Southern Oceans (e.g., *Kiwa* yeti crabs, *Rimicaris* shrimp, *Neolepas* barnacles, *Bathymodiolus* mussels, *Lepetodrilus* limpets, *Alviniconcha* gastropods, *Hesiolyra* polychaetes; Bachraty et al. 2009, Copley et al. 2016, Roterman et al. 2018, Zhou et al. 2018, Gerdes et al. 2019, Wang et al. 2020).

Within the Indian Ocean province, there is a documented trend of continuous biogeographic transitions between the CR, CIR, and SWIR (Zhou et al. 2022). Distinct biogeographic subdivisions of the Indian Ocean province have been inferred accordingly, based on community structure and genetic differentiation of species located across multiple vent fields (Table 3). The distinctive clustering of vent communities between the proposed sub-provinces of the CR, CIR (including the Tiancheng vent field on the northern SWIR), and southern SWIR, is suggested to be linked to the numerous transform faults and fractures that scar these ridge systems (Sun et al. 2020, Zhou et al. 2022). A further phylogeographic boundary was also proposed between the Onnuri and Solitaire vent fields on the CIR based on the genetic divergence of *Alviniconcha* gastropods (Jang et al. 2023). At a local scale, faunal communities at Indian Ocean vent fields follow patterns of spatial zonation from chimneys with high hydrothermal activity, to areas of diffuse flow, to the vent periphery (Watanabe and Beedessee 2014, Perez et al. 2021), as observed at other global vent fields (e.g., Marsh et al. 2012). For example, endemic *Rimicaris* shrimp often dominate the active venting chimneys, with abundant chemosymbiont-hosting *C. squamiferum* and *Gigantopelta* gastropods in areas of diffuse flow venting (Copley et al. 2016, Wang et al. 2021). In contrast, the vent periphery is largely host to organic-enrichment opportunists that are also capable of colonising other reducing environments, rather than vent specialists, such as the holothurian *Chiridota hydrothermica* (Thomas et al. 2020, Thomas et al. 2022).

The faunal communities at inactive hydrothermal vents are far less understood than active sites, despite the probability that these inactive sites will be the primary focus of prospective

polymetallic sulfide mining activity (Van Dover 2011, Gerdes et al. 2019). Inactive and diffuse venting sites are often closely associated with active vent fields, which hampers their exploration as they are rarely prioritised. Inactive vents are those that do not currently exhibit fluid flow but are distinguished from extinct vents by their geological connection to an active source, such that they may potentially become active again in future (Jamieson and Gartman 2020). The Duanqiao vent field on the SWIR was initially thought to be inactive, until vent-endemic taxa were discovered colonising an edifice with weak, but active, fluid venting (Zhou et al. 2018, Van Dover 2019).

There are indications that inactive vent sites may host their own specialist communities (Gerdes et al. 2019, Van Dover 2019, Kniesz et al. 2022). For example, two previously unknown species of *Amphisamytha* polychaetes collected from an inactive chimney at the Longqi vent field have not been recorded elsewhere and are potentially specific to this environment (Zhou et al. 2018, Zhou et al. 2019). By studying a stand-alone inactive vent field, Gerdes et al. (2019) confirmed that the faunal assemblage at the hydrothermally extinct Gauss field is statistically distinct from the communities at the adjacent Edmond vent field, its periphery, and the surrounding hard- and soft-substrate non-vent areas. This discovery supports the existence of unique faunal assemblages related to inactive vent sites (Gerdes et al. 2019). Furthermore, inactive polymetallic sulfides at the Kairei and Pelagia vent fields were found to host abundant microbial communities including microorganisms that are unique to the site or rare elsewhere (Han et al. 2018). Further known inactive hydrothermal fields in the Indian Ocean include the MESO Zone on the CIR and Mount Jourdanne on the SWIR, however, the true number of inactive deposits remains unknown and requires more exploration of the Indian Ocean's ridge systems, especially further away from the ridge crest (Jamieson and Gartman 2020).

Table 3: Known active hydrothermal vent fields in the Indian Ocean, adapted from van der Most et al. 2023 and Jang et al. 2023. Colours denote the proposed biogeographical subdivision of hydrothermal vent communities by Zhou et al. 2022 and Jang et al. 2023, with dashed lines denoting proposed geographic boundaries within a ridge system. Mining contracts – Korea: Government of Korea, BGR: Federal Institute for Geosciences and Natural Resources of Germany, India: Government of India, COMRA: China Ocean Mineral Resources and Development Association. Biogeographic sub-provinces – CR: Carlsberg Ridge, (n/s)CIR: (north/south) Central Indian Ridge, (n/s)SWIR: (north/south) Southwest Indian Ridge, SEIR: Southeast Indian Ridge.

Ridge	Vent Field	Depth (m)	Mining Contract	Biogeographic Sub-province		Key References
				Zhou et al. 2022	Jang et al. 2023	
Carlsberg Ridge	Daxi	3,450	-	CR	CR	Wang et al. 2021, Zhou et al. 2022
	Wocan	2,973-3,105	-	CR	CR	Wang et al. 2017, Zhou et al. 2022
Central Indian Ridge	Tianxiu	3,500	-	CR	CR	Zhou et al. 2022, Qiu et al. 2023
	Onnare	2,984	Korea	-	nCIR	Kim et al. 2020, Jang et al. 2023
	Onbada	2,523	Korea	-	nCIR	Kim et al. 2020, Jang et al. 2023
	Onnuri	1,990-2,170	Korea	CIR-nSWIR	nCIR	Kim et al. 2020, Jang et al. 2023
	Dodo	2,745	°	CIR-nSWIR	sCIR	Nakamura et al. 2012
	Solitaire	2,606	°	CIR-nSWIR	sCIR	Nakamura et al. 2012
	Edmond	3,290-3,320	BGR	CIR-nSWIR	sCIR	Van Dover et al. 2001
	Kairei	2,415-2,460	BGR	CIR-nSWIR	sCIR	Hashimoto et al. 2001
Southwest Indian Ridge	Tiancheng	2,682-2,729	India	CIR-nSWIR	nSWIR	Tao et al. 2014, Zhou et al. 2018, Sun et al. 2020
	Duanqiao	1,732	COMRA	sSWIR	sSWIR	Tao et al. 2014, Zhou et al. 2018
	Longqi	2,755	COMRA	sSWIR	sSWIR	Tao et al. 2012, Copley et al. 2016, Zhou et al. 2018
Southeast Indian Ridge	Pelagia	3,659	BGR	-	SEIR	Gerdes et al. 2019
	Site 21*	~2,800	-	-	SEIR	Scheirer et al. 1998, Watanabe et al. 2018

*Note that vent activity has not been directly observed at Site 21 and the presence of a faunal community is evidenced only by the collection of *Neolepas* barnacles by dredge sampling (Scheirer et al. 1998). °The Dodo and Solitaire vent fields lie within the Mauritius EEZ under national jurisdiction.

Subduction trenches

The deepest known point in the Indian Ocean is located in the Java Trench which is a subduction trench that reaches a maximum depth of 7,187 m (± 13 m) (Bongiovanni et al. 2022).

Subduction trenches form at convergent plate boundaries. The Java Trench is formed as the Indo-Australian Plate subducts beneath the Eurasian Plate (Jamieson et al. 2022). Of all the trenches known globally, the Java Trench is the only subduction trench extending into the hadal zone in the Indian Ocean, making it a relatively geographically isolated hadal ecosystem (Jamieson et al. 2022). Consequently, hierarchical cluster analysis of hadal 'isolation' grouped the ~3,200 km long Java Trench with West Pacific hadal sites rather than others in the Indian Ocean (Stewart and Jamieson 2018).

Note that not all seafloor features named as trenches are associated with the hadal zone nor active subduction. For example, the Amirante Trench is a ~5,200 m deep structure located south of Seychelles within the Mascarene Basin which has attracted the interest of geoscientists due to its enigmatic morphology. The Amirante Ridge/Trough Complex shares characteristics with subduction zones, transform faults and mid-oceanic ridges (Damuth and Johnson 1989, Stephens et al. 2009, Rodriguez 2020) and early theories posited it might be a potential asteroidal impact site (Hartnady 1986). Recent research suggests that its origin may be linked to spreading along the Carlsberg Ridge and the India-Seychelles breakup events occurring around 63-65 Ma (Mukhopadhyay et al. 2012, Rodriguez 2020). A full-coverage high-resolution map of the Amirante Trench was produced in 2020 (Caladan Oceanic 2020), however, further research is required to uncover the ecosystems and biodiversity the Amirante Trench hosts.

Jamieson et al. (2022) summarised the biodiversity of the Java Trench, recording over 70 putative species across 10 phyla and 21 classes. A five-day expedition to the Java Trench in 2019 represents the only modern biological survey of this hadal trench, yielding several new species records and range extensions, including: the first record of hadal liparid snailfish in the Indian Ocean (Jamieson et al. 2022), first hadal cephalopod (Figure 9F; Jamieson and Vecchione

2020), deepest global larvaceans (Jamieson and Linley 2021), and deepest Indian Ocean decapods (Swan et al. 2021). Additionally, the first hydrographic profiles of the trench were also recorded, providing bottom water conditions of 1.54°C for in-situ temperature and 34.70 for practical salinity (Kolbusz et al. 2024). The diverse biological community of the Java Trench is characterised by elasipodid holothurians, amphipods including the supergiant *Alicellea gigantea*, *Bassozetus* cusk eels, and hexactinellid sponge aggregations (Jamieson et al. 2021, Jamieson et al. 2022). These faunal patterns are typical of hadal trenches (e.g., Jamieson et al. 2013, Mironov et al. 2019). There are, however, some noticeable distinctions between Java and other global trench systems, such as the absence of *Hirondellea* amphipods despite their prevalence elsewhere at hadal depths (Weston and Jamieson 2022, Jamieson and Weston 2023). The Java Trench seafloor can be categorised into six different habitat types which drives benthic species distributions within the trench (Jamieson et al. 2022). For example, faunal community shifts are apparent between the flat, fine-grain sediment-covered trench and plateau floors, and the steeper, bedrock-exposed slopes, with holothurians and brittle stars or anemones and asteroids dominating these respective communities (Figure 7); Jamieson et al. 2022). Patchiness in species distributions can also be linked to this habitat heterogeneity, exemplified by the discovery of the dense hexactinellid sponge aggregations only beneath overhangs of the upper trench slope (Jamieson et al. 2022).

Likewise, spatial patchiness may be a feature of the potential hadal chemosynthesis observed in the Java Trench. Numerous putative chemosynthetic bacterial mats were observed exploiting the structural weaknesses in the exposed bedrock of the forearc escarpment (Figure 7D,E), however, it remains to be seen whether they are restricted to these steep, structurally complex areas within the trench (Jamieson et al. 2022). Similar bacterial mats were recorded in association with outcrops and talus at Sirena Deep in the Mariana Trench (Hand et al. 2020) and it is possible the hadal chemosynthesis phenomenon may occur in other areas of structurally

weak bedrock on the fore-arcs of global subduction trenches (Blankenship-Williams and Levin 2009, Jamieson et al. 2022).

Sedimentation rates vary within and between trench systems, in turn influencing this habitat heterogeneity (Stewart and Jamieson 2018). For example, the accumulation of sediment in the deepest part of trenches (see ‘trench resource accumulation depth’ in Jamieson et al. 2010) often supports higher species abundances, typified in the Java Trench by increases in the number of holothurians and anemones observed at the trench floor and towards the base of the trench slope, respectively (Jamieson et al. 2022). However, despite lying within the ultra-productive Sunda-Arafura Sea Coastal biogeochemical province (Longhurst 2007), particulate organic carbon (POC) flux models suggest the Java Trench receives lower annual POC input than > 50% of other major global trenches (Lutz et al. 2007, Stewart and Jamieson 2018).

Transform faults and fracture zones

Transform faults are the active perpendicular offsets between two ridge segments of a mid-ocean spreading centre which extend into inactive fracture zones either side of the main ridge system (Hensen et al. 2019), collectively referred to here as fracture zones. Some of these fracture zones can reach hadal depths (> 6,000 m), of which there are eight in the Indian Ocean (Table 4, Figure 10). Only two of these have been mapped and their hadal status confirmed: the Wallaby-Zenith Fracture Zone (WZFFZ) to the east (6,675 m, Niyazi et al. 2024), and the Diamantina Fracture Zone (DFZ) in the southeast (7,019 ± 17 m, Bongiovanni et al. 2022). There are several other named fracture zones that intersect the mid-Indian Ocean ridges (Appendix 1).

Compared to subduction trenches, the WZFFZ is a small hadal feature (Weston et al. 2021) but is an area of tectonic importance as a consequence of the breakup of Greater India and Australia-Antarctica subcontinents and the opening of the Indian Ocean (Ali and Aitchison 2005, Gibbons et al. 2012, Olierook et al. 2015). Published accounts of the fauna at WZFFZ to date include amphipods (Weston et al. 2020;2021), fish (Jamieson et al. 2021), decapods (Swan et al. 2021), and the functional feeding strategies of demersal and benthic macrofauna (Niyazi et al. 2024).

Weston et al. (2020) described a new genus and two new species of amphipod from the WZfZ from abyssal depths within the fracture zone. Furthermore, cluster analysis of community structure compared to those of other hadal features in the Pacific Ocean suggested that the assemblage hadal amphipods at WZfZ is merely an extension of the abyssal community, despite the abundant presence of the hadal specialist *Bathycallisoma schellenbergi* (Weston et al. 2021; 2022). As such, and conversely to community structure shifts observed in subduction trenches, Weston et al. (2021) postured that the WZfZ represents an area of transitional community composition from abyssal to hadal, likely owing to the comparable seafloor topography between the upper Zenith Plateau and the flat bottom of the hadal fracture zone.

First discovered in 1960 (Jempson 2019), the DFZ is also an area of geological importance for studying ancient tectonic patterns, including the breakup of Australia from Antarctica (Markl 1978, Mammerrickx and Sandwell 1986). Like the WZfZ, the DFZ is one of the smallest hadal areas globally (Stewart and Jamieson 2018). It is, however, the deepest of the Indian Ocean's fracture zones and its deepest point, the Dordrecht Deep, was once believed to be the deepest part of the Indian Ocean (Stewart and Jamieson 2019). For this reason, the Five Deeps Expedition, which mapped the deepest points in each ocean, surveyed the Dordrecht Deep in 2019 and found it to be less than 200 m shallower than the Java Trench (Bongiovanni et al. 2022). A single lander deployment made at the Dordrecht Deep recovered amphipods, including the cosmopolitan hadal species *B. schellenbergi*, which is also known from WZfZ and the Java Trench (Weston et al. 2022). Due to the surveys of Geoscience Australia (2017), the Five Deeps Expedition (Bongiovanni et al. 2022), and multibeam data from the search for flight MH370 (Picard et al. 2018) the WZfZ and DFZ are remarkably well-mapped deep-sea areas of Indian Ocean.

There are no available biological data for any of the other six hadal fracture zones in the Indian Ocean and, other than the Vema Trench. Contrary to its moniker, the Vema Trench is in fact a fracture zone which offsets the Central Indian Ridge by ~300 km between 7°40' and 10°40' S,

ranging from depths of 3,000-4,000 m at its peripheries to > 6,000 m at its deepest point (Dhawaskar et al. 2018). The Vema Trench was once thought to be the deepest point in the western Indian Ocean (Heezen and Nafe 1964); however, other features appear to be deeper based on modern bathymetric data (Table 4). There are no published biological surveys of the area, and most geological surveys are from the 1960-70s (e.g., Heezen and Nafe 1964, von Herzen and Vacquier 1966, Fisher et al. 1967; 1971). Incidentally, one of the most prominent fracture zones in the Atlantic Ocean is also named Vema (Devey et al. 2018); both sites seemingly named after the RV *Vema* (Heezen and Nafe 1964).

Many hadal areas in the central and western Indian Ocean have never been explored and their depths, currently derived from satellite altimetry data, have not been verified by multibeam echosounder bathymetry. This variation in the quality of available bathymetric data demands caution in the interpretation of the extent and maximum depth of hadal features (Table 4), especially in lesser-known areas where values may be overestimated, such as the western Indian Ocean (Jamieson et al. in review).

The paucity of biological studies for the hadal fracture zones in the Indian Ocean demonstrates how non-subduction hadal features are rarely prioritised for biological research even by modern deep-sea exploration programmes. Studies of the hadal zone have largely focussed solely on subduction trenches to date, overlooking non-subduction features such as fracture zones and troughs, despite their importance in unravelling the ecology and evolution of hadal biodiversity (Weston et al. 2021). Although less conspicuous and often shallower than subduction trenches, the discovery of endemic hadal species from these relatively small and isolated fracture zones is suggestive of their importance in the global connectivity of hadal environments (Jamieson and Stewart 2021, Weston et al. 2022). Since 2021, the Minderoo-UWA Deep-Sea Research Centre have undertaken research cruises to both the WZFD (DSSV *Pressure Drop* 'Indomitable' expedition) and DFZ (*Pangea Ocean Explorer*) to fill these knowledge gaps. Using baited landers and a manned submersible, these cruises aimed to map habitat types

(Figure 7A,I) and study the biological communities in these areas (Figures 7-9; Niyazi et al. 2024). However, no such expeditions have been conducted or planned for the hadal fracture zones of the central and southwestern Indian Ocean.

Deep troughs and basins

As well as subduction trenches and fracture zones, hadal areas are also found scattered across Indian Ocean basins (Figure 10). A deep trough is defined here as a “long depression generally wide and flat bottomed with symmetrical and parallel sides, and basin is defined a “depression more or less equidimensional in plan and of variable extent” as per the definitions in IHO (2008). Despite their names, features such as the Chagos Trough (~5,900 m, Raghukumar et al. 2004) are not considered here as they do not reach hadal depths. Of the Indian Ocean’s hadal features, the Wharton Basin in the northeast is the largest in terms of total area > 6,000 m (Table 4, Figure 10; Weston et al. 2021). The significant hadal footprint of the Wharton Basin, across ~10 degrees of latitude, is comparable to the extensive hadal basins of the northwest Pacific (Jamieson et al. in review). Indeed, deep troughs and basins account for 65% of the total hadal area in the Indian Ocean (Table 4) yet remain understudied. They could provide complex connectivity networks for hadal-obligate species, sustaining gene flow between isolated hadal features (Weston et al. 2021). For example, the eight north-south trending compartments of the Wharton Basin may act as dispersal corridors between the Java Trench and the Wallaby-Zenith and Diamantina Fracture Zones, with ~1.4 million km² of seafloor deeper than 5,600 m between the three features (Jacob et al. 2014, Weston et al. 2021). Given the range of different hadal features all within close proximity, including troughs, basins, fracture zones and a trench (Figure 10), the East Indian Ocean represents one of the most interesting opportunities to explore these connectivity patterns, globally (Jamieson et al. in review). In contrast, the deep troughs and basins of the central and western Indian Ocean are much smaller in area and, as with lesser-known fracture zones, require further investigation to verify their status as hadal features (Table 4). As greater recognition is given to their significance for hadal research

(Jamieson and Stewart 2021, Jamieson et al. in review), these aseismic hadal features should be prioritised for future research expeditions.

Journal Pre-proof

Table 4: Hadal features (> 6,000 m) of the Indian Ocean. We determine approximately 20 seafloor features with associated hadal areas in the Indian Ocean (see Figure 10). Maximum depths are derived from GEBCO_2023 bathymetry or from confirmed studies where definitive multibeam bathymetry is available. Areas were calculated using GEBCO data in QGIS.

Feature type	Feature name	Total hadal area (km ²)	Max. depth (m)		References
			GEBCO	Confirmed	
Subduction trenches	Java Trench	78,303	7,210	7,187	Bongiovanni et al. 2022, GEBCO_2023
	Total (trenches)	78,303 (28%)			
Fracture zones	Diamantina Fracture Zone	12,670	7,174	7,019	Bongiovanni et al. 2022, GEBCO_2023
	Wallaby-Zenith Fracture Zone	1,357	6,649	6,675	Niyazi et al. 2024, GEBCO_2023
	Vema Trench Fracture Zone	284	6,315	-	GEBCO_2023
	Melville Fracture Zone	722	6,653	-	GEBCO_2023
	Atlantis II Fracture Zone	490	6,450	-	GEBCO_2023
	Gazelle Fracture Zone	54	6,223	-	GEBCO_2023
	Gallieni Fracture Zone	477	6,374	-	GEBCO_2023
	Andrew Bain Fracture Zone	1,826	6,673	-	GEBCO_2023
	Total (fracture zones)	19,391 (7%)			
Troughs, basins and other features	Wharton Basin	174,430	7,371 [†]	-	GEBCO_2023
	Langseth Trough	164	6,607	-	GEBCO_2023
	Brouwer Trough	1,427	6,404	-	GEBCO_2023
	South Australian Basin	57	6,573	-	GEBCO_2023
	Madagascar Basin	844	6,766	-	GEBCO_2023
	Mozambique Basin	567	6,253	-	GEBCO_2023
	Agulhas Basin	1,091	6,325	-	GEBCO_2023
	Weddell Basin	79	6,091	-	GEBCO_2023
	Zenith Plateau*	209	6,088	-	GEBCO_2023
	Lost Dutchmen Ridge*	382	6,345	-	GEBCO_2023
	Dirck Hartog Ridge*	1,511	6,363	-	GEBCO_2023
	Total (basins)	179,250 (65%)			
	Total (all)	276,944			

*Note that, in the absence of definite names, these hadal areas are clustered under adjacent seafloor features and are not necessarily associated with the geological formation of these features. [†]Note that this max. depth is altimetrically derived and is unlikely to be the deepest point in the IO.




Figure 10: Hadal features of the Indian Ocean, with hadal depths > 6,000 m and hadal features marked by dashed lines. AB: Agulhas Basin, ABFZ: Andrew Bain Fracture Zone, AIFZ: Atlantis II Fracture Zone, BT: Brouwer Trough, DFZ: Diamantina Fracture Zone, DHR: Dirck Hartog Ridge, GFZI: Gazelle Fracture Zone, GFZII: Gallieni Fracture Zone, JT: Java Trench, LDR: Lost Dutchmen Ridge, LT: Langseth Trough, MAB: Madagascar Basin, MFZ: Melville Fracture Zone, MOB: Mozambique Basin, SAB: South Australian Basin, VTFZ: Vema Trench Fracture Zone, WB: Wharton Basin, WEB: Weddell Basin, WZFZ: Wallaby-Zenith Fracture Zone, ZP: Zenith Plateau.

Pelagic environments

It is widely recognised that global knowledge of the pelagic ocean, that is the mid-water environment between the ocean's surface and the seafloor, is extremely poor (Webb et al. 2010). To date, pelagic research in the Indian Ocean has been largely restricted to the upper 1,000 m, with most pelagic surveys occurring at epipelagic (0-200 m) and mesopelagic (200-1,000 m) depths (Vipin et al. 2011, Wafar et al. 2011, Obura 2015, Bernal et al. 2020a, 2020b). Similarly, existing demersal surveys for groundfish in the Indian Ocean are generally limited to depths < 1,000 m (Venu 2013, Aneesh Kumar et al. 2017). Moreover, a large proportion of the available pelagic Indian Ocean literature is focussed on the summits of shallow seamounts,

charismatic megafauna, or species targeted by fisheries (Vipin et al. 2011, Ebert 2013, Menard et al. 2013, Ebert 2014, Rogers 2017, Rogers et al. 2017, Cherel et al. 2020, Marsac et al. 2020, Roberts et al. 2020). Meanwhile, limited sampling of invertebrates and fish at all depths of the water column renders the systematic assessment of total deep pelagic biodiversity impossible (Obura 2015).

While seminal in the promotion and compilation of Indian Ocean research, the International Indian Ocean Expedition (IIOE, 1960-1965) primarily focussed on understanding biological oceanographic processes, primary productivity, and monsoon seasonality (Ryther 1963, Hood et al. 2016, Urban Jr. 2020). The Second IIOE (IIOE-2, 2015-2025) likewise aimed to advance understanding of oceanic, atmospheric, and geologic process and interactions in the Indian Ocean (Hood et al. 2015, Hood et al. 2016, see special issues for IIOE-2 research in Deep Sea Research Part II: Topical Studies in Oceanography – e.g., Hood and Beckley 2019 volume 1). Thus, while the IIOE programs provided a significant scientific foundation for our modern understanding of the Indian Ocean including insights into the pelagic realm (Zeitzschel and Gerlach 2012, Hood and Beckley 2019), they do not directly address the biological communities of the deep pelagic Indian Ocean.

Synthesis

Twenty years ago, Demopoulos et al. (2003) concluded that the deep Indian Ocean remains poorly known and is still waiting to be discovered and understood. Now, 20 years later, we review how deep-sea ecosystem research has advanced, synthesising our updated knowledge of deep-sea habitats and biodiversity in the Indian Ocean, and presenting outstanding problems for this field of research going forward.

Most deep-sea research in the Indian Ocean has been focused on certain ecosystems in specific regions. In the eastern Indian Ocean, the best characterised deep-sea ecosystems are the Australian continental margin especially its submarine canyons, the Java Trench, and other geologically complex features such as the Diamantina and Wallaby-Zenith fracture zones.

Conversely, the varying ecosystems of the Wharton and South Australian basins and Southeast Indian Ridge have scarcely been surveyed and require targeted future exploration. Similarly, the remote southern Indian Ocean, especially the Australian-Antarctic Basin, is thoroughly lacking in deep-sea research, save for limited surveying at the Kerguelen and Crozet plateaus.

Meanwhile, the primary deep-sea research focus in the western Indian Ocean has been the seamounts and hydrothermal vents of the Southwest Indian Ridge, as well some industry-driven research at the continental margins of eastern African countries such as Tanzania and Kenya. Despite this, there is a disparity in deep-sea research between the western and eastern Indian Ocean exemplified by the lack of hadal exploration for the western Indian Ocean's numerous fracture zones. Hydrothermal vent research continues northwards along the Central Indian Ridge and Carlsberg Ridge to the Arabian Sea, where existing research has focussed on the faunal communities and productivity patterns at the abyssal seafloor. Cold seeps and oxygen minimum zones dominate deep-sea research interest in the northern Indian Ocean, alongside the brine pools of the Red Sea, while the primary research target in the central Indian Ocean is the polymetallic nodule field of the Mid-Indian Ocean Basin. In contrast, the deep-sea ecosystems of major seafloor features such as the Mascarene Plateau, and Chagos-Laccadive and Ninetyeast ridges remain unexplored.

Inconsistencies over spatial scales

Spatial inconsistencies pose a significant hurdle to the advancement of deep-sea research in the Indian Ocean, both in research extent and effort. Of the countries bordering the Indian Ocean, India and Australia are leading the way for deep-sea research within EEZs, likely owing to greater access to resources including equipment and ship time. Meanwhile exploration in ABNJ has generally been restricted to specific sites of interest such as seamounts and hydrothermal vents along the mid-ocean ridge systems and potential industry targets such as the Mid-Indian Ocean basin. There are also bathymetric inconsistencies in sampling effort whereby the majority of biological data collected are from shallower bathyal depths (Figure 6), a trend seen

across other global oceans and likely linked to the increase in resources and technology required for deeper surveys (Bridges et al. 2023). This bias towards shallower sampling limits our ability to gain a holistic understanding of the deep Indian Ocean system.

At a regional scale, a lack of consistency in the defining limits of the Indian Ocean means data from holistic Indian Ocean studies are not directly comparable. For example, a global study of ophiuroid diversity unconventionally delimited the Indian Ocean at $\sim 90^\circ$ E and $\sim 30^\circ$ S, such that records from the majority of the eastern Indian Ocean are attributed instead to the Indo-Pacific region while records from the southern Indian Ocean are split between the 'South Africa' and 'South Pacific' regions (Stöhr et al. 2012). A further study reviewing the state of knowledge of coastal and marine biodiversity in the Indian Ocean similarly considered only the northwestern region of the Indian Ocean (Wafar et al. 2011). Meanwhile, the Australian Hydrographic Office regards the south Australian margin, including the Great Australian Bight, as the northerly limit of the Southern Ocean rather than the internationally accepted 60° S.

To achieve a spatially comprehensive and accurate understanding of large-scale ecological and biodiversity patterns in the deep Indian Ocean requires:

- 1) wider access to technologies for deep-sea exploration within EEZs via capacity building initiatives and national collaborations (Gates et al. 2017, Woodall et al. 2021),
- 2) targeted research initiatives in ABNJ, especially regions that are known to be under-sampled (e.g., SEIR),
- 3) improved open-access data via the use of global databases like OBIS and the promotion of FAIR data principles (Klein et al. 2019, Levin et al. 2019), and
- 4) consistency in the use of terminology and defining limits used for ocean basin-wide studies to allow for regional and global comparisons.

Data gaps over temporal scales

At present, there is no targeted long-term monitoring of deep-sea ecosystems in the Indian Ocean. Deep Indian Ocean research is consequently lacking data over temporal scales whereby

existing discrete data are unlikely to capture any seasonal or annual shifts in the faunal communities and abiotic features of deep-sea habitats that may occur in response to natural events or anthropogenic impacts (Wafar et al. 2011). Some research programmes have conducted multi-year sampling to study such trends, including the INDEX project examining the environmental impacts following a simulated mining event in the Mid-Indian Ocean Basin (1997-2001; Ingole et al. 2005), and the BIGSET project's study of the coupling between monsoon-driven productivity cycles and benthic processes in the Arabian Sea (1995-1998; Lochte and Pfannkuche 2000). However, there remains a paucity of continually sustained monitoring and time-series studies essential for recording temporal change in deep-sea ecosystems to the degree seen in other ocean basins (Glover et al. 2010).

Elsewhere, long-term monitoring via observatories or repeated surveying have facilitated the collection of valuable time series datasets. The Porcupine Abyssal Plain Sustained Observatory (PAP) in the North Atlantic and Station M in the North Pacific represent two of the best examples of long-term observing in the deep ocean, with over 30 years of *in situ* data collected from abyssal depths (PAP: 4,850 m, Station M: ~4,000 m; Lampitt et al. 2010, Smith Jr. et al. 2017, Smith Jr. et al. 2020, Hartman et al. 2021). Meanwhile, cabled observatories such as the Monterey Accelerated Research System (MARS) provide the power and network accessibility of an onshore laboratory thus enabling long-term, real-time experiments and testing of new deep-sea equipment like the benthic rover (Henthorn et al. 2010), however they are often restricted to shallower depths (e.g., MARS: 891 m). Indeed, other than the Indian Ocean and South Pacific, some form of long-term multidisciplinary deep-sea monitoring has been undertaken via observatory systems in all other global oceans (Glover et al. 2010), including MABEL – Southern Ocean (Cenedese et al. 2004, Gerber and Clauss 2010), HAUSGARTEN – Arctic Ocean (Soltwedel et al. 2005, Bergmann et al. 2011), and DELOS – South Atlantic (Vardaro et al. 2013). The integration of long-term observatories and repeated surveying would foster a new era of deep-sea exploration and increase research capacity in the deep Indian Ocean.

Unexplored deep pelagic environments

In keeping with the available literature, much of this review is dedicated to the benthic habitats and communities of the Indian Ocean. The chronic under-exploration of the deep pelagic ocean and lack of a biodiversity baseline for this environment leaves a substantial gap in our knowledge of the deep Indian Ocean (Robison 2009, Webb et al. 2010). Establishing a biodiversity baseline for the Indian Ocean's (and indeed the Earth's) largest habitat by volume is essential to identify the regional patterns in community composition and the main environmental or ecological drivers across different depth gradients (Haddock and Choy 2024). As such, a greater focus is required for the mid-water communities of bathypelagic, abyssopelagic and hadal depths (Webb et al. 2010), as well as the mesopelagic zone (Haddock and Choy 2024), to fully catalogue and understand the deep-sea biodiversity patterns of the Indian Ocean. The recent drive to prioritise deep pelagic research and mid-water ecology, as well as the integration of new sampling techniques such as eDNA, has resulted in knowledge advancements for biodiversity distribution and population monitoring in the Pacific and Atlantic Oceans (Li et al. 2022). Similar effort should now be given to prioritise deep pelagic research in the Indian Ocean.

Placing the deep Indian Ocean in a global context

The final consideration for the future promotion of deep-sea research in the Indian Ocean is the need to consider the Indian Ocean in a global context. The Indian Ocean is too often overlooked in global studies or disregarded for the supposition of its lack of data. While there are certainly data limitations, as summarised above, this review highlights that the Indian Ocean's deep-sea ecosystems are better-known than commonly considered. Furthermore, studying the Indian Ocean in isolation neglects the interconnectivity of global ocean processes and faunal biogeography. For example, several studies have demonstrated the importance of the Indian Ocean as a potential connectivity pathway for hydrothermal vent fauna between the Atlantic and Pacific Oceans, whether as a dispersal corridor supporting modern gene flow (German et al.

1998, Hwang et al. 2022) or as an evolutionary corridor creating a melting pot of Atlantic and western Pacific vent species in the Indian Ocean (Perez et al. 2021). Likewise, on a global scale, the Indian Ocean is often considered a single biogeographic province (e.g., lower bathyal province BY11 and abyssal province AB8 – Watling et al. 2013; hydrothermal vents – Rogers et al. 2012). However, recent studies have suggested multiple biogeographic subdivisions for hydrothermal vent fauna, effectively dividing the Indian Ocean's ridges into four subregions (Table 3; Zhou et al. 2022, Jang et al. 2023). Future research should consider whether similar biogeographic patterns arise for other deep-sea ecosystems in the Indian Ocean. Greater recognition of the regional and local heterogeneity of the Indian Ocean's deep-sea ecosystems, as well as the uniqueness of its abiotic characteristics, would highlight its importance for understanding global ocean processes thus promoting further research interest and prioritisation in these areas.

As with our recommendations for improving data inconsistencies over spatial scales, newly collected data should be published in open-access data repositories so that they are readily available for use in future global studies (Klein et al. 2019). This need is exemplified by the comparative lack of OBIS records for the deep Indian Ocean (> 1,000 m), with ~5-16 times the number of records per 1 million km² available from other global oceans, pre-quality checks and data cleaning (Appendix 2; <https://mapper.obis.org/>, accessed 03/04/2024). Ensuring these data are available for future studies will provide greater international recognition of the significant deep-sea research effort that has arisen from Indian Ocean exploration to date. Utilising this existing research in conjunction with global oceanographic and geomorphic patterns will provide the background to ensure the deep Indian Ocean is not overlooked nor misrepresented on a global stage.

Conclusion

We provide the most comprehensive review of deep-sea ecosystem research in the Indian Ocean to date. By recognising and championing the deep-sea research effort undertaken thus

far, this review provides the necessary platform to inform future studies and ensure the Indian Ocean is prioritised for deep-sea exploration. Addressing the knowledge gaps and outstanding problems identified by this review will foster the advancement of deep-sea research in the Indian Ocean during the UN Ocean Decade and beyond.

Journal Pre-proof

Acknowledgements

This review was supported by the Minderoo-UWA Deep-Sea Research Centre, funded by the Minderoo Foundation and Inkfish. We thank the two anonymous reviewers for their helpful feedback to improve this manuscript.

Author Contributions

EAT: Conceptualization, Investigation, Visualization, Writing – original draft, Writing – review & editing, Project administration. **TB:** Writing – original draft, Writing – review & editing. **JLK:** Writing – original draft, Writing – review & editing. **YN:** Visualization, Writing – original draft, Writing – review & editing. **DJBS:** Writing – original draft, Writing – review & editing. **AJJ:** Conceptualization, Writing – review & editing, Supervision.

Journal Pre-proof

References

- Ahusan, M., Rico-Seijo, N., Amjad, F., Gress, E., Naeem, S., Samaai, T., Samimi-Namin, K., Woodall, L., Stefanoudis, P., 2023. The Nekton Maldives taxonomic workshop: Exploring the biodiversity of shallow, mesophotic and deep-sea communities in Maldives. *Research Ideas and Outcomes* 9, e114370. <https://doi.org/10.3897/rio.9.e114370>
- Ali, J.R., Aitchison, J.C., 2005. Greater India. *Earth-Science Reviews* 72, 169–188. <https://doi.org/10.1016/j.earscirev.2005.07.005>
- Amon, D.J., Copley, J.T., Dahlgren, T.G., Horton, T., Kemp, K.M., Rogers, A.D., Glover, A.G., 2017. Observations of fauna attending wood and bone deployments from two seamounts on the Southwest Indian Ridge. *Deep Sea Research Part II: Topical Studies in Oceanography, The Pelagic Ecology of Seamounts of the South West Indian Ocean* 136, 122–132. <https://doi.org/10.1016/j.dsr2.2015.07.003>
- Aneesh Kumar, K.V., Nikki, R., Oxona, K., Hashim, M., Sudhakar, M., 2017. Relationships between fish and otolith size of nine deep-sea fishes from the Andaman and Nicobar waters, North Indian Ocean. *Journal of Applied Ichthyology* 33, 1187–1195. <https://doi.org/10.1111/jai.13467>
- Ansari, Z.A., 2000. Distribution of Deep-Sea Benthos in the Proposed Mining Area of Central Indian Basin. *Marine Georesources & Geotechnology* 18, 201–207. <https://doi.org/10.1080/10641190009353788>
- Antunes, A., Ngugi, D.K., Stingl, U., 2011. Microbiology of the Red Sea (and other) deep-sea anoxic brine lakes. *Environmental Microbiology Reports* 3, 416–433. <https://doi.org/10.1111/j.1758-2229.2011.00264.x>
- Atmanand, M.A., 2019. Ocean Technology Capacity Building in India. *J Geol Soc India* 94, 447–452. <https://doi.org/10.1007/s12594-019-1340-4>
- Bachraty, C., Legendre, P., Desbruyères, D., 2009. Biogeographic relationships among deep-sea hydrothermal vent faunas at global scale. *Deep Sea Research Part I: Oceanographic Research Papers* 56, 1371–1378. <https://doi.org/10.1016/j.dsr.2009.01.009>
- Baco, A.R., 2007. Exploration for Deep-Sea Corals on North Pacific Seamounts and Islands. *Oceanography* 20, 108–117.
- Bailey, D.M., King, N.J., Priede, I.G., 2007. Cameras and carcasses: historical and current methods for using artificial food falls to study deep-water animals. *Marine Ecology Progress Series* 350, 179–192.
- Baker, M.C., Ramirez-Llodra, E.Z., Tyler, P.A., German, C.R., Boetius, A., Cordes, E.E., Dubilier, N., Fisher, C.R., Levin, L.A., Metaxas, A., Rowden, A.A., Santos, R.S., Shank, T.M., Van Dover, C.L., Young, C.M., Warén, A., 2010. Biogeography, Ecology, and Vulnerability of Chemosynthetic Ecosystems in the Deep Sea, in: *Life in the World's Oceans*. John Wiley & Sons, Ltd, pp. 161–182. <https://doi.org/10.1002/9781444325508.ch9>
- Barrett, J., Chase, Z., Zhang, J., Holl, M.M.B., Willis, K., Williams, A., Hardesty, B.D., Wilcox, C., 2020. Microplastic Pollution in Deep-Sea Sediments From the Great Australian Bight. *Frontiers in Marine Science* 7.
- Batang, Z.B., Papathanassiou, E., Al-Suwailem, A., Smith, C., Salomidi, M., Petihakis, G., Alikunhi, N.M., Smith, L., Mallon, F., Yapici, T., Fayad, N., 2012. First discovery of a cold seep on the

continental margin of the central Red Sea. *Journal of Marine Systems* 94, 247–253.

<https://doi.org/10.1016/j.jmarsys.2011.12.004>

Baumann, A., Richter, H., Schoell, M., 1973. Suakin deep: Brines and hydrothermal sediments in the deepest part of the Red Sea. *Geol Rundsch* 62, 684–697.

<https://doi.org/10.1007/BF01820956>

Beaulieu, S.E., Szafranski, K., 2020. InterRidge Global Database of Active Submarine Hydrothermal Vent Fields, Version 3.4. [WWW Document]. URL <http://vents-data.interridge.org> (accessed 2.12.22).

Begg, G.A., 2018. The Great Australian Bight Research Program. Deep Sea Research Part II: Topical Studies in Oceanography, Great Australian Bight Research Program - a whole of system investigation 157–158, 1–2. <https://doi.org/10.1016/j.dsr2.2018.11.017>

Belkin, I.M., Gordon, A., 1996. Southern Ocean fronts from the Greenwich meridian to Tasmania. *Journal of Geophysical Research C: Oceans* 101, 3675–3696.

Bell, K.L.C., Quinzin, M.C., Amon, D., Poulton, S., Hope, A., Sarti, O., Cañete, T.E., Smith, A.M., Baldwin, H.I., Lira, D.M., Cambroner-Solano, S., Chung, T.-R.A., Brady, B., 2023. Exposing inequities in deep-sea exploration and research: results of the 2022 Global Deep-Sea Capacity Assessment. *Frontiers in Marine Science* 10.

Bénard, F., Callot, J.P., Vially, R., Schmitz, J., Roest, W., Patriat, M., Loubrieu, B., 2010a. The Kerguelen plateau: Records from a long-living/composite microcontinent. *Marine and Petroleum Geology* 27, 633–649. <https://doi.org/10.1016/j.MARPETGEO.2009.08.011>

Bénard, F., Callot, J.-P., Vially, R., Schmitz, J., Roest, W., Patriat, M., Loubrieu, B., 2010b. The Kerguelen plateau: Records from a long-living/composite microcontinent. *Marine and Petroleum Geology, The link between shallow and deep processes in sedimentary basins* 27, 633–649. <https://doi.org/10.1016/j.marpetgeo.2009.08.011>

Bergmann, M., Soltwedel, T., Klages, M., 2011. The interannual variability of megafaunal assemblages in the Arctic deep sea: Preliminary results from the HAUSGARTEN observatory (79°N). *Deep Sea Research Part I: Oceanographic Research Papers* 58, 711–723. <https://doi.org/10.1016/j.dsr.2011.03.007>

Bergstad, O.A., Falkenhaug, T., Astthorsson, O.S., Byrkjedal, I., Gebruk, A.V., Piatkowski, U., Priede, I.G., Santos, R.S., Vecchione, M., Lorange, P., Gordon, J.D.M., 2008. Towards improved understanding of the diversity and abundance patterns of the mid-ocean ridge macro- and megafauna. *Deep Sea Research Part II: Topical Studies in Oceanography, Mid-Atlantic Ridge Habitats and Biodiversity* 55, 1–5. <https://doi.org/10.1016/j.dsr2.2007.10.001>

Bernal, A., Olivar, M.P., Beckley, L.E., 2020a. Dietary composition of myctophid larvae off Western Australia. *Deep Sea Research Part II: Topical Studies in Oceanography, The Second International Indian Ocean Expedition (IIOE-2): Motivating New Exploration in a Poorly Understood Ocean Basin Volume 3* 179, 104841. <https://doi.org/10.1016/j.dsr2.2020.104841>

Bernal, A., Toresen, R., Riera, R., 2020b. Mesopelagic fish composition and diets of three myctophid species with potential incidence of microplastics, across the southern tropical gyre. *Deep Sea Research Part II: Topical Studies in Oceanography, The Second International Indian Ocean Expedition (IIOE-2): Motivating New Exploration in a Poorly Understood Ocean Basin Volume 3* 179, 104706. <https://doi.org/10.1016/j.dsr2.2019.104706>

- Bernardino, A.F., Smith, C.R., Baco, A., Altamira, I., Sumida, P.Y.G., 2010. Macrofaunal succession in sediments around kelp and wood falls in the deep NE Pacific and community overlap with other reducing habitats. *Deep Sea Research Part I: Oceanographic Research Papers* 57, 708–723. <https://doi.org/10.1016/j.dsr.2010.03.004>
- Bezrunov, P.L., 1963. Research in the Indian Ocean by the survey vessel *Vitiaz* on its thirty-third voyage. *Deep Sea Research and Oceanographic Abstracts* 10, 59–66. [https://doi.org/10.1016/0011-7471\(63\)90186-4](https://doi.org/10.1016/0011-7471(63)90186-4)
- Bienhold, C., Ristova, P.P., Wenzhöfer, F., Dittmar, T., Boetius, A., 2013. How Deep-Sea Wood Falls Sustain Chemosynthetic Life. *PLOS ONE* 8, e53590. <https://doi.org/10.1371/journal.pone.0053590>
- Billett, D.S.M., Bett, B.J., Jacobs, C.L., Rouse, I.P., Wigham, B.D., 2006. Mass deposition of jellyfish in the deep Arabian Sea. *Limnology and Oceanography* 51, 2077–2083. <https://doi.org/10.4319/lo.2006.51.5.2077>
- Blain, S., Quéguiner, B., Armand, L., Belviso, S., Bombled, B., Bopp, L., Bowie, A., Brunet, C., Brussaard, C., Carlotti, F., Christaki, U., Corbière, A., Durand, I., Ebersbach, F., Fuda, J.L., Garcia, N., Gerringa, L., Griffiths, B., Guigue, C., Guillermin, C., Jacquet, S., Jeandel, C., Laan, P., Lefèvre, D., Lo Monaco, C., Malits, A., Mosseri, J., Obernosterer, I., Park, Y.H., Picheral, M., Pondaven, P., Remenyi, T., Sandroni, V., Sarthou, G., Savoye, N., Scouarnec, L., Souhaut, M., Thuiller, D., Timmermans, K., Trull, T., Uitz, J., Van Beek, P., Veldhuis, M., Vincent, D., Viollier, E., Vong, L., Wagener, T., 2007. Effect of natural iron fertilization on carbon sequestration in the Southern Ocean. *Nature* 446, 1070–1074. <https://doi.org/10.1038/NATURE05700>
- Blankenship-Williams, L.E., Levin, L.A., 2009. Living Deep: A Synopsis of Hadal Trench Ecology. *Marine Technology Society Journal* 43, 137–143. <https://doi.org/10.4031/MTSJ.43.5.23>
- Bond, T., Jamieson, A., 2022. The extent and protection of Australia's deep sea. *Mar. Freshwater Res.* 73, 1520–1526. <https://doi.org/10.1071/MF22156>
- Bond, T., Niyazi, Y., Kolbusz, J.L., Jamieson, A.J., 2023. Habitat and benthic fauna of the Wallaby-Cuvier escarpment, SE Indian ocean. *Deep Sea Research Part II: Topical Studies in Oceanography* 210, 105299. <https://doi.org/10.1016/j.dsr2.2023.105299>
- Bongiovanni, C., Stewart, H.A., Jamieson, A.J., 2022. High-resolution multibeam sonar bathymetry of the deepest place in each ocean. *Geoscience Data Journal* 9, 108–123. <https://doi.org/10.1002/gdj3.122>
- Bridges, A.E.H., Howell, K.L., Amaro, T., Atkinson, L., Barnes, D.K.A., Bax, N., Bell, J.B., Bernardino, A.F., Beuck, L., Braga-Henriques, A., Brandt, A., Bravo, M.E., Brix, S., Butt, S., Carranza, A., Doti, B.L., Elegbede, I.O., Esquete, P., Freiwald, A., Gaudron, S.M., Guilhon, M., Hebbeln, D., Horton, T., Kainge, P., Kaiser, S., Lauretta, D., Limongi, P., McQuaid, K.A., Milligan, R.J., Miloslavich, P., Narayanaswamy, B.E., Orejas, C., Paulus, S., Pearman, T.R.R., Perez, J.A., Ross, R.E., Saeedi, H., Shimabukuro, M., Sink, K., Stevenson, A., Taylor, A., Titschack, J., Vieira, R.P., Vinha, B., Wienberg, C., 2023. Review of the Central and South Atlantic offshore and deep-sea benthos: Science, policy and management, in: Hawkins, S.J., Todd, P.A., Russell, B.D. (Eds.), *Oceanography and Marine Biology: An Annual Review*. CRC Press, Taylor & Francis Group, Boca Raton, pp. 127–218.
- Bristow, L.A., Callbeck, C.M., Larsen, M., Altabet, M.A., Dekaezemacker, J., Forth, M., Gauns, M., Glud, R.N., Kuypers, M.M.M., Lavik, G., Milucka, J., Naqvi, S.W.A., Pratihary, A., Revsbech, N.P., Thamdrup, B., Treusch, A.H., Canfield, D.E., 2017. N₂ production rates limited by nitrite

availability in the Bay of Bengal oxygen minimum zone. *Nature Geosci* 10, 24–29.
<https://doi.org/10.1038/ngeo2847>

Budnichenko, V.A., Gubanov, E.P., Demidov, V.F., Ivanin, N.A., Romanov, A.V., Romanov, E.V., Pinchukov, M.A., Piotrovsky, A.S., Prutko, V.G., 1997. Results of YugNIRO studies of fish resources in the Indian and Southern Oceans, Результаты изучения ЮгНИРО сырьевых ресурсов в Индийском и Южном океанах. *Proceedings of the Southern Scientific Research Institute of Marine Fisheries and Oceanography* 43, 28–47.

Cai, W., Cowan, T., Raupach, M., 2009. Positive Indian Ocean Dipole events precondition southeast Australia bushfires. *Geophysical Research Letters* 36.
<https://doi.org/10.1029/2009GL039902>

Caladan Oceanic, 2020. Caladan Oceanic Completes First Mapping of Key Deep-Sea Features in Indian Ocean.

Carroll, E., Patenaude, N., Alexander, A., Steel, D., Harcourt, R., Childerhouse, S., Smith, S., Bannister, J., Constantine, R., Baker, C.S., 2011. Population structure and individual movement of southern right whales around New Zealand and Australia. *Marine Ecology Progress Series* 432, 257–268. <https://doi.org/10.3354/meps09145>

Cenedese, S., Gasparoni, F., Calcara, M., D'Anna, G., Favali, P., Evers, K.-U., 2004. MABEL – The First Seafloor Observatory For Multidisciplinary Long-term Monitoring In Polar Environment. Presented at the The Fourteenth International Offshore and Polar Engineering Conference, OnePetro.

Ceramicola, S., Dupré, S., Somoza, L., Woodside, J., 2018. Cold Seep Systems, in: Micallef, A., Krastel, S., Savini, A. (Eds.), *Submarine Geomorphology*, Springer Geology. Springer International Publishing, Cham, pp. 367–387. https://doi.org/10.1007/978-3-319-57852-1_19

Chambault, P., Fossette, S., Heide-Jørgensen, M.P., Jouannet, D., Vély, M., 2021. Predicting seasonal movements and distribution of the sperm whale using machine learning algorithms. *Ecology and Evolution* 11, 1432–1445. <https://doi.org/10.1002/ece3.7154>

Chen, C., Linse, K., Copley, J.T., Rogers, A.D., 2015. The ‘scaly-foot gastropod’: a new genus and species of hydrothermal vent-endemic gastropod (Neomphalina: Peltospiridae) from the Indian Ocean. *Journal of Molluscan Studies* 81, 322–334. <https://doi.org/10.1093/mollus/eyv013>

Chen, C., Zhou, Y., Wang, C., Copley, J.T., 2017. Two New Hot-Vent Peltospirid Snails (Gastropoda: Neomphalina) from Longqi Hydrothermal Field, Southwest Indian Ridge. *Frontiers in Marine Science* 4.

Cherel, Y., Duhamel, G., Gasco, N., 2004. Cephalopod fauna of subantarctic islands: new information from predators. *Marine Ecology Progress Series* 266, 143–156.

Cherel, Y., Romanov, E.V., Annasawmy, P., Thibault, D., Ménard, F., 2020. Micronektonic fish species over three seamounts in the southwestern Indian Ocean. *Deep Sea Research Part II: Topical Studies in Oceanography, Bio-Physical coupling around three shallow seamounts in the South Western Indian Ocean, with regional comparisons based on modelling, remote sensing and observational studies* 176, 104777. <https://doi.org/10.1016/j.dsr2.2020.104777>

Christiansen, B., Boetius, A., 2000. Mass sedimentation of the swimming crab *Charybdis smithii* (Crustacea: Decapoda) in the deep Arabian Sea. *Deep Sea Research Part II: Topical Studies in Oceanography* 47, 2673–2685. [https://doi.org/10.1016/S0967-0645\(00\)00044-8](https://doi.org/10.1016/S0967-0645(00)00044-8)

- Clark, M.R., Rowden, A.A., Schlacher, T., Williams, A., Consalvey, M., Stocks, K.I., Rogers, A.D., O'Hara, T.D., White, M., Shank, T.M., Hall-Spencer, J.M., 2010. The Ecology of Seamounts: Structure, Function, and Human Impacts. *Annual Review of Marine Science* 2, 253–278. <https://doi.org/10.1146/annurev-marine-120308-081109>
- Clark, M.R., Vinnichenko, V.I., Gordon, J.D.M., Beck-Bulat, G.Z., Kukharev, N.N., Kakora, A.F., 2007. Large-Scale Distant-Water Trawl Fisheries on Seamounts, in: *Seamounts: Ecology, Fisheries & Conservation*. John Wiley & Sons, Ltd, pp. 361–399. <https://doi.org/10.1002/9780470691953.ch17>
- Clift, P.D., Shimizu, N., Layne, G.D., Blusztajn, J.S., Gaedicke, C., Schlüter, H.-U., Clark, M.K., Amjad, S., 2001. Development of the Indus Fan and its significance for the erosional history of the Western Himalaya and Karakoram. *GSA Bulletin* 113, 1039–1051. [https://doi.org/10.1130/0016-7606\(2001\)113<1039:DOTIFA>2.0.CO;2](https://doi.org/10.1130/0016-7606(2001)113<1039:DOTIFA>2.0.CO;2)
- Conlan, K.E., Currie, D.R., Dittmann, S., Sorokin, S.J., Hendrycks, E., 2015. Macrofaunal Patterns in and around the Couedic and Bonney Submarine Canyons, South Australia. *PLOS ONE* 10, e0143921. <https://doi.org/10.1371/journal.pone.0143921>
- Consalvey, M., Clark, M.R., Rowden, A.A., Stocks, K.I., 2010. Life on Seamounts, in: *Life in the World's Oceans*. John Wiley & Sons, Ltd, pp. 123–138. <https://doi.org/10.1002/9781444325508.ch7>
- Copley, J.T., Marsh, L., Glover, A.G., Hühnerbach, V., Nye, V.E., Reid, W.D.K., Sweeting, C.J., Wigham, B.D., Wiklund, H., 2016. Ecology and biogeography of megafauna and macrofauna at the first known deep-sea hydrothermal vents on the ultraslow-spreading Southwest Indian Ridge. *Sci Rep* 6, 39158. <https://doi.org/10.1038/srep39158>
- Cousins, N., Horton, T., Wigham, B., Bagley, P., 2013. Abyssal scavenging demersal fauna at two areas of contrasting productivity on the Subantarctic Crozet Plateau, southern Indian Ocean. *African Journal of Marine Science* 35, 299–306. <https://doi.org/10.2989/1814232X.2013.802747>
- Cousins, N.J., Priede, I.G., 2012. Abyssal demersal fish fauna composition in two contrasting productivity regions of the Crozet Plateau, Southern Indian Ocean. *Deep Sea Research Part I: Oceanographic Research Papers* 64, 71–77. <https://doi.org/10.1016/J.DSR.2012.02.003>
- Cross, I.A., Gebruk, A., Rogacheva, A., Billett, D.S.M., 2009. *Peniagone crozeti* a new species of elasipodid holothurian from abyssal depths off the Crozet Islands in the Southern Indian Ocean. *Zootaxa* 2096, 484–488. <https://doi.org/10.11646/zootaxa.2096.1.30>
- Curry, J.R., Emmel, F.J., Moore, D.G., 2003. The Bengal Fan: morphology, geometry, stratigraphy, history and processes. *Marine and Petroleum Geology* 19, 1191–1223. [https://doi.org/10.1016/S0264-8172\(03\)00035-7](https://doi.org/10.1016/S0264-8172(03)00035-7)
- Currie, D.R., Sorokin, S.J., 2014. Megabenthic biodiversity in two contrasting submarine canyons on Australia's southern continental margin. *Marine Biology Research* 10, 97–110. <https://doi.org/10.1080/17451000.2013.797586>
- Damuth, J.E., Johnson, D.A., 1989. Morphology, sediments and structure of the Amirante Trench, Western Indian Ocean: implications for trench origin. *Marine and Petroleum Geology* 6, 232–242. [https://doi.org/10.1016/0264-8172\(89\)90003-2](https://doi.org/10.1016/0264-8172(89)90003-2)

Daniell, J., Jorgensen, D., Anderson, T., Borissova, I., Burq, S., Heap, A., Hughes, M., Mantle, D., Nelson, G., Nichol, S., Nicholson, C., Payne, D., Przeslawski, R., Radke, L., Siwabessy, P.J., Smith, C., Party, S., 2009. Frontier Basins of the West Australian Continental Margin: Post-survey Report of Marine Reconnaissance and Geological Sampling Survey GA2476, Geosci. Aust., Rec. Geosci. Aust.

Daniell, J., Jorgensen, D.C., Anderson, T., Borissova, I., Burq, S., Heap, A.D., Hughes, M., Mantle, D., Nelson, G., Nichol, S., Nicholson, C., Payne, D., Przeslawski, R., Radke, L., Siwabessy, J., Smith, C., Party, S., 2010. Frontier basins of the West Australian continental margin: post-survey report of marine reconnaissance and geological sampling survey GA2476, RV Sonne, October 2008-January 2009 229.

Das, P., Iyer, S.D., Kodagali, V.N., 2007. Morphological characteristics and emplacement mechanism of the seamounts in the Central Indian Ocean Basin. *Tectonophysics* 443, 1–18. <https://doi.org/10.1016/j.tecto.2007.08.002>

Davis, J.K., Lawver, L.A., Norton, I.O., Gahagan, L.M., 2016. New Somali Basin magnetic anomalies and a plate model for the early Indian Ocean. *Gondwana Research* 34, 16–28. <https://doi.org/10.1016/j.gr.2016.02.010>

Dawson, M.N., 2016. Island and island-like marine environments. *Global Ecology and Biogeography* 25, 831–846. <https://doi.org/10.1111/geb.12314>

De Deckker, P., 2016. The Indo-Pacific Warm Pool: critical to world oceanography and world climate. *Geoscience Letters* 3, 20. <https://doi.org/10.1186/s40562-016-0054-3>

De Leo, F.C., Smith, C.R., Rowden, A.A., Bowden, D.A., Clark, M.R., 2010. Submarine canyons: hotspots of benthic biomass and productivity in the deep sea. *Proceedings of the Royal Society B: Biological Sciences* 277, 2783–2792. <https://doi.org/10.1098/rspb.2010.0462>

Demopoulos, A.W.J., Smith, C.R., Tyler, P.A., 2003. The Deep Indian Ocean Floor, in: *Ecosystems of the Deep Oceans*. Elsevier, pp. 219–237.

Devey, C.W., Augustin, N., Brandt, A., Brenke, N., Köhler, J., Lins, L., Schmidt, C., Yeo, I.A., 2018. Habitat characterization of the Vema Fracture Zone and Puerto Rico Trench. *Deep Sea Research Part II: Topical Studies in Oceanography, Bathymetry of the Vema-Fracture Zone and Puerto Rico Trench Abyssal Atlantic Biodiversity Study (Vema-TRANSIT)* 148, 7–20. <https://doi.org/10.1016/j.dsr2.2018.02.003>

Dewangan, P., Sriram, G., Kumar, A., Mazumdar, A., Peketi, A., Mahale, V., Reddy, S.S.C., Babu, A., 2021. Widespread occurrence of methane seeps in deep-water regions of Krishna-Godavari basin, Bay of Bengal. *Marine and Petroleum Geology* 124, 104783. <https://doi.org/10.1016/j.marpetgeo.2020.104783>

Dhawaskar, P., Ganguly, S., Mukhopadhyay, R., Manikyamba, C., Iyer, S.D., Karisiddaiah, S.M., Mahender, K., 2020. Geochemical heterogeneity along the Vema Fracture Zone, Indian Ocean: Mixing of melts from the Reunion plume and the Central Indian Ridge. *Geological Journal* 55, 330–343. <https://doi.org/10.1002/gj.3395>

Dick, H.J.B., Lin, J., Schouten, H., 2003. An ultraslow-spreading class of ocean ridge. *Nature* 426, 405–412. <https://doi.org/10.1038/nature02128>

Djurhuus, A., Read, J.F., Rogers, A.D., 2017. The spatial distribution of particulate organic carbon and microorganisms on seamounts of the South West Indian Ridge. *Deep Sea Research Part II*:

Topical Studies in Oceanography, The Pelagic Ecology of Seamounts of the South West Indian Ocean 136, 73–84. <https://doi.org/10.1016/j.dsr2.2015.11.015>

Douglas, C., 2021. Diel and Depth Patterns in Fish and Invertebrates Along a Subsea Pipeline (Masters). The University of Western Australia.

Duhamel, G., Williams, R., 2011. History of whaling, sealing, fishery and aquaculture trials in the area of the Kerguelen Plateau, in: The Kerguelen Plateau: Marine Ecosystem and Fisheries. pp. 15–28. <https://doi.org/10.13140/2.1.2059.7446>

Duperron, S., Gaudron, S.M., Rodrigues, C.F., Cunha, M.R., Decker, C., Olu, K., 2013. An overview of chemosynthetic symbioses in bivalves from the North Atlantic and Mediterranean Sea. Biogeosciences 10, 3241–3267. <https://doi.org/10.5194/bg-10-3241-2013>

Durden, J.M., Bett, B.J., Jones, D.O.B., Huvenne, V.A.I., Ruhl, H.A., 2015. Abyssal hills – hidden source of increased habitat heterogeneity, benthic megafaunal biomass and diversity in the deep sea. Progress in Oceanography 137, 209–218. <https://doi.org/10.1016/j.pocean.2015.06.006>

Ebbe, B., Billett, D., Brandt, A., Ellingsen, K., Glover, A., Keller, S., Malyutina, M., Mart, P., Arbizu, N., Molodtsova, T., Rex, M., Smith, C., Tselepidis, A., 2010. Diversity of Abyssal Marine Life, in: Life in the World's Oceans: Diversity, Distribution, and Abundance. pp. 139–160. <https://doi.org/10.1002/9781444325508.ch8>

Ebert, D.A., 2014. DEEP-SEA CARTILAGINOUS FISHES OF THE INDIAN OCEAN. FOOD AND AGRICULTURE ORGANIZATION OF THE UNITED NATIONS, Rome.

Ebert, D.A., 2013. DEEP-SEA CARTILAGINOUS FISHES OF THE INDIAN OCEAN. FOOD AND AGRICULTURE ORGANIZATION OF THE UNITED NATIONS, Rome.

Enge, A.J., Witte, U., Kucera, M., Heinz, P., 2014. Uptake of phytodetritus by benthic foraminifera under oxygen depletion at the Indian margin (Arabian Sea). Biogeosciences 11, 2017–2026. <https://doi.org/10.5194/bg-11-2017-2014>

Exon, N.F., Raven, M.D., Carlo, E.H.D., 2002. Ferromanganese Nodules and Crusts from the Christmas Island Region, Indian Ocean. Marine Georesources & Geotechnology 20, 275–297. <https://doi.org/10.1080/03608860290051958>

Farrelly, C., Ah Yong, S., 2019. Deepwater decapod, stomatopod and lophogastrid Crustacea from Eastern Australia and the Great Australian Bight collected in 2015–2017: preliminary identifications of 191 species. Museum Victoria Science Reports 21, 1–97. <https://doi.org/10.24199/j.mvsr.2019.21>

Faure, J., Gasco, N., Bonillo, C., Munaron, J.M., Cherel, Y., Péron, C., 2023. Feeding ecology of two deep-sea skates bycaught on demersal longlines off Kerguelen Islands, Southern Indian Ocean. Deep-Sea Research Part I: Oceanographic Research Papers 194, 103980. <https://doi.org/10.1016/j.dsr.2023.103980>

Feng, M., Benthuyssen, J., Zhang, N., Slawinski, D., 2015. Freshening anomalies in the Indonesian throughflow and impacts on the Leeuwin Current during 2010–2011. Geophysical Research Letters 42, 8555–8562. <https://doi.org/10.1002/2015GL065848>

Fischer, D., Sahling, H., Nöthen, K., Bohrmann, G., Zabel, M., Kasten, S., 2012. Interaction between hydrocarbon seepage, chemosynthetic communities, and bottom water redox at cold seeps of

the Makran accretionary prism: insights from habitat-specific pore water sampling and modeling. *Biogeosciences* 9, 2013–2031. <https://doi.org/10.5194/bg-9-2013-2012>

FISHER, R.L., JOHNSON, G.L., HEEZEN, B.C., 1967. Mascarene Plateau, Western Indian Ocean. *GSA Bulletin* 78, 1247–1266. [https://doi.org/10.1130/0016-7606\(1967\)78\[1247:MPWIO\]2.0.CO;2](https://doi.org/10.1130/0016-7606(1967)78[1247:MPWIO]2.0.CO;2)

FISHER, R.L., SCLATER, J.G., MCKENZIE, D.P., 1971. Evolution of the Central Indian Ridge, Western Indian Ocean. *GSA Bulletin* 82, 553–562. [https://doi.org/10.1130/0016-7606\(1971\)82\[553:EOTCIR\]2.0.CO;2](https://doi.org/10.1130/0016-7606(1971)82[553:EOTCIR]2.0.CO;2)

Flanders Marine Institute, 2023. Maritime Boundaries Geodatabase: Maritime Boundaries and Exclusive Economic Zones (200NM).

Forster, G.R., Badcock, J.R., Longbottom, M.R., Merrett, N.R., Thomson, K.S., Fraser, F.C., 1970. Results of the Royal Society Indian Ocean Deep Slope Fishing Expedition, 1969. *Proceedings of the Royal Society of London. Series B. Biological Sciences* 175, 367–404. <https://doi.org/10.1098/rspb.1970.0028>

Fossette, S., Heide-Jørgensen, M.-P., Jensen, M.V., Kiszka, J., Bérubé, M., Bertrand, N., Vély, M., 2014. Humpback whale (*Megaptera novaeangliae*) post breeding dispersal and southward migration in the western Indian Ocean. *Journal of Experimental Marine Biology and Ecology, Charismatic marine mega-fauna* 450, 6–14. <https://doi.org/10.1016/j.jembe.2013.10.014>

Foucher, J.-P., Westbrook, G.K., Boetius, A., Ceramicola, S., Dupré, S., Mascle, J., Mienert, J., Pfannkuche, O., Pierre, C., Praeg, D., 2009. Structure and Drivers of Cold Seep Ecosystems. *Oceanography* 22, 92–109.

Fournier, M., Chamot-Rooke, N., Petit, C., Huchon, P., Al-Kathiri, A., Audin, L., Beslier, M.-O., d'Acremont, E., Fabbri, O., Fleury, J.-M., Khanbari, K., Lepvrier, C., Leroy, S., Maillot, B., Merkouriev, S., 2010. Arabia-Somalia plate kinematics, evolution of the Aden-Owen-Carlsberg triple junction, and opening of the Gulf of Aden. *Journal of Geophysical Research: Solid Earth* 115. <https://doi.org/10.1029/2008JB006257>

Frakes, L.A., 1982. Metal chemistry of manganese nodules from the Cape Leeuwin field, southeast Indian Ocean. *Marine Geology* 47, M1–M10. [https://doi.org/10.1016/0025-3227\(82\)90014-7](https://doi.org/10.1016/0025-3227(82)90014-7)

Frakes, L.A., Exon, N.F., Granath, J.W., 1977. Preliminary studies of the Cape Leeuwin manganese nodule deposit off Western Australia. *BMR Journal of Australian Geology & Geophysics* 2, 66–69.

Frey, F.A., Coffin, M.F., Wallace, P.J., Weis, D., Zhao, X., Wise, S.W., Wähnert, V., Teagle, D.A.H., Saccocia, P.J., Reusch, D.N., Pringle, M.S., Nicolaysen, K.E., Neal, C.R., Müller, R.D., Moore, C.L., Mahoney, J.J., Keszthelyi, L., Inokuchi, H., Duncan, R.A., Delius, H., Damuth, J.E., Damasceno, D., Coxall, H.K., Borre, M.K., Boehm, F., Barling, J., Arndt, N.T., Antretter, M., 2000a. Origin and evolution of a submarine large igneous province: The Kerguelen Plateau and Broken Ridge, southern Indian Ocean. *Earth and Planetary Science Letters* 176, 73–89. [https://doi.org/10.1016/S0012-821X\(99\)00315-5](https://doi.org/10.1016/S0012-821X(99)00315-5)

Frey, F.A., Coffin, M.F., Wallace, P.J., Weis, D., Zhao, X., Wise, S.W., Wähnert, V., Teagle, D.A.H., Saccocia, P.J., Reusch, D.N., Pringle, M.S., Nicolaysen, K.E., Neal, C.R., Müller, R.D., Moore, C.L., Mahoney, J.J., Keszthelyi, L., Inokuchi, H., Duncan, R.A., Delius, H., Damuth, J.E., Damasceno, D., Coxall, H.K., Borre, M.K., Boehm, F., Barling, J., Arndt, N.T., Antretter, M., 2000b. Origin and evolution of a submarine large igneous province: the Kerguelen Plateau and Broken Ridge,

southern Indian Ocean. *Earth and Planetary Science Letters* 176, 73–89.
[https://doi.org/10.1016/S0012-821X\(99\)00315-5](https://doi.org/10.1016/S0012-821X(99)00315-5)

Fulton, E.A., Bulman, C.M., Pethybridge, H., Goldsworthy, S.D., 2018. Modelling the Great Australian Bight Ecosystem. *Deep Sea Research Part II: Topical Studies in Oceanography*, Great Australian Bight Research Program - a whole of system investigation 157–158, 211–235.
<https://doi.org/10.1016/j.dsr2.2018.11.002>

Gasco, N., 2011. Contributions to marine science by fishery observers in the French EEZ of Kerguelen. *Cybiu* 35(SP), 93–98. <https://doi.org/10.26028/CYBIUM/2011-35SP-009>

Gates, A.R., 2016. Deep-sea life of Tanzania. National Oceanography Centre, Southampton.

Gates, A.R., Benfield, M.C., Booth, D.J., Fowler, A.M., Skropeta, D., Jones, D.O.B., 2017a. Deep-sea observations at hydrocarbon drilling locations: Contributions from the SERPENT Project after 120 field visits. *Deep Sea Research Part II: Topical Studies in Oceanography*, Advances in deep-sea biology: biodiversity, ecosystem functioning and conservation 137, 463–479.
<https://doi.org/10.1016/j.dsr2.2016.07.011>

Gates, A.R., Durden, J.M., Richmond, M.D., Muhando, C.A., Khamis, Z.A., Jones, D.O.B., 2021. Ecological considerations for marine spatial management in deep-water Tanzania. *Ocean & Coastal Management* 210, 105703. <https://doi.org/10.1016/j.ocecoaman.2021.105703>

Gates, A.R., Morris, K.J., Jones, D.O.B., Sulak, K.J., 2017b. An association between a cusk eel (*Bassozetus* sp.) and a black coral (*Schizopathes* sp.) in the deep western Indian Ocean. *Mar Biodiv* 47, 971–977. <https://doi.org/10.1007/s12526-016-0516-z>

Geodekyan, A.A., Rozanov, A.G., 1984. Manganese nodules off Mozambique [WWW Document]. URL https://hero.epa.gov/hero/index.cfm/reference/details/reference_id/8349796 (accessed 4.16.24).

Georgeson, L., Rigby, C.L., Emery, T.J., Fuller, M., Hartog, J., Williams, A.J., Hobday, A.J., Duffy, C.A.J., Simpfendorfer, C.A., Okuda, T., Stobutzki, I.C., Nicol, S.J., 2020. Ecological risks of demersal fishing on deepwater chondrichthyan populations in the Southern Indian and South Pacific Oceans. *ICES Journal of Marine Science* 77, 1711–1727.
<https://doi.org/10.1093/icesjms/fsaa019>

Geoscience Australia, 2017. Diamantina Fracture Zone and Naturaliste Plateau Multibeam Bathymetry (Tiles SI48, SJ48, SK48, SL48, SI47, SJ47, SK47, SL47, SJ46, SK46, SL46, SK45 and SL45).

Gerber, H.W., Clauss, G.F., 2010. MABEL: Recovery Operation of the First Long-Term Heavy Benthic Laboratory in the Deep Sea of Antarctica. Presented at the ASME 2009 28th International Conference on Ocean, Offshore and Arctic Engineering, American Society of Mechanical Engineers Digital Collection, pp. 195–203. <https://doi.org/10.1115/OMAE2009-80251>

Gerdes, K., Martínez Arbizu, P., Schwarz-Schampera, U., Schwentner, M., Kihara, T.C., 2019. Detailed Mapping of Hydrothermal Vent Fauna: A 3D Reconstruction Approach Based on Video Imagery. *Frontiers in Marine Science* 6.

Gerdes, K.H., Martínez Arbizu, P., Schwentner, M., Freitag, R., Schwarz-Schampera, U., Brandt, A., Kihara, T.C., 2019. Megabenthic assemblages at the southern Central Indian Ridge – Spatial segregation of inactive hydrothermal vents from active-, periphery- and non-vent sites. *Marine Environmental Research* 151, 104776. <https://doi.org/10.1016/j.marenvres.2019.104776>

German, C.R., Baker, E.T., Mevel, C., Tamaki, K., the FUJI Science Team, 1998. Hydrothermal activity along the southwest Indian ridge. *Nature* 395, 490–493.
<https://doi.org/10.1038/26730>

German, C.R., Ramirez-Llodra, E., Baker, M.C., Tyler, P.A., Committee, and the C.S.S., 2011. Deep-Water Chemosynthetic Ecosystem Research during the Census of Marine Life Decade and Beyond: A Proposed Deep-Ocean Road Map. *PLOS ONE* 6, e23259.
<https://doi.org/10.1371/journal.pone.0023259>

Gibbons, Ana D., Barckhausen, U., van den Bogaard, P., Hoernle, K., Werner, R., Whittaker, J.M., Müller, R.D., 2012. Constraining the Jurassic extent of Greater India: Tectonic evolution of the West Australian margin. *Geochemistry, Geophysics, Geosystems* 13.
<https://doi.org/10.1029/2011GC003919>

Gibbons, A.D., Whittaker, J.M., Müller, R.D., 2013. The breakup of East Gondwana: Assimilating constraints from Cretaceous ocean basins around India into a best-fit tectonic model. *Journal of Geophysical Research: Solid Earth* 118, 808–822. <https://doi.org/10.1002/jgrb.50079>

Gibbons, Ana D, Van Den Bogaard, Paul, Hoernle, Kaj, Werner, Reinhard, Whittaker, Joanne M, Müller, R Dietmar, Gibbons, A D, Barckhausen, U., Van Den Bogaard, P, Hoernle, K, Werner, R, Whittaker, J M, Müller, R D, 2012. Constraining the Jurassic extent of Greater India: Tectonic evolution of the West Australian margin. *Geochemistry, Geophysics, Geosystems* 13, 5–13.
<https://doi.org/10.1029/2011GC003919>

Glasby, G.P., 1973. Distribution of manganese nodules and lebensspuren in underwater photographs from the Carlsberg Ridge, Indian Ocean. *New Zealand Journal of Geology and Geophysics* 16, 1–17. <https://doi.org/10.1080/00288306.1973.10425383>

Glover, A.G., Gooday, A.J., Bailey, D.M., Billett, D.S.M., Chevaldonné, P., Colaço, A., Copley, J., Cuvelier, D., Desbruyères, D., Kalogeropoulou, V., Klages, M., Lampadariou, N., Lejeusne, C., Mestre, N.C., Paterson, G.L.J., Perez, T., Ruhl, H., Sarrazin, J., Soltwedel, T., Soto, E.H., Thatje, S., Tselepidis, A., Van Gaeve, S., Vanreusel, A., 2010. Temporal Change in Deep-Sea Benthic Ecosystems: A Review of the Evidence From Recent Time-Series Studies, in: Lesser, M. (Ed.), *Advances in Marine Biology*. Academic Press, pp. 1–95. <https://doi.org/10.1016/B978-0-12-381015-1.00001-0>

Gonsalves, M.-J., Tiwari, S., Padate, V.P., Trivedi, J.N., Samuel, V.D., Viswanathan, C., 2022. First Record of Galatheid Squat Lobster, *Shinkaia crosnieri* Baba & Williams, 1998 (Decapoda: Galatheaidea) From Cold-seep Environment of the Indian Ocean. *Thalassas* 38, 1245–1252.
<https://doi.org/10.1007/s41208-022-00467-9>

Gooday, A.J., Bett, B.J., Escobar, E., Ingole, B., Levin, L.A., Neira, C., Raman, A.V., Sellanes, J., 2010. Habitat heterogeneity and its influence on benthic biodiversity in oxygen minimum zones. *Marine Ecology* 31, 125–147. <https://doi.org/10.1111/j.1439-0485.2009.00348.x>

Gooday, A.J., Levin, L.A., Aranda da Silva, A., Bett, B.J., Cowie, G.L., Dissard, D., Gage, J.D., Hughes, D.J., Jeffreys, R., Lamont, P.A., Larkin, K.E., Murty, S.J., Schumacher, S., Whitcraft, C., Woulds, C., 2009. Faunal responses to oxygen gradients on the Pakistan margin: A comparison of foraminiferans, macrofauna and megafauna. *Deep Sea Research Part II: Topical Studies in Oceanography, Benthic Biological and Biogeochemical Patterns and Processes Across an Oxygen Minimum Zone (Pakistan Margin, NE Arabian Sea)* 56, 488–502.
<https://doi.org/10.1016/j.dsr2.2008.10.003>

GOSLIN, J., SEGOUFIN, J., SCHLICH, R., FISHER, R.L., 1980. Submarine topography and shallow structure of the Madagascar Ridge, western Indian Ocean. *GSA Bulletin* 91, 741–753. [https://doi.org/10.1130/0016-7606\(1980\)91<741:STASSO>2.0.CO;2](https://doi.org/10.1130/0016-7606(1980)91<741:STASSO>2.0.CO;2)

Grorud-Colvert, K., Sullivan-Stack, J., Roberts, C., Constant, V., Horta e Costa, B., Pike, E.P., Kingston, N., Laffoley, D., Sala, E., Claudet, J., Friedlander, A.M., Gill, D.A., Lester, S.E., Day, J.C., Gonçalves, E.J., Ahmadi, G.N., Rand, M., Villagomez, A., Ban, N.C., Gurney, G.G., Spalding, A.K., Bennett, N.J., Briggs, J., Morgan, L.E., Moffitt, R., Deguignet, M., Pikitch, E.K., Darling, E.S., Jessen, S., Hameed, S.O., Di Carlo, G., Guidetti, P., Harris, J.M., Torre, J., Kizilkaya, Z., Agardy, T., Cury, P., Shah, N.J., Sack, K., Cao, L., Fernandez, M., Lubchenco, J., 2021. The MPA Guide: A framework to achieve global goals for the ocean. *Science* 373, eabf0861. <https://doi.org/10.1126/science.abf0861>

Haddock, S.H.D., Choy, C.A., 2024. Life in the Midwater: The Ecology of Deep Pelagic Animals. *Annual Review of Marine Science* 16, 383–416. <https://doi.org/10.1146/annurev-marine-031623-095435>

Halbach, P.E., Jahn, A., Cherkashov, G., 2017. Marine Co-Rich Ferromanganese Crust Deposits: Description and Formation, Occurrences and Distribution, Estimated World-wide Resources, in: Sharma, R. (Ed.), *Deep-Sea Mining: Resource Potential, Technical and Environmental Considerations*. Springer International Publishing, Cham, pp. 65–141. https://doi.org/10.1007/978-3-319-52557-0_3

Halpin, J.A., Crawford, A.J., Direen, N.G., Coffin, M.F., Forbes, C.J., Borissova, I., 2008. Naturaliste Plateau, offshore Western Australia: A submarine window into Gondwana assembly and breakup. *Geology* 36, 807–810. <https://doi.org/10.1130/G25059A.1>

Han, Y., Gonnella, G., Adam, N., Schippers, A., Burkhardt, L., Kurtz, S., Schwarz-Schampera, U., Franke, H., Perner, M., 2018. Hydrothermal chimneys host habitat-specific microbial communities: analogues for studying the possible impact of mining seafloor massive sulfide deposits. *Sci Rep* 8, 10386. <https://doi.org/10.1038/s41598-018-28613-5>

Hand, K.P., Bartlett, D.H., Fryer, P., Peoples, L., Williford, K., Hofmann, A.E., Cameron, J., 2020. Discovery of novel structures at 10.7 km depth in the Mariana Trench may reveal chemolithoautotrophic microbial communities. *Deep Sea Research Part I: Oceanographic Research Papers* 160, 103238. <https://doi.org/10.1016/j.dsr.2020.103238>

Harris, P.T., Baker, E.K., 2020. Chapter 60 - GeoHab Atlas of seafloor geomorphic features and benthic habitats—synthesis and lessons learned, in: Harris, P.T., Baker, E. (Eds.), *Seafloor Geomorphology as Benthic Habitat (Second Edition)*. Elsevier, pp. 969–990. <https://doi.org/10.1016/B978-0-12-814960-7.00060-9>

Harris, P.T., Macmillan-Lawler, M., Rupp, J., Baker, E.K., 2014. Geomorphology of the oceans. *Marine Geology, 50th Anniversary Special Issue* 352, 4–24. <https://doi.org/10.1016/j.margeo.2014.01.011>

Hartman, S.E., Bett, B.J., Durden, J.M., Henson, S.A., Iversen, M., Jeffreys, R.M., Horton, T., Lampitt, R., Gates, A.R., 2021. Enduring science: Three decades of observing the Northeast Atlantic from the Porcupine Abyssal Plain Sustained Observatory (PAP-SO). *Progress in Oceanography* 191, 102508. <https://doi.org/10.1016/j.pocean.2020.102508>

Hartnady, C.J.H., 1986. Amirante Basin, western Indian Ocean: Possible impact site of the Cretaceous/Tertiary extinction bolide? *Geology* 14, 423–426. [https://doi.org/10.1130/0091-7613\(1986\)14<423:ABWIOP>2.0.CO;2](https://doi.org/10.1130/0091-7613(1986)14<423:ABWIOP>2.0.CO;2)

Hashimoto, J., Ohta, S., Gamo, T., Chiba, H., Yamaguchi, T., Tsuchida, S., Okudaira, T., Watabe, H., Yamanaka, T., Kitazawa, M., 2001. First Hydrothermal Vent Communities from the Indian Ocean Discovered. *J. Zool.* 18, 717–721. <https://doi.org/10.2108/zsj.18.717>

Heap, A.D., Edwards, J., Fountain, L., Spinnocchia, M., Hughes, M., Mathews, E., Griffin, J., Borissova, I., Blevin, J., Mitchell, C., Krassay, A., 2008. Geomorphology, Sedimentology, and Stratigraphy of Submarine Canyons on the SW Australian Slope.

Heezen, B.C., Nafe, J.E., 1964. Vema trench: western Indian Ocean. *Deep Sea Research and Oceanographic Abstracts* 11, 79–84. [https://doi.org/10.1016/0011-7471\(64\)91083-6](https://doi.org/10.1016/0011-7471(64)91083-6)

Hein, J.R., Mizell, K., Koschinsky, A., Conrad, T.A., 2013. Deep-ocean mineral deposits as a source of critical metals for high- and green-technology applications: Comparison with land-based resources. *Ore Geology Reviews* 51, 1–14. <https://doi.org/10.1016/j.oregeorev.2012.12.001>

Hendon, H.H., Salby, M.L., 1994. The Life Cycle of the Madden–Julian Oscillation. *Journal of the Atmospheric Sciences* 51, 2225–2237. [https://doi.org/10.1175/1520-0469\(1994\)051<2225:TLCOTM>2.0.CO;2](https://doi.org/10.1175/1520-0469(1994)051<2225:TLCOTM>2.0.CO;2)

Hensen, C., Duarte, J.C., Vannucchi, P., Mazzini, A., Lever, M.A., Terrinha, P., Géli, L., Henry, P., Villinger, H., Morgan, J., Schmidt, M., Gutscher, M.-A., Bartolome, R., Tomonaga, Y., Polonia, A., Gràcia, E., Tinivella, U., Lupi, M., Çağatay, M.N., Elvert, M., Sakellariou, D., Matias, L., Kipfer, R., Karageorgis, A.P., Ruffine, L., Liebetrau, V., Pierre, C., Schmidt, C., Batista, L., Gasperini, L., Burwicz, E., Neres, M., Nuzzo, M., 2019. Marine Transform Faults and Fracture Zones: A Joint Perspective Integrating Seismicity, Fluid Flow and Life. *Frontiers in Earth Science* 7.

Henthorn, R.G., Hobson, B.W., McGill, P.R., Sherman, A.D., Smith, K.L., 2010. MARS Benthic Rover: In-situ rapid proto-testing on the Monterey Accelerated Research System, in: *OCEANS 2010 MTS/IEEE SEATTLE*. Presented at the OCEANS 2010 MTS/IEEE SEATTLE, pp. 1–7. <https://doi.org/10.1109/OCEANS.2010.5664381>

Hermes, J.C., Masumoto, Y., Beal, L.M., Roxy, M.K., Vialard, J., Andres, M., Annamalai, H., Behera, S., D’Adamo, N., Doi, T., Feng, M., Han, W., Hardman-Mountford, N., Hendon, H., Hood, R., Kido, S., Lee, C., Lee, T., Lengaigne, M., Li, J., Lumpkin, R., Navaneeth, K.N., Milligan, B., McPhaden, M.J., Ravichandran, M., Shinoda, T., Singh, A., Sloyan, B., Strutton, P.G., Subramanian, A.C., Thurston, S., Tozuka, T., Ummenhofer, C.C., Unnikrishnan, A.S., Venkatesan, R., Wang, D., Wiggert, J., Yu, L., Yu, W., 2019. A Sustained Ocean Observing System in the Indian Ocean for Climate Related Scientific Knowledge and Societal Needs. *Frontiers in Marine Science* 6.

Hernandez, I., Davies, J.S., Huvenne, V.A.I., Dissanayake, A., 2022. Marine litter in submarine canyons: A systematic review and critical synthesis. *Front. Mar. Sci.* 9. <https://doi.org/10.3389/fmars.2022.965612>

Hestetun, J.T., Rapp, H.T., Xavier, J., 2017. Carnivorous sponges (Porifera, Cladorhizidae) from the Southwest Indian Ocean Ridge seamounts. *Deep Sea Research Part II: Topical Studies in Oceanography, Advances in deep-sea biology: biodiversity, ecosystem functioning and conservation* 137, 166–189. <https://doi.org/10.1016/j.dsr2.2016.03.004>

Higgs, N.D., Gates, A.R., Jones, D.O.B., 2014. Fish Food in the Deep Sea: Revisiting the Role of Large Food-Falls. *PLOS ONE* 9, e96016. <https://doi.org/10.1371/journal.pone.0096016>

Hilario, A., Cunha, M.R., Génio, L., Marçal, A.R., Ravara, A., Rodrigues, C.F., Wiklund, H., 2015. First clues on the ecology of whale falls in the deep Atlantic Ocean: results from an experiment using cow carcasses. *Marine Ecology* 36, 82–90. <https://doi.org/10.1111/maec.12246>

- Hodel, F., Grespan, R., de Rafélis, M., Dera, G., Lezin, C., Nardin, E., Rouby, D., Aretz, M., Steinman, M., Buatier, M., Lacan, F., Jeandel, C., Chavagnac, V., 2021. Drake Passage gateway opening and Antarctic Circumpolar Current onset 31 Ma ago: The message of foraminifera and reconsideration of the Neodymium isotope record. *Chemical Geology* 570, 120171. <https://doi.org/10.1016/j.chemgeo.2021.120171>
- Hoffman, L., Kniesz, K., Arbizu, P.M., Kihara, T.C., 2022. Abyssal vent field habitats along plate margins in the Central Indian Ocean yield new species in the genus *Anatoma* (Vetigastropoda: Anatomidae). *European Journal of Taxonomy* 826, 135–162. <https://doi.org/10.5852/ejt.2022.826.1841>
- Hood, R., et al., 2015. Science Plan of the Second International Indian Ocean Expedition (IIOE-2): a Basin-Wide Research Program. (2015-2020)., IIOE-2-DOC_AD_2. Scientific Committee on Oceanic Research, Newark, Delaware.
- Hood, R.R., Beckley, L.E., 2019. The Second International Indian Ocean Expedition (IIOE-2): Motivating New Exploration in a Poorly Understood Ocean Basin (Volume 1). *Deep Sea Research Part II: Topical Studies in Oceanography, The 2nd International Indian Ocean Expedition (IIOE-2): Motivating New Exploration in a Poorly Understood Ocean Basin (Volume 1)* 161, 2–4. <https://doi.org/10.1016/j.dsr2.2019.04.009>
- Hood, R.R., Beckley, L.E., Vialard, J., Gaye, B., 2022. The Second International Indian Ocean Expedition (IIOE-2): New exploration in a poorly understood ocean basin (volume 4). *Deep Sea Research Part II: Topical Studies in Oceanography* 197, 105047. <https://doi.org/10.1016/j.dsr2.2022.105047>
- Hood, R.R., Beckley, L.E., Wiggert, J.D., 2017. Biogeochemical and ecological impacts of boundary currents in the Indian Ocean. *Progress in Oceanography* 156, 290–325. <https://doi.org/10.1016/j.pocean.2017.04.011>
- Hood, R.R., Urban, E.R., McPhaden, M.J., Su, D., Raes, E., 2016. The 2nd International Indian Ocean Expedition (IIOE-2): Motivating New Exploration in a Poorly Understood Basin. *Limnology and Oceanography Bulletin* 25, 117–124. <https://doi.org/10.1002/lob.10149>
- Hook, S.E., Revill, A.T., Mondon, J., Corbett, P., Armstrong, E.K., Song, J., Tanner, J.E., Stalvies, C., Ross, A.S., Williams, A., 2018. Naturally occurring hydrocarbon content and baseline condition of deep-sea benthic fauna from the Great Australian Bight. *Deep Sea Research Part II: Topical Studies in Oceanography, Great Australian Bight Research Program - a whole of system investigation* 157–158, 106–120. <https://doi.org/10.1016/j.dsr2.2018.08.005>
- Horii, T., Ueki, I., Ando, K., Mizuno, K., 2013. Eastern Indian Ocean warming associated with the negative Indian Ocean dipole: A case study of the 2010 event. *Journal of Geophysical Research: Oceans* 118, 536–549. <https://doi.org/10.1002/jgrc.20071>
- Hughes, D.J., Lamont, P.A., Levin, L.A., Packer, M., Feeley, K., Gage, J.D., 2009. Macrofaunal communities and sediment structure across the Pakistan margin Oxygen Minimum Zone, North-East Arabian Sea. *Deep Sea Research Part II: Topical Studies in Oceanography, Benthic Biological and Biogeochemical Patterns and Processes Across an Oxygen Minimum Zone (Pakistan Margin, NE Arabian Sea)* 56, 434–448. <https://doi.org/10.1016/j.dsr2.2008.05.030>
- Hughes, J.A., Smith, T., Chaillan, F., Bett, B.J., Billett, D.S.M., Boorman, B., Fisher, E.H., Frenz, M., Wolff, G.A., 2007. Two abyssal sites in the Southern Ocean influenced by different organic matter inputs: Environmental characterization and preliminary observations on the benthic foraminifera. *Deep Sea Research Part II: Topical Studies in Oceanography, The Crozet Natural*

Iron Bloom and Export Experiment 54, 2275–2290.

<https://doi.org/10.1016/j.dsr2.2007.06.006>

Huussen, T.N., Naveira-Garabato, A.C., Bryden, H.L., McDonagh, E.L., 2012. Is the deep Indian Ocean MOC sustained by breaking internal waves? *Journal of Geophysical Research: Oceans* 117. <https://doi.org/10.1029/2012JC008236>

Huvenne, V.A.I., Davies, J.S., 2014. Towards a new and integrated approach to submarine canyon research. *Deep Sea Research Part II: Topical Studies in Oceanography, Submarine Canyons: Complex Deep-Sea Environments Unravelling by Multidisciplinary Research* 104, 1–5. <https://doi.org/10.1016/j.dsr2.2013.09.012>

Ihara, C., Kushnir, Y., Cane, M.A., 2008. Warming Trend of the Indian Ocean SST and Indian Ocean Dipole from 1880 to 2004. *Journal of Climate* 21, 2035–2046. <https://doi.org/10.1175/2007JCLI1945.1>

Iken, K., Brey, T., Wand, U., Voigt, J., Junghans, P., 2001. Food web structure of the benthic community at the Porcupine Abyssal Plain (NE Atlantic): a stable isotope analysis. *Progress in Oceanography, High resolution temporal and spatial study of the benthic biology and geochemistry of a North-Eastern Atlantic abyssal locality (BENGAL)* 50, 383–405. [https://doi.org/10.1016/S0079-6611\(01\)00062-3](https://doi.org/10.1016/S0079-6611(01)00062-3)

Ingole, B.S., Ansari, Z.A., Rathod, V., Rodrigues, N., 2001. Response of deep-sea macrobenthos to a small-scale environmental disturbance. *Deep Sea Research Part II: Topical Studies in Oceanography, Indian Deep-sea Environment Experiment (INDEX)* 48, 3401–3410. [https://doi.org/10.1016/S0967-0645\(01\)00048-0](https://doi.org/10.1016/S0967-0645(01)00048-0)

Ingole, B.S., Ansari, Z.A., Rathod, V., Rodrigues, N., 2000. Response of Meiofauna to Immediate Benthic Disturbance in the Central Indian Ocean Basin. *Marine Georesources & Geotechnology* 18, 263–272. <https://doi.org/10.1080/10641190009353794>

Ingole, B.S., Koslow, J.A., 2005. Deep-sea ecosystems of the Indian Ocean. *Indian Journal of Marine Sciences* 34, 27–34.

Ingole, B.S., Pavithran, S., Ansari, Z.A., 2005. Restoration of Deep-Sea Macrofauna after Simulated Benthic Disturbance in the Central Indian Basin. *Marine Georesources & Geotechnology* 23, 267–288. <https://doi.org/10.1080/10641190500446573>

Ingole, B.S., Sautya, S., Sivadas, S., Singh, R., Nanajkar, M., 2010. Macrofaunal community structure in the western Indian continental margin including the oxygen minimum zone. *Marine Ecology* 31, 148–166. <https://doi.org/10.1111/j.1439-0485.2009.00356.x>

International Hydrographic Organization (IHO), 1953. Limits of Oceans and Seas [WWW Document]. International Hydrographic Organization. URL <https://epic.awi.de/id/eprint/29772/> (accessed 8.15.23).

International Seabed Authority, 2019. Current Status of the Reserved Areas with the International Seabed Authority.

ISA, 2022a. Exploration areas - International Seabed Authority. URL <https://www.isa.org.jm/exploration-contracts/exploration-areas/> (accessed 11.3.23).

ISA, 2022b. Reserved Areas - International Seabed Authority. URL <https://www.isa.org.jm/exploration-contracts/reserved-areas/> (accessed 11.3.23).

- Iyer, S.D., Das, P., Kalangutkar, N.G., Mehta, C.M., 2012. Seamounts – windows of opportunities and the Indian scenario. *Current Science* 102, 1382–1391.
- Jacob, J., Dymant, J., Yatheesh, V., 2014. Revisiting the structure, age, and evolution of the Wharton Basin to better understand subduction under Indonesia. *Journal of Geophysical Research: Solid Earth* 119, 169–190. <https://doi.org/10.1002/2013JB010285>
- Jacques, G., Hauff, F., Hoernle, K., Werner, R., Uenzelmann-Neben, G., Garbe-Schönberg, D., Fischer, M., 2019. Nature and origin of the Mozambique Ridge, SW Indian Ocean. *Chemical Geology* 507, 9–22. <https://doi.org/10.1016/j.chemgeo.2018.12.027>
- Jamieson, A., 2015. *The Hadal Zone: Life in the Deepest Oceans*. Cambridge University Press, Cambridge. <https://doi.org/10.1017/CBO9781139061384>
- Jamieson, A.J., 2016. Landers, in: *Biological Sampling in the Deep Sea*. John Wiley & Sons, Ltd, pp. 228–259. <https://doi.org/10.1002/9781118332535.ch11>
- Jamieson, Alan J., Boorman, B., Jones, D.O.B., 2013. Deep-Sea Benthic Sampling, in: *Methods for the Study of Marine Benthos*. John Wiley & Sons, Ltd, pp. 285–347. <https://doi.org/10.1002/9781118542392.ch7>
- Jamieson, A.J., Fujii, T., Mayor, D.J., Solan, M., Priede, I.G., 2010. Hadal trenches: the ecology of the deepest places on Earth. *Trends in Ecology & Evolution* 25, 190–197. <https://doi.org/10.1016/j.tree.2009.09.009>
- Jamieson, A.J., Giles, G.T., Stewart, H.A., in review. Hadal zones of the Southwest Pacific and East Indian oceans.
- Jamieson, A. J., Lacey, N.C., Lörz, A.-N., Rowden, A.A., Piortney, S.B., 2013. The supergiant amphipod *Alicella gigantea* (Crustacea: Alicellidae) from hadal depths in the Kermadec Trench, SW Pacific Ocean. *Deep Sea Research Part II: Topical Studies in Oceanography, Deep-Sea Biodiversity and Life History Processes* 92, 107–113. <https://doi.org/10.1016/j.dsr2.2012.12.002>
- Jamieson, A.J., Lindsay, D.J., Kitazato, H., 2023. Maximum depth extensions for Hydrozoa, Tunicata and Ctenophora. *Mar Biol* 170, 33. <https://doi.org/10.1007/s00227-023-04177-5>
- Jamieson, A.J., Linley, T.D., 2021. Hydrozoans, scyphozoans, larvaceans and ctenophores observed in situ at hadal depths. *Journal of Plankton Research* 43, 20–32. <https://doi.org/10.1093/plankt/fbaa062>
- Jamieson, A.J., Linley, T.D., Eigler, S., Macdonald, T., 2021. A global assessment of fishes at lower abyssal and upper hadal depths (5000 to 8000 m). *Deep Sea Research Part I: Oceanographic Research Papers* 178, 103642. <https://doi.org/10.1016/j.dsr.2021.103642>
- Jamieson, A.J., Onda, D.F.L., 2022. Lebensspuren and müllspuren: Drifting plastic bags alter microtopography of seafloor at full ocean depth (10,000 m, Philippine Trench). *Continental Shelf Research* 250, 104867. <https://doi.org/10.1016/j.csr.2022.104867>
- Jamieson, A.J., Stewart, H.A., 2021. Hadal zones of the Northwest Pacific Ocean. *Progress in Oceanography* 190, 102477. <https://doi.org/10.1016/j.pocean.2020.102477>
- Jamieson, A.J., Stewart, H.A., Weston, J.N.J., Lahey, P., Vescovo, V.L., 2022. Hadal Biodiversity, Habitats and Potential Chemosynthesis in the Java Trench, Eastern Indian Ocean. *Frontiers in Marine Science* 9.

- Jamieson, A.J., Vecchione, M., 2020. First in situ observation of Cephalopoda at hadal depths (Octopoda: Opisthoteuthidae: Grimpoteuthis sp.). *Mar Biol* 167, 82. <https://doi.org/10.1007/s00227-020-03701-1>
- Jamieson, A.J., Weston, J.N.J., 2023. Amphipoda from depths exceeding 6,000 meters revisited 60 years on. *Journal of Crustacean Biology* 43, ruad020. <https://doi.org/10.1093/jcbiol/ruad020>
- Jamieson, J.W., Gartman, A., 2020. Defining active, inactive, and extinct seafloor massive sulfide deposits. *Marine Policy* 117, 103926. <https://doi.org/10.1016/j.marpol.2020.103926>
- Jang, S.-J., Cho, S.-Y., Li, C., Zhou, Y., Wang, H., Sun, J., Patra, A.K., Won, Y.-J., 2023. Geographical subdivision of Alviniconcha snail populations in the Indian Ocean hydrothermal vent regions. *Frontiers in Marine Science* 10.
- Janßen, F., Treude, T., Witte, U., 2000. Scavenger assemblages under differing trophic conditions: a case study in the deep Arabian Sea. *Deep Sea Research Part II: Topical Studies in Oceanography* 47, 2999–3026. [https://doi.org/10.1016/S0967-0645\(00\)00056-4](https://doi.org/10.1016/S0967-0645(00)00056-4)
- Jempson, I., 2019. Hmas “Diamantina” – Oceanographic Research. *The Great Circle* 41, 15–37.
- Jenner, K.C.S., Jenner, M.-N.M., McCabe, K.A., 2001. GEOGRAPHICAL AND TEMPORAL MOVEMENTS OF HUMPBACK WHALES IN WESTERN AUSTRALIAN WATERS. *The APPEA Journal* 41, 749–765. <https://doi.org/10.1071/aj00044>
- Johannes, R.E., Pearce, A.F., Wiebe, W.J., Crossland, C.J., Rimmer, D.W., Smith, D.F., Manning, C., 1994. Nutrient Characteristics of Well-mixed Coastal Waters off Perth, Western Australia. *Estuarine, Coastal and Shelf Science* 39, 273–285. <https://doi.org/10.1006/ecss.1994.1064>
- Jones, D.O.B., Kaiser, S., Sweetman, A.K., Smith, C.R., Menot, L., Vink, A., Trueblood, D., Greinert, J., Billett, D.S.M., Arbizu, P.M., Radziejewska, T., Singh, R., Ingole, B., Stratmann, T., Simon-Lledó, E., Durden, J.M., Clark, M.R., 2017. Biological responses to disturbance from simulated deep-sea polymetallic nodule mining. *PLOS ONE* 12, e0171750. <https://doi.org/10.1371/journal.pone.0171750>
- Kara, A.B., Rochford, P.A., Hurlburt, H.E., 2003. Mixed layer depth variability over the global ocean. *Journal of Geophysical Research: Oceans* 108. <https://doi.org/10.1029/2000JC000736>
- Kennedy, B.R.C., Rotjan, R.D., 2023. Mind the gap: comparing exploration effort with global biodiversity patterns and climate projections to determine ocean areas with greatest exploration needs. *Frontiers in Marine Science* 10.
- Kennett, J.P., Watkins, N.D., 1975. Deep-Sea Erosion and Manganese Nodule Development in the Southeast Indian Ocean. *Science* 188, 1011–1013. <https://doi.org/10.1126/science.188.4192.1011>
- Kitchingman, A., Lai, S., Morato, T., Pauly, D., 2007. How Many Seamounts are There and Where are They Located?, in: *Seamounts: Ecology, Fisheries & Conservation*. John Wiley & Sons, Ltd, pp. 26–40. <https://doi.org/10.1002/9780470691953.ch2>
- Klein, E., Appeltans, W., Provoost, P., Saeedi, H., Benson, A., Bajona, L., Peralta, A.C., Bristol, R.S., 2019. OBIS Infrastructure, Lessons Learned, and Vision for the Future. *Front. Mar. Sci.* 6. <https://doi.org/10.3389/fmars.2019.00588>
- Kniesz, K., Jażdżewska, A.M., Martínez Arbizu, P., Kihara, T.C., 2022. DNA Barcoding of Scavenging Amphipod Communities at Active and Inactive Hydrothermal Vents in the Indian Ocean. *Front. Mar. Sci.* 8. <https://doi.org/10.3389/fmars.2021.752360>

Knudsen, J., 1967. The Deep-Sea Bivalvia. The John Murray Expedition, 1933-34, Scientific Reports; Voume XI, No. 3. J. Knudsen. Scientific Reports 11, 343–344.
<https://doi.org/10.1086/405869>

Kolbusz, J., Zika, J., Pattiaratchi, C., Jamieson, A., 2024. Water properties and bottom water patterns in hadal trench environments. *Ocean Science* 20, 123–140.
<https://doi.org/10.5194/os-20-123-2024>

Kuhn, T., Wegorzewski, A., Rühlemann, C., Vink, A., 2017. Composition, Formation, and Occurrence of Polymetallic Nodules, in: Sharma, R. (Ed.), *Deep-Sea Mining: Resource Potential, Technical and Environmental Considerations*. Springer International Publishing, Cham, pp. 23–63. https://doi.org/10.1007/978-3-319-52557-0_2

Kumagai, H., Nakamura, K., Toki, T., Morishita, T., Okino, K., Ishibashi, J. -i., Tsunogai, U., Kawagucci, S., Gamo, T., Shibuya, T., Sawaguchi, T., Neo, N., Joshima, M., Sato, T., Takai, K., 2008. Geological background of the Kairei and Edmond hydrothermal fields along the Central Indian Ridge: Implications of their vent fluids' distinct chemistry. *Geofluids* 8, 239–251.
<https://doi.org/10.1111/j.1468-8123.2008.00223.x>

Lampitt, R.S., Billett, D.S.M., Martin, A.P., 2010. The sustained observatory over the Porcupine Abyssal Plain (PAP): Insights from time series observations and process studies. *Deep Sea Research Part II: Topical Studies in Oceanography, Water Column and Seabed Studies at the PAP Sustained Observatory in the Northeast Atlantic* 57, 1267–1271.
<https://doi.org/10.1016/j.dsr2.2010.01.003>

Laptikhovsky, V., Boersch-Supan, P., Bolstad, K., Kemp, K., Letessier, T., Rogers, A.D., 2017. Cephalopods of the Southwest Indian Ocean Ridge: A hotspot of biological diversity and absence of endemism. *Deep Sea Research Part II: Topical Studies in Oceanography, The Pelagic Ecology of Seamounts of the South West Indian Ocean* 136, 98–107.
<https://doi.org/10.1016/j.dsr2.2015.07.002>

Laran, S., Authier, M., Van Canneyt, O., Dorémus, G., Watremez, P., Ridoux, V., 2017. A Comprehensive Survey of Pelagic Megafauna: Their Distribution, Densities, and Taxonomic Richness in the Tropical Southwest Indian Ocean. *Front. Mar. Sci.* 4.
<https://doi.org/10.3389/fmars.2017.00139>

Lee, H., Galy, V., Feng, X., Ponton, C., Galy, A., France-Lanord, C., Feakins, S.J., 2019. Sustained wood burial in the Bengal Fan over the last 19 My. *Proceedings of the National Academy of Sciences* 116, 22518–22525. <https://doi.org/10.1073/pnas.1913714116>

Letessier, T.B., De Grave, S., Boersch-Supan, P.H., Kemp, K.M., Brierley, A.S., Rogers, A.D., 2017. Seamount influences on mid-water shrimps (Decapoda) and gnathophausiids (Lophogastridea) of the South-West Indian Ridge. *Deep Sea Research Part II: Topical Studies in Oceanography, The Pelagic Ecology of Seamounts of the South West Indian Ocean* 136, 85–97.
<https://doi.org/10.1016/j.dsr2.2015.05.009>

Levchenko, O.V., Sushchevskaya, N.M., Marinova, Yu.G., 2021. The Nature and Evolution of the Ninetyeast Ridge: A Key Tectonic and Magmatic Feature of the East Indian Ocean. *Geotecton.* 55, 194–218. <https://doi.org/10.1134/S0016852121020060>

Levin, L.A., Bett, B.J., Gates, A.R., Heimbach, P., Howe, B.M., Janssen, F., McCurdy, A., Ruhl, H.A., Snelgrove, P., Stocks, K.I., Bailey, D., Baumann-Pickering, S., Beaverson, C., Benfield, M.C., Booth, D.J., Carreiro-Silva, M., Colaço, A., Eblé, M.C., Fowler, A.M., Gjerde, K.M., Jones, D.O.B., Katsumata, K., Kelley, D., Le Bris, N., Leonardi, A.P., Lejzerowicz, F., Macreadie, P.I., McLean, D., Meitz, F.,

- Morato, T., Netburn, A., Pawlowski, J., Smith, C.R., Sun, S., Uchida, H., Vardaro, M.F., Venkatesan, R., Weller, R.A., 2019. Global Observing Needs in the Deep Ocean. *Frontiers in Marine Science* 6.
- Levin, L.A., Gage, J.D., Martin, C., Lamont, P.A., 2000. Macrobenthic community structure within and beneath the oxygen minimum zone, NW Arabian Sea. *Deep Sea Research Part II: Topical Studies in Oceanography* 47, 189–226. [https://doi.org/10.1016/S0967-0645\(99\)00103-4](https://doi.org/10.1016/S0967-0645(99)00103-4)
- Levin, L.A., McGregor, A.L., Mendoza, G.F., Woulds, C., Cross, P., Witte, U., Gooday, A.J., Cowie, G., Kitazato, H., 2013. Macrofaunal colonization across the Indian margin oxygen minimum zone. *Biogeosciences* 10, 7161–7177. <https://doi.org/10.5194/bg-10-7161-2013>
- Levin, L.A., Sibuet, M., 2012. Understanding Continental Margin Biodiversity: A New Imperative. *Annual Review of Marine Science* 4, 79–112. <https://doi.org/10.1146/annurev-marine-120709-142714>
- Levin, L.A., Whitcraft, C.R., Mendoza, G.F., Gonzalez, J.P., Cowie, G., 2009. Oxygen and organic matter thresholds for benthic faunal activity on the Pakistan margin oxygen minimum zone (700–1100m). *Deep Sea Research Part II: Topical Studies in Oceanography, Benthic Biological and Biogeochemical Patterns and Processes Across an Oxygen Minimum Zone (Pakistan Margin, NE Arabian Sea)* 56, 449–471. <https://doi.org/10.1016/j.dsr2.2008.05.032>
- Li, C., Long, H., Yang, S., Zhang, Y., Tang, F., Jin, W., Wang, G., Chang, W., Pi, Y., Gao, L., Ma, L., Zhao, M., Zheng, H., Gong, Y., Liu, Y., Jiang, K., 2022. eDNA assessment of pelagic fish diversity, distribution, and abundance in the central Pacific Ocean. *Regional Studies in Marine Science* 56, 102661. <https://doi.org/10.1016/j.rsma.2022.102661>
- Li, Q., Liu, Y., Li, G., Wang, Z., Zheng, Z., Sun, Y., Lei, N., Li, Qi, Zhang, W., 2022. Review of the Impact of Whale Fall on Biodiversity in Deep-Sea Ecosystems. *Frontiers in Ecology and Evolution* 10.
- Lochte, K., Pfannkuche, O., 2000. How the deep Arabian Sea reflects the productivity of surface waters. *U.S. JGOFS News Vol. 10*, 10–11.
- Longhurst, A.R., 2007. *Ecological Geography of the Sea*. Elsevier.
- Lord, C., Duhamel, G., Pruvost, P., 2006. The Patagonian toothfish (*Dissostichus eleginoides*) fishery in the Kerguelen Islands (Indian Ocean sector of the Southern Ocean). *CCAMLR Science* 13, 1–25.
- Lörz, A.-N., Schwentner, M., Bober, S., Jażdżewska, A.M., 2023. Multi-ocean distribution of a brooding predator in the abyssal benthos. *Sci Rep* 13, 15867. <https://doi.org/10.1038/s41598-023-42942-0>
- Lourey, M.J., Dunn, J.R., Waring, J., 2006. A mixed-layer nutrient climatology of Leeuwin Current and Western Australian shelf waters: Seasonal nutrient dynamics and biomass. *Journal of Marine Systems* 59, 25–51. <https://doi.org/10.1016/j.jmarsys.2005.10.001>
- Lutz, M.J., Caldeira, K., Dunbar, R.B., Behrenfeld, M.J., 2007. Seasonal rhythms of net primary production and particulate organic carbon flux to depth describe the efficiency of biological pump in the global ocean. *Journal of Geophysical Research: Oceans* 112. <https://doi.org/10.1029/2006JC003706>
- Ma, Y., Cochran, J.R., 1996. Transitions in axial morphology along the Southeast Indian Ridge. *Journal of Geophysical Research: Solid Earth* 101, 15849–15866. <https://doi.org/10.1029/95JB03038>

- MacIntosh, H., Althaus, F., Williams, A., Tanner, J.E., Alderslade, P., Ah Yong, S.T., Bax, N., Criscione, F., Crowther, A.L., Farrelly, C.A., Finn, J.K., Goudie, L., Gowlett-Holmes, K., Hosie, A.M., Kupriyanova, E., Mah, C., McCallum, A.W., Merrin, K.L., Miskelly, A., Mitchell, M.L., Molodtsova, T., Murray, A., O'Hara, T.D., O'Loughlin, P.M., Paxton, H., Reid, A.L., Sorokin, S.J., Staples, D., Walker-Smith, G., Whitfield, E., Wilson, R.S., 2018. Invertebrate diversity in the deep Great Australian Bight (200–5000 m). *Marine Biodiversity Records* 11, 23. <https://doi.org/10.1186/s41200-018-0158-x>
- Macreadie, P.I., McLean, D.L., Thomson, P.G., Partridge, J.C., Jones, D.O.B., Gates, A.R., Benfield, M.C., Collin, S.P., Booth, D.J., Smith, L.L., Techera, E., Skropeta, D., Horton, T., Pattiaratchi, C., Bond, T., Fowler, A.M., 2018a. Eyes in the sea: Unlocking the mysteries of the ocean using industrial, remotely operated vehicles (ROVs). *Science of the Total Environment* 634, 1077–1091. <https://doi.org/10.1016/j.scitotenv.2018.04.049>
- Macreadie, P.I., McLean, D.L., Thomson, P.G., Partridge, J.C., Jones, D.O.B., Gates, A.R., Benfield, M.C., Collin, S.P., Booth, D.J., Smith, L.L., Techera, E., Skropeta, D., Horton, T., Pattiaratchi, C., Bond, T., Fowler, A.M., 2018b. Eyes in the sea: Unlocking the mysteries of the ocean using industrial, remotely operated vehicles (ROVs). *Science of The Total Environment* 634, 1077–1091. <https://doi.org/10.1016/j.scitotenv.2018.04.049>
- Madden, R.A., Julian, P.R., 1971. Detection of a 40–50 Day Oscillation in the Zonal Wind in the Tropical Pacific. *Journal of the Atmospheric Sciences* 28, 702–708. [https://doi.org/10.1175/1520-0469\(1971\)028<0702:DOADOI>2.0.CO;2](https://doi.org/10.1175/1520-0469(1971)028<0702:DOADOI>2.0.CO;2)
- Mammerickx, J., Sandwell, D., 1986. Rifting of old oceanic lithosphere. *Journal of Geophysical Research: Solid Earth* 91, 1975–1988. <https://doi.org/10.1029/JB091iB02p01975>
- Markl, R.G., 1978. Basement morphology and rift geometry near the former junction of India, Australia, and Antarctica. *Earth and Planetary Science Letters* 39, 211–225. [https://doi.org/10.1016/0012-821X\(78\)90197-8](https://doi.org/10.1016/0012-821X(78)90197-8)
- Marsac, F., Annasawmy, P., Noyon, M., Demarcq, H., Soria, M., Rabearisoa, N., Bach, P., Cherel, Y., Grelet, J., Romanov, E., 2020a. Seamount effect on circulation and distribution of ocean taxa in the vicinity of La Pérouse, a shallow seamount in the southwestern Indian Ocean. *Deep Sea Research Part II: Topical Studies in Oceanography, Bio-Physical coupling around three shallow seamounts in the South Western Indian Ocean, with regional comparisons based on modelling, remote sensing and observational studies* 176, 104806. <https://doi.org/10.1016/j.dsr2.2020.104806>
- Marsac, F., Galletti, F., Ternon, J.-F., Romanov, E.V., Demarcq, H., Corbari, L., Bouchet, P., Roest, W.R., Jorry, S.J., Olu, K., Loncke, L., Roberts, M.J., Ménard, F., 2020b. Seamounts, plateaus and governance issues in the southwestern Indian Ocean, with emphasis on fisheries management and marine conservation, using the Walters Shoal as a case study for implementing a protection framework. *Deep Sea Research Part II: Topical Studies in Oceanography, Bio-Physical coupling around three shallow seamounts in the South Western Indian Ocean, with regional comparisons based on modelling, remote sensing and observational studies* 176, 104715. <https://doi.org/10.1016/j.dsr2.2019.104715>
- Marsh, L., Copley, J.T., Huvenne, V.A.I., Linse, K., Reid, W.D.K., Rogers, A.D., Sweeting, C.J., Tyler, P.A., 2012. Microdistribution of Faunal Assemblages at Deep-Sea Hydrothermal Vents in the Southern Ocean. *PLOS ONE* 7, e48348. <https://doi.org/10.1371/journal.pone.0048348>

- Martin, A., Blettery, J., Dettal, A., Rosset, N., Gousseau, Y., 2023. Bendima: A database for marine macro-invertebrate bycatch data designed to improve reproducibility in benthic ecology. *Cybium* 47, 325–334. <https://doi.org/10.26028/cybium/2023-020>
- Mazumdar, A., Dewangan, P., João, H.M., Peketi, A., Khosla, V.R., Kocherla, M., Badesab, F.K., Joshi, R.K., Roxanne, P., Ramamurty, P.B., Karisiddaiah, S.M., Patil, D.J., Dayal, A.M., Ramprasad, T., Hawkesworth, C.J., Avanzinelli, R., 2009. Evidence of paleo–cold seep activity from the Bay of Bengal, offshore India. *Geochemistry, Geophysics, Geosystems* 10. <https://doi.org/10.1029/2008GC002337>
- Mazumdar, A., Dewangan, P., Peketi, A., Badesaab, F., Sadique, M., Sivan, K., Mathai, J., Ghosh, A., Zatale, A., Pillutla, S.P.K., Uma, C., Mishra, C.K., Fernandes, W., Tyagi, A., Paul, T., 2021. The first record of the genus *Lamellibrachia* (Siboglinidae) tubeworm along with associated organisms in a chemosynthetic ecosystem from the Indian Ocean: A report from the Cauvery–Mannar Basin. *J Earth Syst Sci* 130, 94. <https://doi.org/10.1007/s12040-021-01587-1>
- Mazumdar, A., Dewangan, P., Peketi, A., Gullapalli, S., Kalpana, M.S., Naik, G.P., Shetty, D., Pujari, S., Pillutla, S.P.K., Gaikwad, V.V., Nazareth, D., Sangodkar, N.S., Dakara, G., Kumar, A., Mishra, C.K., Singha, P., Reddy, R., 2019. The first record of active methane (cold) seep ecosystem associated with shallow methane hydrate from the Indian EEZ. *J Earth Syst Sci* 128, 18. <https://doi.org/10.1007/s12040-018-1044-y>
- Mazumdar, A., Joshi, R.K., Peketi, A., Kocherla, M., 2011. Occurrence of faecal pellet-filled simple and composite burrows in cold seep carbonates: A glimpse of a complex benthic ecosystem. *Marine Geology* 289, 117–121. <https://doi.org/10.1016/j.margeo.2011.09.003>
- McClain, C.R., Nunnally, C., Dixon, R., Rouse, G.W., Benfield, M., 2019. Alligators in the abyss: The first experimental reptilian food fall in the deep ocean. *PLOS ONE* 14, e0225345. <https://doi.org/10.1371/journal.pone.0225345>
- McCreary, J.P., Yu, Z., Hood, R.R., Vinaychandran, P.N., Furue, R., Ishida, A., Richards, K.J., 2013. Dynamics of the Indian-Ocean oxygen minimum zones. *Progress in Oceanography* 112–113, 15–37. <https://doi.org/10.1016/j.pocean.2013.03.002>
- McEnnulty, F.R., Gowlett-Holmes, K.L., Williams, A., Althaus, F., Fromont, J., Poore, G.C.B., O'Hara, T.D., Marsh, L., Kott, P., Slack-Smith, S., Alderslade, P., Kitahara, M.V., 2011. The deepwater megabenthic invertebrates on the western continental margin of Australia (100–1100 m depths): composition, distribution and novelty. *RECORDS OF THE WESTERN AUSTRALIAN MUSEUM Supplement 80 : Deepwater megabenthos of south-western Australia*, 1–191. <https://doi.org/10.18195/issn.0313-122x.80.2011.001-191>
- McLean, D., Speed, C.W., Birt, M.J., Colquhoun, J., Case, M., Stowar, M., Bond, T., Ierodiaconou, D., Whitmarsh, S.K., Taylor, M.D., Wines, S., Booth, D.J., Fowler, A.M., Vaughan, B.I., 2022. Habitat value of subsea wells and pipelines for fishery target species in Australia. *Frontiers in Marine Science* 9, 1537. <https://doi.org/10.3389/fmars.2022.960496>
- McPhaden, M.J., Foltz, G.R., 2013. Intraseasonal variations in the surface layer heat balance of the central equatorial Indian Ocean: The importance of zonal advection and vertical mixing. *Geophysical Research Letters* 40, 2737–2741. <https://doi.org/10.1002/grl.50536>
- Ménard, F., Potier, M., Jaquemet, S., Romanov, E., Sabatié, R., Cherel, Y., 2013. Pelagic cephalopods in the western Indian Ocean: New information from diets of top predators. *Deep Sea Research Part II: Topical Studies in Oceanography, The Role of Squids in Pelagic Ecosystems* 95, 83–92. <https://doi.org/10.1016/j.dsr2.2012.08.022>

- Mendenhall, E., Tiller, R., Nyman, E., 2023. The ship has reached the shore: The final session of the 'Biodiversity Beyond National Jurisdiction' negotiations. *Marine Policy* 155, 105686. <https://doi.org/10.1016/j.marpol.2023.105686>
- Menot, L., Sibuet, M., Carney, R.S., Levin, L.A., Rowe, G.T., Billett, D.S.M., Poore, G., Kitazato, H., Vanreusel, A., Galéron, J., Lavrado, H.P., Sellanes, J., Ingole, B., Krylova, E., 2010. New Perceptions of Continental Margin Biodiversity, in: *Life in the World's Oceans*. John Wiley & Sons, Ltd, pp. 79–102. <https://doi.org/10.1002/9781444325508.ch5>
- Milkov, A.V., 2000. Worldwide distribution of submarine mud volcanoes and associated gas hydrates. *Marine Geology* 167, 29–42. [https://doi.org/10.1016/S0025-3227\(00\)00022-0](https://doi.org/10.1016/S0025-3227(00)00022-0)
- Miller, K.A., Thompson, K.F., Johnston, P., Santillo, D., 2018. An Overview of Seabed Mining Including the Current State of Development, Environmental Impacts, and Knowledge Gaps. *Frontiers in Marine Science* 4.
- Mironov, A.N., Dilman, A.B., Gebruk, A.V., Kremenetskaia, A.V., Minin, K.V., Smirnov, I.S., 2019. Echinoderms of the Kuril-Kamchatka Trench. *Progress in Oceanography* 179, 102217. <https://doi.org/10.1016/j.pocean.2019.102217>
- Monaco Explorations, n.d. INDIAN OCEAN 2022 MISSION. Monaco Explorations. URL <https://www.monacoexplorations.org/en/missions/indian-ocean-2022-mission/> (accessed 3.11.24).
- Moore, A.B.M., Gates, A.R., 2015. Deep-water observation of scalloped hammerhead *Sphyrna lewini* in the western Indian Ocean off Tanzania. *Marine Biodiversity Records* 8, e91. <https://doi.org/10.1017/S1755267215000627>
- Mukhopadhyay, R., Ghosh, A.K., 2010. Dynamics of formation of ferromanganese nodules in the Indian Ocean. *Journal of Asian Earth Sciences* 37, 394–398. <https://doi.org/10.1016/j.jseaes.2009.09.003>
- Mukhopadhyay, R., Ghosh, A.K., Iyer, S.D., 2018a. Seafloor Environment, in: *The Indian Ocean Nodule Field*. Elsevier, pp. 181–230. <https://doi.org/10.1016/B978-0-12-805474-1.00005-1>
- Mukhopadhyay, R., Ghosh, A.K., Iyer, S.D., 2018b. *The Indian Ocean Nodule Field: Geology and Resource Potential*, 2nd ed. Elsevier.
- Mukhopadhyay, R., Iyer, S.D., Ghosh, A.K., 2002. The Indian Ocean Nodule Field: petrotectonic evolution and ferromanganese deposits. *Earth-Science Reviews* 60, 67–130. [https://doi.org/10.1016/S0012-8252\(02\)00078-8](https://doi.org/10.1016/S0012-8252(02)00078-8)
- Mukhopadhyay, R., Karisiddaiah, S.M., Ghosh, A.K., 2012. Geodynamics of the Amirante Ridge and Trench Complex, Western Indian Ocean. *International Geology Review* 54, 81–92. <https://doi.org/10.1080/00206814.2010.504076>
- Mukhopadhyay, R., Naik, S., De Souza, S., Dias, O., Iyer, S.D., Ghosh, A.K., 2019. The economics of mining seabed manganese nodules: A case study of the Indian Ocean nodule field. *Marine Georesources & Geotechnology* 37, 845–851. <https://doi.org/10.1080/1064119X.2018.1504149>
- Müller, R.D., Zahirovic, S., Williams, S.E., Cannon, J., Seton, M., Bower, D.J., Tetley, M.G., Heine, C., Le Breton, E., Liu, S., Russell, S.H.J., Yang, T., Leonard, J., Gurnis, M., 2019. A Global Plate Model Including Lithospheric Deformation Along Major Rifts and Orogens Since the Triassic. *Tectonics* 38, 1884–1907. <https://doi.org/10.1029/2018TC005462>

- Munsch, M., Schlich, R., 1989. The Rodriguez Triple Junction (Indian Ocean): Structure and evolution for the past one million years. *Marine Geophysical Researches* 11, 1–14. <https://doi.org/10.1007/BF00286244>
- Murton, B.J., Rona, P.A., 2015. Carlsberg Ridge and Mid-Atlantic Ridge: Comparison of slow spreading centre analogues. *Deep Sea Research Part II: Topical Studies in Oceanography, Exploring New Frontiers in Deep-Sea Research: In Honor and Memory of Peter A. Rona* 121, 71–84. <https://doi.org/10.1016/j.dsr2.2015.04.021>
- Murty, S.J., Bett, B.J., Gooday, A.J., 2009. Megafaunal responses to strong oxygen gradients on the Pakistan margin of the Arabian Sea. *Deep Sea Research Part II: Topical Studies in Oceanography, Benthic Biological and Biogeochemical Patterns and Processes Across an Oxygen Minimum Zone (Pakistan Margin, NE Arabian Sea)* 56, 472–487. <https://doi.org/10.1016/j.dsr2.2008.05.029>
- Nagender Nath, B., Balaram, V., Sudhakar, M., Plüger, W.L., 1992. Rare earth element geochemistry of ferromanganese deposits from the Indian Ocean. *Marine Chemistry* 38, 185–208. [https://doi.org/10.1016/0304-4203\(92\)90034-8](https://doi.org/10.1016/0304-4203(92)90034-8)
- Nagura, M., McPhaden, M.J., 2010. Dynamics of zonal current variations associated with the Indian Ocean dipole. *Journal of Geophysical Research: Oceans* 115. <https://doi.org/10.1029/2010JC006423>
- Narayanaswamy, B.E., Rea, T., Serpetti, N., Lamont, P.A., 2017. What lies within; Annelid polychaetes found in micro-habitats of coral/carbonate material from SW Indian ocean seamounts. *Deep Sea Research Part II: Topical Studies in Oceanography, Advances in deep-sea biology: biodiversity, ecosystem functioning and conservation* 137, 157–165. <https://doi.org/10.1016/j.dsr2.2016.06.018>
- Nekton, 2024. MALDIVES [WWW Document]. Nekton. URL <https://nektonmission.org/missions/maldives> (accessed 3.11.24).
- Neyman, A.A., Sokolova, M.N., Vinogradova, N.G., Pasternak, F.A., 1973. Some Patterns of the Distribution of Bottom Fauna in the Indian Ocean, in: Zeitzschel, B., Gerlach, S.A. (Eds.), *The Biology of the Indian Ocean, Ecological Studies*. Springer, Berlin, Heidelberg, pp. 467–473. https://doi.org/10.1007/978-3-642-65468-8_42
- Niyazi, Y., Bond, T., Kolbusz, J.L., Maroni, P.J., Stewart, H.A., Jamieson, A.J., 2024. Deep-sea benthic structures and substrate types influence the distribution of functional groups in the Wallaby-Zenith Fracture Zone (East Indian Ocean). *Deep Sea Research Part I: Oceanographic Research Papers* 206, 104268. <https://doi.org/10.1016/j.dsr.2024.104268>
- Obura, D., 2015. Deep sea and Offshore/Pelagic Habitats, in: *Regional State of the Coast Report: Western Indian Ocean*. United Nations.
- OceanX, 2024. <https://oceanx.org/press-room/article/oceanx-launches-scientific-research-mission-in-the-seychelles-islands-in-partners> [WWW Document]. URL <https://oceanx.org/press-room/article/oceanx-launches-scientific-research-mission-in-the-seychelles-islands-in-partners> (accessed 3.11.24).
- O'Hara, T., 2022. IN2022_V08 Voyage Summary.pdf.
- Ohta, S., 1984. Star-shaped feeding traces produced by echiuran worms on the deep-sea floor of the Bay of Bengal. *Deep Sea Research Part A. Oceanographic Research Papers* 31, 1415–1432. [https://doi.org/10.1016/0198-0149\(84\)90080-3](https://doi.org/10.1016/0198-0149(84)90080-3)

- Olierook, H.K.H., Merle, R.E., Jourdan, F., Sircombe, K., Fraser, G., Timms, N.E., Nelson, G., Dadd, K.A., Kellerson, L., Borissova, I., 2015. Age and geochemistry of magmatism on the oceanic Wallaby Plateau and implications for the opening of the Indian Ocean. *Geology* 43, 971–974. <https://doi.org/10.1130/G37044.1>
- Oliver, P.G., Vestheim, H., Antunes, A., Kaartvedt, S., 2015. Systematics, functional morphology and distribution of a bivalve (*Apachecorbula muriatica* gen. et sp. nov.) from the rim of the 'Valdivia Deep' brine pool in the Red Sea. *Journal of the Marine Biological Association of the United Kingdom* 95, 523–535. <https://doi.org/10.1017/S0025315414001234>
- O'Loughlin, P.M., Mackenzie, M., VandenSpiegel, D., 2013. New sea cucumber species from the seamounts on the Southwest Indian Ocean Ridge (Echinodermata: Holothuroidea: Aspidochirotida, Elasipodida, Dendrochirotida). *Mem. Mus. Vic.* 70, 37–50. <https://doi.org/10.24199/j.mmv.2013.70.04>
- Ovechkina, M.N., Watkeys, M.K., Mostovski, M.B., Kretzinger, W., Perritt, S.M., 2021. Calcareous nannofossils identify the age and precipitation rates of manganese deposits of the Mozambique Ridge and Mozambique Basin, SW Indian Ocean. *Geo-Mar Lett* 41, 36. <https://doi.org/10.1007/s00367-021-00707-6>
- Palomares, M.L.D., Pauly, D., 2011. A Brief History of Fishing in the Kerguelen Islands, France (No. 4), Fisheries catch reconstructions: Islands, Part II.
- Parin, N.V., Timokhin, I.G., Novikov, N.P., Shcherbachev, Yu.N., 2008. On the composition of talassobathyal ichthyofauna and commercial productivity of Mozambique Seamount (the Indian Ocean). *J. Ichthyol.* 48, 361–366. <https://doi.org/10.1134/S0032945208050019>
- Park, Y.H., Gasco, N., Duhamel, G., 2008. Slope currents around the Kerguelen Islands from demersal longline fishing records. *Geophysical Research Letters* 35. <https://doi.org/10.1029/2008GL033660>
- Parsieglä, N., Gohl, K., Uenzelmann-Neben, G., 2008. The Agulhas Plateau: Structure and evolution of a Large Igneous Province. *Geophysical Journal International* 174, 336–350. <https://doi.org/10.1111/j.1365-246X.2008.03808.x>
- Parson, L.M., Evans, A.J., 2005. Seafloor topography and tectonic elements of the Western Indian Ocean. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* 363, 15–24. <https://doi.org/10.1098/rsta.2004.1472>
- Parulekar, A.H., Harkantra, S.N., Ansari, Z.A., Matondkar, S.G.P., 1982. Abyssal benthos of the central Indian Ocean. *Deep Sea Research Part A. Oceanographic Research Papers* 29, 1531–1537. [https://doi.org/10.1016/0198-0149\(82\)90041-3](https://doi.org/10.1016/0198-0149(82)90041-3)
- Parulekar, A.H., Ingole, B.S., Harkantra, S.N., Ansari, Z.A., 1992. Deep sea benthos of the western and central Indian Ocean.
- Patriat, P., Sauter, D., Munsch, M., Parson, L., 1997. A Survey of the Southwest Indian Ridge Axis Between Atlantis II Fracture Zone and the Indian Ocean Triple Junction: Regional Setting and Large Scale Segmentation. *Marine Geophysical Researches* 19, 457–480. <https://doi.org/10.1023/A:1004312623534>
- Pearce, J.R., Linley, T.D., Bond, T., Jamieson, A.J., 2023. Depth distribution of the bigeye hound shark *Iago omanensis* and other deep-sea species observed by baited-camera in the Red Sea. *Journal of the Marine Biological Association of the United Kingdom* 103, e8. <https://doi.org/10.1017/S002531542200100X>

- Peketi, A., Mazumdar, A., Sawant, B., Manaskanya, A., Zatale, A., 2022. Biogeochemistry and trophic structure of a cold seep ecosystem, offshore Krishna-Godavari basin (east coast of India). *Marine and Petroleum Geology* 138, 105542. <https://doi.org/10.1016/j.marpetgeo.2022.105542>
- Perez, M., Sun, J., Xu, Q., Qian, P.-Y., 2021. Structure and Connectivity of Hydrothermal Vent Communities Along the Mid-Ocean Ridges in the West Indian Ocean: A Review. *Frontiers in Marine Science* 8.
- Periasamy, R., Ingole, B., 2022. Three New Carnivorous sponge species (Demospongiae: Cladorhizidae) from the Seamounts of the Central Indian Ridge. *Zootaxa* 5162, 451–486. <https://doi.org/10.11646/ZOOTAXA.5162.5.1>
- Periasamy, R., Kurian, J., Ingole, B., 2023a. Two Black Corals (Anthozoa: Antipatharia) from the Seamount of Central Indian Ridge (1917 m) and Deeper Southwest Indian Ridge (4007 m). *Thalassas*. <https://doi.org/10.1007/s41208-023-00597-8>
- Periasamy, R., Kurian, P., Ingole, B., 2023b. A new deep-water coral species *Telestula ridgensis* sp. nov (Scleralcyonacea: Sarcodictyonidae) from the seamount of the Central Indian Ridge. *Zootaxa* 5254. <https://doi.org/10.11646/zootaxa.5254.2.4>
- Periasamy, R., Kurian, P.J., Ingole, B., 2023c. Two new deep-water species of squat lobsters (Crustacea: Anomura: Galatheaidea) from the Central and Southwest Indian Ridge. *Zootaxa* 5231, 3. <https://doi.org/10.11646/ZOOTAXA.5231.2.3>
- Periasamy, R., Kurian, P.J., Ingole, B., 2023d. Two new bamboo corals species (Octocorallia: Keratoisididae) from the seamounts of slow-spreading Central Indian Ridge. *Deep Sea Research Part I: Oceanographic Research Papers* 201, 104158. <https://doi.org/10.1016/j.dsr.2023.104158>
- Periasamy, R., Kurianurian, P., Ingole, B., 2022. A new report on the deep-sea sponge-associated *Spongicoloides weijiaensis* Xu, Zhou & Wang, 2017 (Decapoda: Spongicolidae) from the Southwest Indian Ocean Ridge. *Zootaxa* 5195, 5. <https://doi.org/10.11646/ZOOTAXA.5195.3.5>
- Periasamy, R., Palayil, J.K., Ingole, B., 2023e. First record of sponge-associated deep-sea polychaete (Polynoidae: Bathymoorea) on the ultraslow-spreading Southwest Indian Ridge. *Deep Sea Research Part I: Oceanographic Research Papers* 191, 103923. <https://doi.org/10.1016/j.dsr.2022.103923>
- Péron, C., Welsford, D.C., Ziegler, P., Lamb, T.D., Gasco, N., Chazeau, C., Sinègre, R., Duhamel, G., 2016. Modelling spatial distribution of Patagonian toothfish through life-stages and sex and its implications for the fishery on the Kerguelen Plateau. *Progress in Oceanography* 141, 81–95. <https://doi.org/10.1016/j.pocean.2015.12.003>
- Pettis, R.W., de Forest, A., 1979. Chemical Composition of ferromanganese Nodules from the Southern Ocean. *Mar. Freshwater Res.* 30, 535–539. <https://doi.org/10.1071/mf9790535>
- Pfannkuche, O., Lochte, K., 2000. The biogeochemistry of the deep Arabian Sea: overview. *Deep Sea Research Part II: Topical Studies in Oceanography* 47, 2615–2628. [https://doi.org/10.1016/S0967-0645\(00\)00041-2](https://doi.org/10.1016/S0967-0645(00)00041-2)
- Phillips, H.E., Tandon, A., Furue, R., Hood, R., Ummenhofer, C.C., Benthuyssen, J.A., Menezes, V., Hu, S., Webber, B., Sanchez-Franks, A., Cherian, D., Shroyer, E., Feng, M., Wijesekera, H., Chatterjee, A., Yu, L., Hermes, J., Murtugudde, R., Tozuka, T., Su, D., Singh, A., Centurioni, L., Prakash, S., Wiggert, J., 2021. Progress in understanding of Indian Ocean circulation, variability, air–sea

exchange, and impacts on biogeochemistry. *Ocean Science* 17, 1677–1751.
<https://doi.org/10.5194/os-17-1677-2021>

Picard, K., Brooke, B.P., Harris, P.T., Siwabessy, P.J.W., Coffin, M.F., Tran, M., Spinoccia, M., Weales, J., Macmillan-Lawler, M., Sullivan, J., 2018. Malaysia Airlines flight MH370 search data reveal geomorphology and seafloor processes in the remote southeast Indian Ocean. *Marine Geology* 395, 301–319. <https://doi.org/10.1016/j.margeo.2017.10.014>

Pickering, K.T., Pouderoux, H., McNeill, L.C., Backman, J., Chemale, F., Kutterolf, S., Milliken, K.L., Mukoyoshi, H., Henstock, T.J., Stevens, D.E., Parnell, C., Dugan, B., 2020. Sedimentology, stratigraphy and architecture of the Nicobar Fan (Bengal–Nicobar Fan System), Indian Ocean: Results from International Ocean Discovery Program Expedition 362. *Sedimentology* 67, 2248–2281. <https://doi.org/10.1111/sed.12701>

Pollard, R., Read, J., 2017. Circulation, stratification and seamounts in the Southwest Indian Ocean. *Deep Sea Research Part II: Topical Studies in Oceanography, The Pelagic Ecology of Seamounts of the South West Indian Ocean* 136, 36–43.
<https://doi.org/10.1016/j.dsr2.2015.02.018>

Pollard, R., Salter, I., Sanders, R., Lucas, M., Moore, C., 2009. Southern Ocean deep-water carbon export enhanced by natural iron fertilization. *Nature* 457.

Pollard, R.T., Venables, H.J., Read, J.F., Allen, J.T., 2007. Large-scale circulation around the Crozet Plateau controls an annual phytoplankton bloom in the Crozet Basin. *Deep Sea Research Part II: Topical Studies in Oceanography* 54, 1915–1929. <https://doi.org/10.1016/J.DSR2.2007.06.012>

Poore, G.C.B., Avery, L., Błażewicz-Paszkowycz, M., Browne, J., Bruce, N.L., Gerken, S., Glasby, C., Greaves, E., McCallum, A.W., Staples, D., Syme, A., Taylor, J., Walker-Smith, G., Warne, M., Watson, C., Williams, A., Wilson, R.S., Woolley, S., 2015. Invertebrate diversity of the unexplored marine western margin of Australia: taxonomy and implications for global biodiversity. *Mar Biodiv* 45, 271–286. <https://doi.org/10.1007/s12526-014-0255-y>

Post, A., Przeslawski, R., Huang, Z., Smith, D., Kirkendale, L., Wilson, N., 2021. An eco-narrative of Gascoyne Marine Park, North-west marine region.

Post, A.L., Przeslawski, R., Huang, Z., Smith, D., Kirkendale, L., Wilson, N., 2020. Gascoyne Marine Park post-survey report, RV Falkor, FK200308 (GA4859). Geoscience Australia.
<https://doi.org/10.11636/Record.2020.030>

Poupin, J., 2015. Deep water Decapoda collected during the BENTHEDI Expedition in Mayotte region (17/03 to 15/04/1977, R/V Suroît). <https://doi.org/10.17882/40956>

Prasanna Kumar, S., Divya David, T., Byju, P., Narvekar, J., Yoneyama, K., Nakatani, N., Ishida, A., Horii, T., Masumoto, Y., Mizuno, K., 2012. Bio-physical coupling and ocean dynamics in the central equatorial Indian Ocean during 2006 Indian Ocean Dipole. *Geophysical Research Letters* 39. <https://doi.org/10.1029/2012GL052609>

Praveen Kumar, B., Vialard, J., Lengaigne, M., Murty, V.S.N., Foltz, G.R., McPhaden, M.J., Pous, S., de Boyer Montégut, C., 2014. Processes of interannual mixed layer temperature variability in the thermocline ridge of the Indian Ocean. *Clim Dyn* 43, 2377–2397.
<https://doi.org/10.1007/s00382-014-2059-y>

Przeslawski, R., Dundas, K., Radke, L., Anderson, T.J., 2012. Deep-sea Lebensspuren of the Australian continental margins. *Deep Sea Research Part I: Oceanographic Research Papers* 65, 26–35. <https://doi.org/10.1016/j.dsr.2012.03.006>

- Purkis, S.J., Shernisky, H., Swart, P.K., Sharifi, A., Oehlert, A., Marchese, F., Benzoni, F., Chimienti, G., Duchâtellier, G., Klaus, J., Eberli, G.P., Peterson, L., Craig, A., Rodrigue, M., Titschack, J., Kolodziej, G., Abdulla, A., 2022. Discovery of the deep-sea NEOM Brine Pools in the Gulf of Aqaba, Red Sea. *Commun Earth Environ* 3, 1–13. <https://doi.org/10.1038/s43247-022-00482-x>
- Raghukumar, C., Raghukumar, S., Sheelu, G., Gupta, S.M., Nagender Nath, B., Rao, B.R., 2004. Buried in time: culturable fungi in a deep-sea sediment core from the Chagos Trench, Indian Ocean. *Deep Sea Research Part I: Oceanographic Research Papers* 51, 1759–1768. <https://doi.org/10.1016/j.dsr.2004.08.002>
- Ramirez-Llodra, E., Brandt, A., Danovaro, R., De Mol, B., Escobar, E., German, C.R., Levin, L.A., Martinez Arbizu, P., Menot, L., Buhl-Mortensen, P., Narayanaswamy, B.E., Smith, C.R., Tittensor, D.P., Tyler, P.A., Vanreusel, A., Vecchione, M., 2010. Deep, diverse and definitely different: unique attributes of the world's largest ecosystem. *Biogeosciences* 7, 2851–2899. <https://doi.org/10.5194/bg-7-2851-2010>
- Ramiro-Sánchez, B., Martin, A., Leroy, B., 2023. The epitome of data paucity: Deep-sea habitats of the Southern Indian Ocean. *Biological Conservation* 283, 110096. <https://doi.org/10.1016/j.biocon.2023.110096>
- Read, J., Pollard, R., 2017. An introduction to the physical oceanography of six seamounts in the southwest Indian Ocean. *Deep Sea Research Part II: Topical Studies in Oceanography, The Pelagic Ecology of Seamounts of the South West Indian Ocean* 136, 44–58. <https://doi.org/10.1016/j.dsr2.2015.06.022>
- Renn, D., Shepard, L., Vancea, A., Karan, R., Arold, S.T., Rueping, M., 2021. Novel Enzymes From the Red Sea Brine Pools: Current State and Potential. *Frontiers in Microbiology* 12.
- Riehl, T., Wöfl, A.-C., Augustin, N., Devey, C.W., Brandt, A., 2020. Discovery of widely available abyssal rock patches reveals overlooked habitat type and prompts rethinking deep-sea biodiversity. *Proceedings of the National Academy of Sciences* 117, 15450–15459. <https://doi.org/10.1073/pnas.1920706117>
- Rixen, T., Cowie, G., Gaye, B., Goes, J., do Rosário Gomes, H., Hood, R.R., Lachkar, Z., Schmidt, H., Segsneider, J., Singh, A., 2020. Reviews and syntheses: Present, past, and future of the oxygen minimum zone in the northern Indian Ocean. *Biogeosciences* 17, 6051–6080. <https://doi.org/10.5194/bg-17-6051-2020>
- Roberts, M.J., Ternon, J.-F., Marsac, F., Noyon, M., Payne, A.I.L., 2020. The MADRidge project: Bio-physical coupling around three shallow seamounts in the South West Indian Ocean. *Deep Sea Research Part II: Topical Studies in Oceanography, Bio-Physical coupling around three shallow seamounts in the South Western Indian Ocean, with regional comparisons based on modelling, remote sensing and observational studies* 176, 104813. <https://doi.org/10.1016/j.dsr2.2020.104813>
- Robertson, C.M., Demopoulos, A.W.J., Bourque, J.R., Mienis, F., Duineveld, G.C.A., Lavaleye, M.S.S., Koivisto, R.K.K., Brooke, S.D., Ross, S.W., Rhode, M., Davies, A.J., 2020. Submarine canyons influence macrofaunal diversity and density patterns in the deep-sea benthos. *Deep Sea Research Part I: Oceanographic Research Papers* 159, 103249. <https://doi.org/10.1016/j.dsr.2020.103249>
- Robison, B.H., 2009. Conservation of Deep Pelagic Biodiversity. *Conservation Biology* 23, 847–858. <https://doi.org/10.1111/j.1523-1739.2009.01219.x>

Rodrigues, N., Sharma, R., Nagender Nath, B., 2001. Impact of benthic disturbance on megafauna in Central Indian Basin. Deep Sea Research Part II: Topical Studies in Oceanography, Indian Deep-sea Environment Experiment (INDEX) 48, 3411–3426. [https://doi.org/10.1016/S0967-0645\(01\)00049-2](https://doi.org/10.1016/S0967-0645(01)00049-2)

Rodriguez, M., 2020. The Amirante Ridge and Trench System in the Indian Ocean: the southern termination of the NW Indian subduction. Comptes Rendus Geosciences 352, 235–245. <https://doi.org/10.5802/crgeos.40>

Rogacheva, A., Cross, I.A., Billett, D.S.M., 2009. Psychropotid holothurians (Echinodermata: Holothuroidea: Elasipodida) collected at abyssal depths from around the Crozet Plateau in the Southern Indian Ocean. Zootaxa 2096, 460–478. <https://doi.org/10.11646/zootaxa.2096.1.28>

Rogers, A., 2012. An ecosystem approach to management of seamounts in the Southern Indian Ocean : volume 1 : overview of seamount ecosystems and biodiversity. IUCN.

Rogers, A.D., 2018. Chapter Four - The Biology of Seamounts: 25 Years on, in: Sheppard, C. (Ed.), Advances in Marine Biology. Academic Press, pp. 137–224. <https://doi.org/10.1016/bs.amb.2018.06.001>

Rogers, A.D., 2017. Pelagic ecology of the South West Indian Ocean Ridge seamounts: Introduction and overview. Deep Sea Research Part II: Topical Studies in Oceanography, The Pelagic Ecology of Seamounts of the South West Indian Ocean 136, 1–4. <https://doi.org/10.1016/j.dsr2.2016.12.009>

Rogers, A.D., 2000. The role of the oceanic oxygen minima in generating biodiversity in the deep sea. Deep Sea Research Part II: Topical Studies in Oceanography 47, 119–148. [https://doi.org/10.1016/S0967-0645\(99\)00107-1](https://doi.org/10.1016/S0967-0645(99)00107-1)

Rogers, A.D., Alvheim, O., Bemanaja, E., Benivary, D., Boersch-Supan, P., Bornman, T.G., Cedras, R., Du Plessis, N., Gotheil, S., Høines, A., Kemp, K., Kristiansen, J., Letessier, T., Mangar, V., Mazungula, N., Mørk, T., Pinet, P., Pollard, R., Read, J., Sonnekus, T., 2017. Pelagic communities of the South West Indian Ocean seamounts: R/V Dr Fridtjof Nansen Cruise 2009-410. Deep Sea Research Part II: Topical Studies in Oceanography, The Pelagic Ecology of Seamounts of the South West Indian Ocean 136, 5–35. <https://doi.org/10.1016/j.dsr2.2016.12.010>

Rogers, A.D., Tyler, P.A., Connelly, D.P., Copley, J.T., James, R., Larter, R.D., Linse, K., Mills, R.A., Garabato, A.N., Pancost, R.D., Pearce, D.A., Polunin, N.V.C., German, C.R., Shank, T., Boersch-Supan, P.H., Alker, B.J., Aquilina, A., Bennett, S.A., Clarke, A., Dinley, R.J.J., Graham, A.G.C., Green, D.R.H., Hawkes, J.A., Hepburn, L., Hilario, A., Huvenne, V.A.I., Marsh, L., Ramirez-Llodra, E., Reid, W.D.K., Roterman, C.N., Sweeting, C.J., Thatje, S., Zwirgmaier, K., 2012. The Discovery of New Deep-Sea Hydrothermal Vent Communities in the Southern Ocean and Implications for Biogeography. PLOS Biology 10, e1001234. <https://doi.org/10.1371/journal.pbio.1001234>

Rogers, E.A.D., Taylor, M.L., n.d. Benthic biodiversity of seamounts in the southwest Indian Ocean, Cruise report – R/V James Cook 066.

Rogers, J., 1995. A comparative study of manganese nodules off southern Africa. South African Journal of Geology 98, 208–216. <https://doi.org/10.10520/EJC-92a096b00>

Rogers, P., Ward, T., van Ruth, P., A, W., BD, B., Connell, S., Currie, D., CR, D., Evans, K., Gillanders, B., Goldsworthy, S., Griffin, D., Hardman-Mountford, N., Ivey, A., Kloser, R., Middleton, J., AE, R., A,

- R., Tanner, J., Young, J., 2013. Physical processes, biodiversity and ecology of the Great Australian Bight region: a literature review.
- Romanov, E.V., 2003. Summary and Review of Soviet and Ukrainian Scientific and Commercial Fishing Operations [WWW Document]. URL <https://www.fao.org/3/y4884e/y4884e00.htm> (accessed 3.12.24).
- Ross, A., Williams, A., Talukder, A., Parr, J., Trefry, C., Kempton, R., Stalvies, C., Althaus, F., Kulpecz, A., Schaefer, C., Rarey, P., 2017. Building the regional understanding of the deep-water geology and benthic ecology of the Great Australian Bight. *The APPEA Journal* 57, 798–805. <https://doi.org/10.1071/AJ16212>
- Ross, A., Williams, A., Talukder, A., Parr, J., Trefry, C., Kempton, R., Stalvies, C., Schinteie, R., Ezzy, A., Kulpecz, A., Schaefer, C., Rarey, P., 2018. Insights into the Great Australian Bight gained through marine geology and benthic ecology studies. *The APPEA Journal* 58, 845–851. <https://doi.org/10.1071/AJ17240>
- Roterman, C.N., Lee, W.-K., Liu, X., Lin, R., Li, X., Won, Y.-J., 2018. A new yeti crab phylogeny: Vent origins with indications of regional extinction in the East Pacific. *PLOS ONE* 13, e0194696. <https://doi.org/10.1371/journal.pone.0194696>
- Roxy, M.K., Ritika, K., Terray, P., Masson, S., 2014. The Curious Case of Indian Ocean Warming. *Journal of Climate* 27, 8501–8509. <https://doi.org/10.1175/JCLI-D-14-00471.1>
- Ryther, J.H., 1963. International Indian Ocean Expedition. *AIBS Bulletin* 13, 48–51. <https://doi.org/10.2307/1292962>
- Saji, N.H., Goswami, B.N., Vinayachandran, P.N., Yamagata, T., 1999. A dipole mode in the tropical Indian Ocean. *Nature* 401, 360–363. <https://doi.org/10.1038/43854>
- Sangodkar, N., Gonsalves, M.J., Nazareth, D.R., 2023. Macrofaunal Distribution, Diversity, and Its Ecological Interaction at the Cold Seep Site of Krishna-Godavari Basin, East Coast of India. *Microb Ecol* 85, 61–75. <https://doi.org/10.1007/s00248-021-01942-1>
- Saraswat, R., Nanajkar, M., Damare, S.R., Khare, N., Lei, Y., 2022. Editorial: Benthic Biodiversity of the Indian Ocean. *Frontiers in Marine Science* 9.
- Saunders, B.J., Galaiduk, R., Inostroza, K., Myers, E.M.V., Goetze, J.S., Westera, M., Twomey, L., McCorry, D., Harvey, E.S., 2021. Quantifying Patterns in Fish Assemblages and Habitat Use Along a Deep Submarine Canyon-Valley Feature Using a Remotely Operated Vehicle. *Frontiers in Marine Science* 8.
- Sauter, D., Cannat, M., 2010. The ultraslow spreading Southwest Indian Ridge, in: Rona, P.A., Devey, C.W., Dymont, J., Murton, B.J. (Eds.), *Geophysical Monograph Series*. American Geophysical Union, Washington, D. C., pp. 153–173. <https://doi.org/10.1029/2008GM000843>
- Sautya, S., Gaikwad, S., Khokher, S., Pradhan, U.K., Chatterjee, S., Choudhury, A., Sahu, B., Attri, S., 2021. Distribution Pattern of the Benthic Meiofaunal Community Along the Depth Gradient of the Western Indian Continental Margin, Including the OMZ and Abyssal Plain. *Frontiers in Marine Science* 8.
- Sautya, S., Ingole, B., Jones, D.O.B., Ray, D., Kameshraj, K.A., 2017. First quantitative exploration of benthic megafaunal assemblages on the mid-oceanic ridge system of the Carlsberg Ridge, Indian Ocean. *Journal of the Marine Biological Association of the United Kingdom* 97, 409–417. <https://doi.org/10.1017/S0025315416000515>

- Sautya, S., Ingole, B., Ray, D., Stöhr, S., Samudrala, K., Raju, K.A.K., Mudholkar, A., 2011a. Megafaunal Community Structure of Andaman Seamounts Including the Back-Arc Basin – A Quantitative Exploration from the Indian Ocean. *PLOS ONE* 6, e16162. <https://doi.org/10.1371/journal.pone.0016162>
- Sautya, S., Tabachnick, K.R., Ingole, B., 2011b. A new genus and species of deep-sea glass sponge (Porifera, Hexactinellida, Aulocalycidae) from the Indian Ocean. *Zookeys* 13–21. <https://doi.org/10.3897/zookeys.136.1626>
- Sayers, J., Borissova, I., Ramsay, D., Symonds, P.A., 2002. Geological framework of the Wallaby Plateau and adjacent areas. *Geoscience Australia* 1–133.
- Scheirer, D.S., Baker, E.T., Johnson, K.T.M., 1998. Detection of hydrothermal plumes along the Southeast Indian Ridge near the Amsterdam-St. Paul Plateau. *Geophysical Research Letters* 25, 97–100. <https://doi.org/10.1029/97GL03443>
- Schettino, A., Scotese, C.R., 2005. Apparent polar wander paths for the major continents (200 Ma to the present day): a palaeomagnetic reference frame for global plate tectonic reconstructions. *Geophysical Journal International* 163, 727–759. <https://doi.org/10.1111/j.1365-246X.2005.02638.x>
- Schneider, N., 1998. The Indonesian Throughflow and the Global Climate System. *Journal of Climate* 11, 676–689. [https://doi.org/10.1175/1520-0442\(1998\)011<0676:TITATG>2.0.CO;2](https://doi.org/10.1175/1520-0442(1998)011<0676:TITATG>2.0.CO;2)
- Schott, F.A., Xie, S.-P., McCreary Jr., J.P., 2009. Indian Ocean circulation and climate variability. *Reviews of Geophysics* 47. <https://doi.org/10.1029/2007RG000245>
- SCLATER, J.G., FISHER, R.L., 1974. Evolution of the East: Central Indian Ocean, with Emphasis on the Tectonic Setting of the Ninetyeast Ridge. *GSA Bulletin* 85, 683–702. [https://doi.org/10.1130/0016-7606\(1974\)85<683:EOTECEI>2.0.CO;2](https://doi.org/10.1130/0016-7606(1974)85<683:EOTECEI>2.0.CO;2)
- Scotese, C.R., 2021. An Atlas of Phanerozoic Paleogeographic Maps: The Seas Come In and the Seas Go Out. *Annual Review of Earth and Planetary Sciences* 49, 679–728. <https://doi.org/10.1146/annurev-earth-081320-064052>
- Scotese, C.R., Gahagan, L.M., Larson, R.L., 1988. Plate tectonic reconstructions of the Cretaceous and Cenozoic ocean basins. *Tectonophysics*, 8th Geodynamics Symposium, “Mesozoic and Cenozoic Plate Reconstructions” 155, 27–48. [https://doi.org/10.1016/0040-1951\(88\)90259-4](https://doi.org/10.1016/0040-1951(88)90259-4)
- Scott, R.B., Goff, J.A., Naveira Garabato, A.C., Nurser, A.J.G., 2011. Global rate and spectral characteristics of internal gravity wave generation by geostrophic flow over topography. *Journal of Geophysical Research: Oceans* 116. <https://doi.org/10.1029/2011JC007005>
- Sempéré, J.-C., Cochran, J.R., 1997. The Southeast Indian Ridge between 88°E and 118°E: Variations in crustal accretion at constant spreading rate. *Journal of Geophysical Research: Solid Earth* 102, 15489–15505. <https://doi.org/10.1029/97JB00171>
- Sempéré, J.-C., Palmer, J., Christie, D.M., Morgan, J.P., Shor, A.N., 1991. Australian-Antarctic discordance. *Geology* 19, 429–432. [https://doi.org/10.1130/0091-7613\(1991\)019<0429:AAD>2.3.CO;2](https://doi.org/10.1130/0091-7613(1991)019<0429:AAD>2.3.CO;2)
- Shotton, R., 2006. Management of demersal fisheries resources of the southern Indian Ocean., *FAO Fisheries Circular*. FAO, Rome, Italy.

- Sibuet, M., Olu, K., 1998. Biogeography, biodiversity and fluid dependence of deep-sea cold-seep communities at active and passive margins. *Deep Sea Research Part II: Topical Studies in Oceanography* 45, 517–567. [https://doi.org/10.1016/S0967-0645\(97\)00074-X](https://doi.org/10.1016/S0967-0645(97)00074-X)
- SIODFA Secretariat, 2016. Southern Indian Ocean Deepwater Fisheries Association (SIODFA) Benthic Protected Areas in the Southern Indian Ocean (No. Technical Report XVII 16/02).
- SIOFA, 2023. Active Conservation and Management Measures (CMMs) of the Southern Indian Ocean Fisheries Agreement. Southern Indian Ocean Fisheries Agreement.
- SIOFA Secretariat, 2023. SIOFA Ecosystem Summary 2023. Southern Indian Ocean Fisheries Agreement (SIOFA).
- SIOFA Secretariat, 2023. SIOFA-Fisheries-Overview-2023_redacted.pdf.
- Sloyan, B.M., 2006. Antarctic bottom and lower circumpolar deep water circulation in the eastern Indian Ocean. *Journal of Geophysical Research: Oceans* 111. <https://doi.org/10.1029/2005JC003011>
- Smallwood, B.J., Wolff, G.A., Bett, B.J., Smith, C.R., Hoover, D., Gage, J.D., Patience, A., 1999. Megafauna Can Control the Quality of Organic Matter in Marine Sediments. *Naturwissenschaften* 86, 320–324. <https://doi.org/10.1007/s001140050624>
- Smith, C.R., Baco, A.R., 2003. ECOLOGY OF WHALE FALLS AT THE DEEP-SEA FLOOR. *Oceanography and Marine Biology: An Annual Review* 41, 311–354.
- Smith, C.R., Glover, A.G., Treude, T., Higgs, N.D., Amon, D.J., 2015. Whale-Fall Ecosystems: Recent Insights into Ecology, Paleoecology, and Evolution. *Annual Review of Marine Science* 7, 571–596. <https://doi.org/10.1146/annurev-marine-010213-135144>
- Smith, D.C., Begg, G.A., Lukatelich, R., Anson, T., Baghurst, B., Ham, J., Lapidge, S., Lewis, R., Ormandy, P., Smith, R., Ward, T., 2018. A whole of systems approach to improved understanding of the environmental, economic and social values of a frontier marine oil and gas field: Establishment, success factors and lessons learnt. *Deep Sea Research Part II: Topical Studies in Oceanography, Great Australian Bight Research Program - a whole of system investigation* 157–158, 3–10. <https://doi.org/10.1016/j.dsr2.2018.09.009>
- Smith Jr., K.L., Huffard, C.L., Ruhl, H.A., 2020. Thirty-year time series study at a station in the abyssal NE Pacific: An introduction. *Deep Sea Research Part II: Topical Studies in Oceanography, Thirty-year time-series study in the abyssal NE Pacific* 173, 104764. <https://doi.org/10.1016/j.dsr2.2020.104764>
- Smith Jr., K.L., Sherman, A.D., McGill, P.R., Henthorn, R.G., Ferreira, J., Huffard, C.L., 2017. Evolution of Monitoring an Abyssal Time-Series Station in the Northeast Pacific Over 28 Years. *Oceanography* 30, 72–81.
- Snelgrove, P., Smith, C., 2002. A riot of species in an environmental calm: The paradox of the species-rich deep-sea floor. *Oceanography and marine biology* 40.
- Soltwedel, T., Bauerfeind, E., Bergmann, M., Budaeva, N., Hoste, E., Jaeckisch, N., von Juterzenka, K., Matthiessen, J., Mokievsky, V., Nöthig, E.-M., Quéric, N.-V., Sablotny, B., Sauter, E., Schewe, I., Urban-Malinga, B., Wegner, J., Wlodarska-Kowalczyk, M., Klages, M., 2005. HAUSGARTEN: Multidisciplinary investigations at a deep-sea, long-term observatory in the Arctic Ocean. *Oceanography*.

- Sonnekus, M.J., Bornman, T.G., Campbell, E.E., 2017. Phytoplankton and nutrient dynamics of six South West Indian Ocean seamounts. *Deep Sea Research Part II: Topical Studies in Oceanography, The Pelagic Ecology of Seamounts of the South West Indian Ocean* 136, 59–72. <https://doi.org/10.1016/j.dsr2.2016.12.008>
- Sreejith, K.M., Unnikrishnan, P., Radhakrishna, M., 2019. Isostasy and crustal structure of the Chagos–Laccadive Ridge, Western Indian Ocean: Geodynamic implications. *J Earth Syst Sci* 128, 157. <https://doi.org/10.1007/s12040-019-1161-2>
- Stephens, W.E., Storey, M., Donaldson, C.H., Ellam, R.M., Lelikov, E., Tararin, G., Garbe-Schoenberg, C., 2009. Age and origin of the Amirante ridge-trench structure, western Indian Ocean 2009, T23A-1885.
- Stewart, H.A., Jamieson, A.J., 2019. The five deeps: The location and depth of the deepest place in each of the world's oceans. *Earth-Science Reviews* 197, 102896. <https://doi.org/10.1016/j.earscirev.2019.102896>
- Stewart, H.A., Jamieson, A.J., 2018. Habitat heterogeneity of hadal trenches: Considerations and implications for future studies. *Progress in Oceanography* 161, 47–65. <https://doi.org/10.1016/j.pocean.2018.01.007>
- Stocks, K.I., Clark, M.R., Rowden, A.A., Consalvey, M., Schlacher, T.A., 2012. CenSeam, an International Program on Seamounts within the Census of Marine Life: Achievements and Lessons Learned. *PLOS ONE* 7, e32031. <https://doi.org/10.1371/journal.pone.0032031>
- Stöhr, S., Sautya, S., Ingole, B., 2012. Brittle stars (Echinodermata: Ophiuroidea) from seamounts in the Andaman Sea (Indian Ocean): first account, with descriptions of new species. *Journal of the Marine Biological Association of the United Kingdom* 92, 1195–1208. <https://doi.org/10.1017/S0025315412000240>
- Strutton, P.G., Coles, V.J., Hood, R.R., Matear, R.J., McPhaden, M.J., Phillips, H.E., 2015. Biogeochemical variability in the central equatorial Indian Ocean during the monsoon transition. *Biogeosciences* 12, 2367–2382. <https://doi.org/10.5194/bg-12-2367-2015>
- Stuart, C.T., Arbizu, P.M., Smith, C.R., Molodtsova, T., Brandt, A., Etter, R.J., Escobar-Briones, E., Fabri, M.-C., Rex, M.A., 2008. CeDAMar global database of abyssal biological sampling. *Aquatic Biology* 4, 143–145. <https://doi.org/10.3354/ab00097>
- Subrahmanyam, V., Krishna, K.S., Ramana, M.V., Murthy, K.S.R., 2008. Marine geophysical investigations across the submarine canyon (Swatch-of-No-Ground), northern Bay of Bengal. *Current Science* 94, 507–513.
- Sun, J., Zhou, Y., Chen, C., Kwan, Y.H., Sun, Y., Wang, X., Yang, L., Zhang, R., Wei, T., Yang, Y., Qu, L., Sun, C., Qian, P.-Y., 2020. Nearest vent, dearest friend: biodiversity of Tiancheng vent field reveals cross-ridge similarities in the Indian Ocean. *Royal Society Open Science* 7, 200110. <https://doi.org/10.1098/rsos.200110>
- Suo, Y., Li, S., Li, X., Guo, L., Wang, Y., 2016. Crustal thickness anomalies in the Indian Ocean inferred from gravity analysis. *Geological Journal* 51, 634–643. <https://doi.org/10.1002/gj.2786>
- Suokhrie, T., Saraswat, R., Nigam, R., 2020. Lack of denitrification causes a difference in benthic foraminifera living in the oxygen deficient zones of the Bay of Bengal and the Arabian Sea. *Marine Pollution Bulletin* 153, 110992. <https://doi.org/10.1016/j.marpolbul.2020.110992>

- Swan, J.A., Jamieson, A.J., Linley, T.D., Yancey, P.H., 2021. Worldwide distribution and depth limits of decapod crustaceans (Penaeoidea, Oplophoroidea) across the abyssal-hadal transition zone of eleven subduction trenches and five additional deep-sea features. *Journal of Crustacean Biology* 41, ruaa102. <https://doi.org/10.1093/jcabiol/ruaa102>
- Swanborn, D.J.B., Huvenne, V.A.I., Malpas, T., Pittman, S.J., Rogers, A.D., Taylor, M.L., Woodall, L.C., 2023a. Seamount seascape composition and configuration shape Southwest Indian Ridge fish assemblages. *Deep Sea Research Part I: Oceanographic Research Papers* 191, 103921. <https://doi.org/10.1016/j.dsr.2022.103921>
- Swanborn, D.J.B., Huvenne, V.A.I., Pittman, S.J., Rogers, A.D., Taylor, M.L., Woodall, L.C., 2023b. Mapping, quantifying and comparing seascape heterogeneity of Southwest Indian Ridge seamounts. *Landsc Ecol* 38, 185–203. <https://doi.org/10.1007/s10980-022-01541-6>
- Symonds, P.A., Planke, S., Frey, O., Skogseid, J., 1998. Volcanic Evolution of the Western Australian Continental Margin and its Implications for Basin Development, in: *The Sedimentary Basins of Western Australia II*. Petroleum Exploration Society of Australia (PESA), pp. 33–55.
- Talley, L.D., 2013. Hydrographic Atlas of the World Ocean Circulation Experiment (WOCE) Volume 4: Indian Ocean. International WOCE Project Office, Southampton UK.
- Talley, L.D., Pickard, G.L., Emery, W.J., Swift, J.H., 2011. Indian Ocean, in: *Descriptive Physical Oceanography*. Elsevier, pp. 363–399. <https://doi.org/10.1016/B978-0-7506-4552-2.10011-3>
- Talma, S., 2022. Understanding local deep-sea knowledge and capacity within the Indian Ocean. *Global Deep-Sea Capacity Assessment*. <https://doi.org/10.21428/cbd17b20.4c7a9f87>
- Tao, C., Wu, G., Deng, X., Qiu, Z., Han, C., Long, Y., 2013. New discovery of seafloor hydrothermal activity on the Indian Ocean Carlsberg Ridge and Southern North Atlantic Ridge—progress during the 26th Chinese COMRA cruise. *Acta Oceanol. Sin.* 32, 85–88. <https://doi.org/10.1007/s13131-013-0345-x>
- Taviani, M., Foglini, F., Castellan, G., Montagna, P., McCulloch, M.T., Trotter, J.A., 2023. First assessment of anthropogenic impacts in submarine canyon systems off southwestern Australia. *Science of The Total Environment* 857, 159243. <https://doi.org/10.1016/j.scitotenv.2022.159243>
- Taylor, M.L., Gwinnett, C., Robinson, L.F., Woodall, L.C., 2016. Plastic microfibre ingestion by deep-sea organisms. *Sci Rep* 6, 33997. <https://doi.org/10.1038/srep33997>
- Taylor, M.L., Rogers, A.D., 2017. Primnoidae (Cnidaria: Octocorallia) of the SW Indian Ocean: new species, genus revisions and systematics. *Zoological Journal of the Linnean Society* 181, 70–97. <https://doi.org/10.1093/zoolinlean/zlx003>
- Thaler, A.D., Amon, D., 2019. 262 Voyages Beneath the Sea: a global assessment of macro- and megafaunal biodiversity and research effort at deep-sea hydrothermal vents. *PeerJ* 7, e7397. <https://doi.org/10.7717/peerj.7397>
- Thomas, E.A., Böhm, M., Pollock, C., Chen, C., Seddon, M., Sigwart, J.D., 2022a. Assessing the extinction risk of insular, understudied marine species. *Conservation Biology* 36, e13854. <https://doi.org/10.1111/cobi.13854>
- Thomas, E.A., Liu, R., Amon, D., Copley, J.T., Glover, A.G., Helyar, S.J., Olu, K., Wiklund, H., Zhang, H., Sigwart, J.D., 2020. Chiridota heheva—the cosmopolitan holothurian. *Mar. Biodivers.* 50, 110. <https://doi.org/10.1007/s12526-020-01128-x>

- Thomas, E.A., Molloy, A., Hanson, N.B., Böhm, M., Seddon, M., Sigwart, J.D., 2021. A Global Red List for Hydrothermal Vent Molluscs. *Frontiers in Marine Science* 8.
- Thomas, E.A., Sigwart, J.D., Helyar, S.J., 2022b. New evidence for a cosmopolitan holothurian species at deep-sea reducing environments. *Mar. Biodivers.* 52, 63. <https://doi.org/10.1007/s12526-022-01298-w>
- Thomassin, B., 1977. BENTHEDI I cruise, Le Suroît R/V. <https://doi.org/10.17600/77003111>
- Thompson, K.F., Miller, K.A., Currie, D., Johnston, P., Santillo, D., 2018. Seabed Mining and Approaches to Governance of the Deep Seabed. *Front. Mar. Sci.* 5. <https://doi.org/10.3389/fmars.2018.00480>
- Thums, M., C. Ferreira, L., Jenner, C., Jenner, M., Harris, D., Davenport, A., Andrews-Goff, V., Double, M., Möller, L., Attard, C.R.M., Bilgmann, K., G. Thomson, P., McCauley, R., 2022. Pygmy blue whale movement, distribution and important areas in the Eastern Indian Ocean. *Global Ecology and Conservation* 35, e02054. <https://doi.org/10.1016/j.gecco.2022.e02054>
- Thurston, M.H., Bett, B.J., Rice, A.L., Jackson, P.A.B., 1994. Variations in the invertebrate abyssal megafauna in the North Atlantic Ocean. *Deep Sea Research Part I: Oceanographic Research Papers* 41, 1321–1348. [https://doi.org/10.1016/0967-0637\(94\)90100-7](https://doi.org/10.1016/0967-0637(94)90100-7)
- Thushara, V., Vinayachandran, P.N., 2016. Formation of summer phytoplankton bloom in the northwestern Bay of Bengal in a coupled physical-ecosystem model. *Journal of Geophysical Research: Oceans* 121, 8535–8550. <https://doi.org/10.1002/2016JC011987>
- Toole, J.M., Warren, B.A., 1993. A hydrographic section across the subtropical South Indian Ocean. *Deep Sea Research Part I: Oceanographic Research Papers* 40, 1973–2019. [https://doi.org/10.1016/0967-0637\(93\)90042-2](https://doi.org/10.1016/0967-0637(93)90042-2)
- Treude, T., Janßen, F., Queisser, W., Witte, U., 2002. Metabolism and decompression tolerance of scavenging lysianassoid deep-sea amphipods. *Deep Sea Research Part I: Oceanographic Research Papers* 49, 1281–1289. [https://doi.org/10.1016/S0967-0637\(02\)00023-7](https://doi.org/10.1016/S0967-0637(02)00023-7)
- Trotter, J., Montagna, P., Taviani, M., Sadekov, A., Skrzypek, G., Mazzoli, C., Remia, A., Hosie, A., Hara, A., Pattiaratchi, C., McCulloch, M., 2021. SCHMIDT OCEAN INSTITUTE CRUISE FK200126.
- Trotter, Julie A., McCulloch, M.T., D’Olivo, J.P., Scott, P., Tisnérat-Laborde, N., Taviani, M., Montagna, P., 2022. Deep-water coral records of glacial and recent ocean-atmosphere dynamics from the Perth Canyon in the southeast Indian Ocean. *Quaternary Science Advances* 6, 100052. <https://doi.org/10.1016/j.qsa.2022.100052>
- Trotter, J.A., Pattiaratchi, C., Montagna, P., Taviani, M., Falter, J., Thresher, R., Hosie, A., Haig, D., Foglini, F., Hua, Q., McCulloch, M.T., 2019. First ROV Exploration of the Perth Canyon: Canyon Setting, Faunal Observations, and Anthropogenic Impacts. *Frontiers in Marine Science* 6.
- Trotter, Julie A, Taviani, M., Foglini, F., Sadekov, A., Skrzypek, G., Mazzoli, C., Remia, A., Santodomingo, N., Castellan, G., McCulloch, M., Pattiaratchi, C., Montagna, P., 2022. Unveiling deep-sea habitats of the Southern Ocean-facing submarine canyons of southwestern Australia. *Progress in Oceanography* 209, 102904. <https://doi.org/10.1016/j.pocean.2022.102904>
- Turnewitsch, R., Witte, U., Graf, G., 2000. Bioturbation in the abyssal Arabian Sea: influence of fauna and food supply. *Deep Sea Research Part II: Topical Studies in Oceanography* 47, 2877–2911. [https://doi.org/10.1016/S0967-0645\(00\)00052-7](https://doi.org/10.1016/S0967-0645(00)00052-7)

- Urban Jr., E.R., 2020. Outcomes of the U.S. Program in Biology of the International Indian Ocean Expedition. Deep Sea Research Part II: Topical Studies in Oceanography, The Second International Indian Ocean Expedition (IIOE-2): Motivating New Exploration in a Poorly Understood Ocean Basin Volume 3) 179, 104780. <https://doi.org/10.1016/j.dsr2.2020.104780>
- van der Most, N., Qian, P.-Y., Gao, Y., Gollner, S., 2023. Active hydrothermal vent ecosystems in the Indian Ocean are in need of protection. *Frontiers in Marine Science* 9.
- Van Dover, C.L., 2019. Inactive Sulfide Ecosystems in the Deep Sea: A Review. *Frontiers in Marine Science* 6.
- Van Dover, C.L., 2011. Mining seafloor massive sulphides and biodiversity: what is at risk? *ICES Journal of Marine Science* 68, 341–348. <https://doi.org/10.1093/icesjms/fsq086>
- Van Dover, C.L., 2000. *The Ecology of Deep-Sea Hydrothermal Vents*. Princeton University Press. <https://doi.org/10.1515/9780691239477>
- Van Dover, C.L., Humphris, S.E., Fornari, D., Cavanaugh, C.M., Collier, R., Goffredi, S.K., Hashimoto, J., Lilley, M.D., Reysenbach, A.L., Shank, T.M., Von Damm, K.L., Banta, A., Gallant, R.M., Götz, D., Green, D., Hall, J., Harmer, T.L., Hurtado, L.A., Johnson, P., McKiness, Z.P., Meredith, C., Olson, E., Pan, I.L., Turnipseed, M., Won, Y., Young, C.R., Vrijenhoek, R.C., 2001. Biogeography and Ecological Setting of Indian Ocean Hydrothermal Vents. *Science* 294, 818–823. <https://doi.org/10.1126/science.1064574>
- van Weering, T.C.E., Helder, W., Schalk, P., 1997. Netherlands Indian Ocean Program 1992-1993: First Results. *Deep-Sea Research II* 44, 1177–1193.
- Vanden Berghe, E., Stocks, K.I., Grassle, J.F., 2010. Data Integration: The Ocean Biogeographic Information System, in: *Life in the World's Oceans*. John Wiley & Sons, Ltd, pp. 331–353. <https://doi.org/10.1002/9781444325508.ch17>
- Vanreusel, A., Hilario, A., Ribeiro, P.A., Menot, L., Arbizu, P.M., 2016. Threatened by mining, polymetallic nodules are required to preserve abyssal epifauna. *Sci Rep* 6, 26808. <https://doi.org/10.1038/srep26808>
- Vardaro, M.F., Bagley, P.M., Bailey, D.M., Bett, B.J., Jones, D.O.B., Milligan, R.J., Priede, I.G., Risien, C.M., Rowe, G.T., Ruhl, H.A., Sangolay, B.B., Smith, K.L., Jr., Walls, A., Clarke, J., 2013. A Southeast Atlantic deep-ocean observatory: first experiences and results. *Limnology and Oceanography: Methods* 11, 304–315. <https://doi.org/10.4319/lom.2013.11.304>
- Venu, S., 2013. Deep-Sea Fish Distribution Along the South-West Region of Indian EEZ, in: Venkataraman, K., Sivaperuman, C., Raghunathan, C. (Eds.), *Ecology and Conservation of Tropical Marine Faunal Communities*. Springer, Berlin, Heidelberg, pp. 261–281. https://doi.org/10.1007/978-3-642-38200-0_17
- Vinayachandran, P.N., Matthews, A.J., Kumar, K.V., Sanchez-Franks, A., Thushara, V., George, J., Vijith, V., Webber, B.G.M., Queste, B.Y., Roy, R., Sarkar, A., Baranowski, D.B., Bhat, G.S., Klingaman, N.P., Peatman, S.C., Parida, C., Heywood, K.J., Hall, R., King, B., Kent, E.C., Nayak, A.A., Neema, C.P., Amol, P., Lotliker, A., Kankonkar, A., Gracias, D.G., Vernekar, S., D'Souza, A.C., Valluvan, G., Pargaonkar, S.M., Dinesh, K., Giddings, J., Joshi, M., 2018. BoBBLE: Ocean–Atmosphere Interaction and Its Impact on the South Asian Monsoon. *Bulletin of the American Meteorological Society* 99, 1569–1587. <https://doi.org/10.1175/BAMS-D-16-0230.1>

- Vipin, P.M., Ravi, R., Jose Fernandez, T., Pradeep, K., Boopendranath, M.R., Remesan, M.P., 2012. Distribution of myctophid resources in the Indian Ocean. *Rev Fish Biol Fisheries* 22, 423–436. <https://doi.org/10.1007/s11160-011-9244-4>
- von Herzen, R.P., Vacquier, V., Hill, M.N., 1997. A discussion concerning the floor of the northwest Indian Ocean - Heat flow and magnetic profiles on the Mid-Indian Ocean Ridge. *Philosophical Transactions of the Royal Society of London. Series A, Mathematical and Physical Sciences* 259, 262–270. <https://doi.org/10.1098/rsta.1966.0011>
- von Rad, U., Berner, U., Delisle, G., Dooze-Rolinski, H., Fechner, N., Linke, P., Lückge, A., Roeser, H.A., Schmaljohann, R., Wiedicke, M., SONNE 122/130 Scientific Parties, Block, M., Damm, V., Erbacher, J., Fritsch, J., Harazim, B., Poggenburg, J., Scheeder, G., Schreckenberger, B., von Mirbach, N., Drews, M., Walter, S., Ali Khan, A., Inam, A., Tahir, M., Tabrez, A.R., Cheema, A.H., Pervaz, M., Ashraf, M., 2000. Gas and fluid venting at the Makran accretionary wedge off Pakistan. *Geo-Marine Letters* 20, 10–19. <https://doi.org/10.1007/s003670000033>
- Wafar, M., Venkataraman, K., Ingole, B., Khan, S.A., LokaBharathi, P., 2011. State of Knowledge of Coastal and Marine Biodiversity of Indian Ocean Countries. *PLOS ONE* 6, e14613. <https://doi.org/10.1371/journal.pone.0014613>
- Wang, Y., Han, X., Zhou, Y., Qiu, Z., Yu, X., Petersen, S., Li, H., Yang, M., Chen, Y., Liu, J., Wu, X., Luo, H., 2021. The Daxi Vent Field: An active mafic-hosted hydrothermal system at a non-transform offset on the slow-spreading Carlsberg Ridge, 6°48'N. *Ore Geology Reviews* 129, 103888. <https://doi.org/10.1016/j.oregeorev.2020.103888>
- Wang, Z., Xu, T., Zhang, Y., Zhou, Y., Liu, Z., Chen, C., Watanabe, H.K., Qiu, J.-W., 2020. Molecular phylogenetic and morphological analyses of the 'monospecific' Hesiolyra (Annelida: Hesionidae) reveal two new species. *Deep Sea Research Part I: Oceanographic Research Papers* 166, 103401. <https://doi.org/10.1016/j.dsr.2020.103401>
- Watanabe, H., Beedessee, G., 2014. Vent Fauna on the Central Indian Ridge, in: Ishibashi, J., Okino, K., Sunamura, M. (Eds.), *Subseafloor Biosphere Linked to Hydrothermal Systems: TAIGA Concept*. Springer Japan, Tokyo, pp. 205–212. https://doi.org/10.1007/978-4-431-54865-2_16
- Watanabe, H.K., Chen, C., Marie, D.P., Takai, K., Fujikura, K., Chan, B.K.K., 2018. Phylogeography of hydrothermal vent stalked barnacles: a new species fills a gap in the Indian Ocean 'dispersal corridor' hypothesis. *Royal Society Open Science* 5, 172408. <https://doi.org/10.1098/rsos.172408>
- Watkins, N.D., Kennett, J.P., 1977. Erosion of deep-sea sediments in the Southern Ocean between longitudes 70°E and 190°E and contrasts in manganese nodule development. *Marine Geology* 23, 103–111. [https://doi.org/10.1016/0025-3227\(77\)90084-6](https://doi.org/10.1016/0025-3227(77)90084-6)
- Watkins, N.D., Kennett, J.P., 1971. Antarctic Bottom Water: Major Change in Velocity during the Late Cenozoic between Australia and Antarctica. *Science* 173, 813–818. <https://doi.org/10.1126/science.173.3999.813>
- Watling, L., Guinotte, J., Clark, M.R., Smith, C.R., 2013. A proposed biogeography of the deep ocean floor. *Progress in Oceanography* 111, 91–112. <https://doi.org/10.1016/j.pocean.2012.11.003>
- Webb, T.J., Berghe, E.V., O'Dor, R., 2010. Biodiversity's Big Wet Secret: The Global Distribution of Marine Biological Records Reveals Chronic Under-Exploration of the Deep Pelagic Ocean. *PLOS ONE* 5, e10223. <https://doi.org/10.1371/journal.pone.0010223>

- Wei, X., Luan, X., Meng, F., Lu, Y., He, H., Qiao, J., Yin, J., Wang, Y., Xue, Y., 2023. Deformation feature and tectonic model of the Timor Trough: New interpretation of the evolution and mechanism of Banda arc-continent collision. *Tectonophysics* 862, 229958. <https://doi.org/10.1016/j.tecto.2023.229958>
- Wessel, P., 2007. Seamount Characteristics, in: *Seamounts: Ecology, Fisheries & Conservation*. John Wiley & Sons, Ltd, pp. 1–25. <https://doi.org/10.1002/9780470691953.ch1>
- Weston, J.N.J., Jamieson, A.J., 2022. The Multi-Ocean Distribution of the Hadal Amphipod, *Hirondellea dubia* Dahl, 1959 (Crustacea, Amphipoda). *Frontiers in Marine Science* 9.
- Weston, J.N.J., Jensen, E.L., Hasoon, M.S.R., Kitson, J.J.N., Stewart, H.A., Jamieson, A.J., 2022. Barriers to gene flow in the deepest ocean ecosystems: Evidence from global population genomics of a cosmopolitan amphipod. *Sci Adv* 8, eabo6672. <https://doi.org/10.1126/sciadv.abo6672>
- Weston, J.N.J., Peart, R.A., Jamieson, A.J., 2020. Amphipods from the Wallaby-Zenith Fracture Zone, Indian Ocean: new genus and two new species identified by integrative taxonomy. *Systematics and Biodiversity* 18, 57–78. <https://doi.org/10.1080/14772000.2020.1729891>
- Weston, J.N.J., Peart, R.A., Stewart, H.A., Ritchie, H., Piertney, S.B., Linley, T.D., Jamieson, A.J., 2021. Scavenging amphipods from the Wallaby-Zenith Fracture Zone: Extending the hadal paradigm beyond subduction trenches. *Mar Biol* 168, 1. <https://doi.org/10.1007/s00227-020-03798-4>
- Wiggert, J.D., Murtugudde, R.G., Christian, J.R., 2006. Annual ecosystem variability in the tropical Indian Ocean: Results of a coupled bio-physical ocean general circulation model. *Deep Sea Research Part II: Topical Studies in Oceanography, The US JGOFS Synthesis and Modeling Project: Phase III* 53, 644–676. <https://doi.org/10.1016/j.dsr2.2006.01.027>
- Wiles, E., Green, A., Watkeys, M., Botes, R., Jokat, W., 2019. Submarine canyons of NW Madagascar: A first geomorphological insight. *Deep Sea Research Part II: Topical Studies in Oceanography, The 2nd International Indian Ocean Expedition (IIOE-2): Motivating New Exploration in a Poorly Understood Ocean Basin (Volume 1)* 161, 5–15. <https://doi.org/10.1016/j.dsr2.2018.06.003>
- Wiles, E., Green, A., Watkeys, M., Jokat, W., 2017. The Zambezi Channel: A new perspective on submarine channel evolution at low latitudes. *Geomorphology* 286, 121–132. <https://doi.org/10.1016/j.geomorph.2017.02.014>
- Wiles, E., Green, A., Watkeys, M., Jokat, W., Krockner, R., 2013. The evolution of the Tugela canyon and submarine fan: A complex interaction between margin erosion and bottom current sweeping, southwest Indian Ocean, South Africa. *Marine and Petroleum Geology* 44, 60–70. <https://doi.org/10.1016/j.marpetgeo.2013.03.012>
- Williams, A., Althaus, F., Dunstan, P.K., Poore, G.C.B., Bax, N.J., Kloser, R.J., McEnnulty, F.R., 2010. Scales of habitat heterogeneity and megabenthos biodiversity on an extensive Australian continental margin (100–1100 m depths). *Marine Ecology* 31, 222–236. <https://doi.org/10.1111/j.1439-0485.2009.00355.x>
- Williams, A., Althaus, F., MacIntosh, H., Loo, M., Gowlett-Holmes, K., Tanner, J.E., Sorokin, S.J., Green, M., 2018a. Characterising the invertebrate megafaunal assemblages of a deep-sea (200–3000 m) frontier region for oil and gas exploration: the Great Australian Bight, Australia. *Deep*

Sea Research Part II: Topical Studies in Oceanography, Great Australian Bight Research Program - a whole of system investigation 157–158, 78–91. <https://doi.org/10.1016/j.dsr2.2018.07.015>

Williams, A., Althaus, F., Pogonoski, J., Osterhage, D., Gomon, M., Graham, K., Appleyard, S.A., Gledhill, D., Bray, D., McMillan, P., Green, M., Doyle, S., Graham, A., Tanner, J., Ross, A., 2018b. Composition, diversity and biogeographic affinities of the deep-sea (200–3000 m) fish assemblage in the Great Australian Bight, Australia. *Deep Sea Research Part II: Topical Studies in Oceanography, Great Australian Bight Research Program - a whole of system investigation* 157–158, 92–105. <https://doi.org/10.1016/j.dsr2.2018.05.005>

Williams, A., Koslow, J.A., Last, P.R., 2001. Diversity, density and community structure of the demersal fish fauna of the continental slope off Western Australia (20 to 35°S). *Marine Ecology Progress Series* 212, 247–263. <https://doi.org/10.3354/meps212247>

Witte, U., 1999. Consumption of large carcasses by scavenger assemblages in the deep Arabian Sea: observations by baited camera. *Marine Ecology Progress Series* 183, 139–147. <https://doi.org/10.3354/meps183139>

Witte, U., 2000. Vertical distribution of metazoan macrofauna within the sediment at four sites with contrasting food supply in the deep Arabian Sea. *Deep Sea Research Part II: Topical Studies in Oceanography* 47, 2979–2997. [https://doi.org/10.1016/S0967-0645\(00\)00055-2](https://doi.org/10.1016/S0967-0645(00)00055-2)

Witte, U., Pfannkuche, O., 2000. High rates of benthic carbon remineralisation in the abyssal Arabian Sea. *Deep Sea Research Part II: Topical Studies in Oceanography* 47, 2785–2804. [https://doi.org/10.1016/S0967-0645\(00\)00049-7](https://doi.org/10.1016/S0967-0645(00)00049-7)

Wohl, E., Iskin, E.P., 2021. Damming the wood falls. *Science Advances* 7, eabj0988. <https://doi.org/10.1126/sciadv.abj0988>

Wolff, G.A., Billett, D.S.M., Bett, B.J., Holtvoeth, J., FitzGeorge-Balfour, T., Fisher, E.H., Cross, I., Shannon, R., Salter, I., Boorman, B., King, N.J., Jamieson, A., Chaillan, F., 2011. The Effects of Natural Iron Fertilisation on Deep-Sea Ecology: The Crozet Plateau, Southern Indian Ocean. *PLOS ONE* 6, e20697. <https://doi.org/10.1371/journal.pone.0020697>

Woodall, L.C., Robinson, L.F., Rogers, A.D., Narayanaswamy, B.E., Paterson, G.L.J., 2015. Deep-sea litter: a comparison of seamounts, banks and a ridge in the Atlantic and Indian Oceans reveals both environmental and anthropogenic factors impact accumulation and composition. *Front. Mar. Sci.* 2. <https://doi.org/10.3389/fmars.2015.00003>


Woodall, L.C., Sanchez-Vidal, A., Canals, M., Paterson, G.L.J., Coppock, R., Sleight, V., Calafat, A., Rogers, A.D., Narayanaswamy, B.E., Thompson, R.C., 2014. The deep sea is a major sink for microplastic debris. *Royal Society Open Science* 1, 140317. <https://doi.org/10.1098/rsos.140317>

Woodall, L.C., Talma, S., Steeds, O., Stefanoudis, P., Jeremie-Muzungaile, M.-M., de Comarmond, A., 2021. Co-development, co-production and co-dissemination of scientific research: a case study to demonstrate mutual benefits. *Biology Letters* 17, 20200699. <https://doi.org/10.1098/rsbl.2020.0699>

Wyrtki, K., 1973. An Equatorial Jet in the Indian Ocean. *Science* 181, 262–264. <https://doi.org/10.1126/science.181.4096.262>

Yesson, C., Clark, M.R., Taylor, M.L., Rogers, A.D., 2011. The global distribution of seamounts based on 30 arc seconds bathymetry data. *Deep Sea Research Part I: Oceanographic Research Papers* 58, 442–453. <https://doi.org/10.1016/j.dsr.2011.02.004>

- Yesson, C., Letessier, T.B., Nimmo-Smith, A., Hosegood, P., Brierley, A.S., Hardouin, M., Proud, R., 2021. Improved bathymetry leads to >4000 new seamount predictions in the global ocean – but beware of phantom seamounts! UCL Open Environment 4. <https://doi.org/10.14324/111.444/ucloe.000030>
- Zang, N., Sprintall, J., Ienny, R., Wang, F., 2021. Seasonality of the Somali Current/Undercurrent system. Deep Sea Research Part II: Topical Studies in Oceanography 191–192, 104953. <https://doi.org/10.1016/j.dsr2.2021.104953>
- Zeitzschel, B., Gerlach, S.A., 2012. The Biology of the Indian Ocean. Springer Science & Business Media.
- Zeppilli, D., Sarrazin, J., Leduc, D., Arbizu, P.M., Fontaneto, D., Fontanier, C., Gooday, A.J., Kristensen, R.M., Ivanenko, V.N., Sørensen, M.V., Vanreusel, A., Thébault, J., Mea, M., Allio, N., Andro, T., Arvigo, A., Castrec, J., Danielo, M., Foulon, V., Fumeron, R., Hermabessiere, L., Hulot, V., James, T., Langonne-Augen, R., Le Bot, T., Long, M., Mahabror, D., Morel, Q., Pantalos, M., Pouplard, E., Raimondeau, L., Rio-Cabello, A., Seite, S., Traisnel, G., Urvoy, K., Van Der Stegen, T., Weyand, M., Fernandes, D., 2015. Is the meiofauna a good indicator for climate change and anthropogenic impacts? Mar Biodiv 45, 505–535. <https://doi.org/10.1007/s12526-015-0359-z>
- Zhou, Y., Chen, C., Sun, Y., Watanabe, H.K., Zhang, R., Wang, C., 2019. Amphisamytha (Annelida: Ampharetidae) from Indian Ocean hydrothermal vents: Biogeographic implications. Deep Sea Research Part I: Oceanographic Research Papers 154, 103148. <https://doi.org/10.1016/j.dsr.2019.103148>
- Zhou, Y., Chen, C., Zhang, D., Wang, Y., Watanabe, H.K., Sun, J., Bissessur, D., Zhang, R., Han, Y., Sun, D., Xu, P., Lu, B., Zhai, H., Han, X., Tao, C., Qiu, Z., Sun, Y., Liu, Z., Qiu, J.-W., Wang, C., 2022. Delineating biogeographic regions in Indian Ocean deep-sea vents and implications for conservation. Diversity and Distributions 28, 2858–2870. <https://doi.org/10.1111/ddi.13535>
- Zhou, Y., Wang, Y., Li, Y., Shen, C., Liu, Z., Wang, C., 2020. First report of Osedax in the Indian Ocean indicative of trans-oceanic dispersal through the Southern Ocean. Mar. Biodivers. 50, 4. <https://doi.org/10.1007/s12526-019-01034-x>
- Zhou, Y., Zhang, D., Zhang, R., Liu, Z., Tao, C., Lu, B., Sun, D., Xu, P., Lin, R., Wang, J., Wang, C., 2018. Characterization of vent fauna at three hydrothermal vent fields on the Southwest Indian Ridge: Implications for biogeography and interannual dynamics on ultraslow-spreading ridges. Deep Sea Research Part I: Oceanographic Research Papers 137, 1–12. <https://doi.org/10.1016/j.dsr.2018.05.001>
- Ziko, L., Adel, M., Malash, M.N., Siam, R., 2019. Insights into Red Sea Brine Pool Specialized Metabolism Gene Clusters Encoding Potential Metabolites for Biotechnological Applications and Extremophile Survival. Marine Drugs 17, 273. <https://doi.org/10.3390/md17050273>



Graphical abstract

Journal Pre-proof

Highlights

1. Over 90% of the Indian Ocean is deeper than 1,000 m but deep sea remains unknown
2. Biological research reviewed for deep-sea ecosystems of the Indian Ocean
3. Historical, socioeconomic, geological, and oceanographic research backgrounds presented
4. Ecosystem knowledge summaries provided for each major seafloor feature
5. Data gaps identified and four outstanding problems presented for future studies